

# FO Model Checking of Interval Graphs<sup>\*</sup>

Robert Ganian<sup>1</sup>, Petr Hliněný<sup>2</sup>, Daniel Král<sup>3</sup>, Jan Obdržálek<sup>2</sup>, Jarett Schwartz<sup>4</sup>, and Jakub Teska<sup>5</sup>

<sup>1</sup> Goethe University, Frankfurt, Germany  
rganian@gmail.com

<sup>2</sup> Masaryk University, Brno, Czech Republic  
{hlineny,obdrzalek}@fi.muni.cz

<sup>3</sup> University of Warwick, Coventry, United Kingdom  
D.Kral@warwick.ac.uk

<sup>4</sup> UC Berkeley, Berkeley, United States  
jarett@cs.berkeley.edu

<sup>5</sup> University of West Bohemia, Pilsen, Czech Republic  
teska@kma.zcu.cz

**Abstract.** We study the computational complexity of the FO model checking problem on interval graphs, i.e., intersection graphs of intervals on the real line. The main positive result is that this problem can be solved in time  $O(n \log n)$  for  $n$ -vertex interval graphs with representations containing only intervals with lengths from a prescribed finite set. We complement this result by showing that the same is not true if the lengths are restricted to any set that is dense in some open subset, e.g., in the set  $(1, 1 + \varepsilon)$ .

**Keywords:** FO model checking; parameterized complexity; interval graph; clique-width

## 1 Introduction

Results on the existence of an efficient algorithm for a class of problems have recently attracted a significant amount of attention. Such results are now referred to as algorithmic meta-theorems, see a recent survey [14]. The most prominent example is a theorem of Courcelle [1] asserting that every MSO property can be model checked in linear time on the class of graphs with bounded tree-width. Another example is a theorem of Courcelle, Makowski and Rotics [2] asserting that the same conclusion holds for graphs with bounded clique-width when quantification is restricted to vertices and their subsets.

In this paper, we focus on more restricted graph properties, specifically those expressible in the first order logic. Clearly, every such property can be tested in

---

<sup>\*</sup> All the authors except for Jarett Schwartz acknowledge support of the Czech Science Foundation under grant P202/11/0196. Jarett Schwartz acknowledges support of the Fulbright and NSF Fellowships.

polynomial time if we allow the degree of the polynomial to depend on the property of interest. But can these properties be tested in so-called *fixed parameter tractable* (FPT [6]) time, i.e., in polynomial time where the degree of the polynomial does not depend on the considered property? The first result in this direction could be that of Seese [20]: Every FO property can be tested in linear time on graphs with bounded maximum degree. A breakthrough result of Frick and Grohe [11] asserts that every FO property can be tested in almost linear time on classes of graphs with locally bounded tree-width. Here, an almost linear algorithm stands for an algorithm running in time  $O(n^{1+\varepsilon})$  for every  $\varepsilon > 0$ . A generalization to graph classes locally excluding a minor (with worse running time) was later obtained by Dawar, Grohe and Kreutzer [4].

Research in this direction so far culminated in establishing that every FO property can be tested in almost linear time on classes of graphs with locally bounded expansion, as shown (independently) by Dawar and Kreutzer [5] (also see [13] for the complete proof), and by Dvořák, Král' and Thomas [7]. The concept of graph classes with bounded expansion has recently been introduced by Nešetřil and Ossona de Mendéz [16,17,18]; examples of such graph classes include classes of graphs with bounded maximum degree or proper minor-closed classes of graphs. A holy grail of this area is establishing the fixed parameter tractability of testing FO properties on nowhere-dense classes of graphs.

In this work, we investigate whether structural properties which do not yield (locally) bounded width parameters could lead to similar results. Specifically, we study the intersection graphs of intervals on the real line, which are also called interval graphs. When we restrict to unit interval graphs, i.e., intersection graphs of intervals with unit lengths, one can easily deduce the existence of a linear time algorithm for testing FO properties from Gaifman's theorem, using the result of Courcelle et al [2] and that of Lozin [15] asserting that every proper hereditary subclass of unit interval graphs, in particular, the class of unit interval graphs with bounded radius, has bounded clique-width. This observation is a starting point for our research presented in this paper.

Let us now give a definition. For a set  $L$  of reals, an interval graph is called an  $L$ -interval graph if it is an intersection graph of intervals with lengths from  $L$ . For example, unit interval graphs are  $\{1\}$ -interval graphs. If  $L$  is a finite set of rationals, then any  $L$ -interval graph with bounded radius has bounded clique-width (see Section 4 for further details). So, FO properties of such graphs can be tested in the fixed parameter way. However, if  $L$  is not a set of rationals, there exist  $L$ -interval graphs with bounded radius and unbounded clique-width, and so the easy argument above does not apply.

Our main algorithmic result says that every FO property can be tested in time  $O(n \log n)$  for  $L$ -interval graphs when  $L$  is any finite set of reals. To prove this result, we employ a well-known relation of FO properties to Ehrenfeucht-Fraïssé games. Specifically, we show using the notion of game trees, which we introduce, that there exists an algorithm transforming an input  $L$ -interval graph to another  $L$ -interval graph that has bounded maximum degree and that satisfies the same properties expressible by FO sentences with bounded quantifier rank.

On the negative side, we show that if  $L$  is an infinite set that is dense in some open set, then  $L$ -interval graphs can be used to model arbitrary graphs. Specifically, we show that  $L$ -interval graphs for such sets  $L$  allow polynomially bounded FO interpretations of all graphs. Consequently, testing FO properties for  $L$ -interval graphs for such sets  $L$  is  $W[2]$ -hard (see Corollary 5.2). In addition, we show that unit interval graphs allow polynomially bounded MSO interpretations of all graphs.

The property of being  $W[2]$ -hard comes from the theory of parameterized complexity [6], and it is equivalent to saying that the considered problem is at least as hard as the  $d$ -dominating set problem, asking for an existence of a dominating set of fixed parameter size  $d$  in a graph. It is known that, unless the Exponential time hypothesis fails,  $W[2]$ -hard problems cannot have polynomial algorithms with the degree a constant independent of the parameter (of the considered FO property in our case).

In Section 2, we introduce the notation and the computational model used in the paper. In the following section, we present an  $O(n \log n)$  algorithm for deciding FO properties of  $L$ -interval graphs for finite sets  $L$ . In Section 4, we present proofs of the facts mentioned above on the clique-width of  $L$ -interval graphs with bounded radius. Finally, we establish FO interpretability of graphs in  $L$ -interval graphs for sets  $L$  which are dense in an open set in Section 5.

## 2 Notation

An *interval graph* is a graph  $G$  such that every vertex  $v$  of  $G$  can be associated with an interval  $J(v) = [\ell(v), r(v))$  such that two vertices  $v$  and  $v'$  of  $G$  are adjacent if and only if  $J(v)$  and  $J(v')$  intersect (it can be easily shown that the considered class of graphs remains the same regardless of whether we consider open, half-open or closed intervals in the definition). We refer to such an assignment of intervals to the vertices of  $G$  as a *representation* of  $G$ . The point  $\ell(v)$  is the *left end point* of the interval  $J(v)$  and  $r(v)$  is its *right end point*.

If  $L$  is a set of reals and  $r(v) - \ell(v) \in L$  for every vertex  $v$ , we say that  $G$  is an  $L$ -interval graph and we say that the representation is an  $L$ -representation of  $G$ . For example, if  $L = \{1\}$ , we speak about unit interval graphs. Finally, if  $r(v) - \ell(v) \in L$  and  $0 \leq \ell(v) \leq r(v) \leq d$  for some real  $d$ , i.e., all intervals are subintervals of  $[0, d)$ , we speak about  $(L, d)$ -interval graphs. Note that if  $G$  is an interval graph of radius  $k$ , then  $G$  is also an  $(L, (2k + 1) \max L)$ -interval graph (we use  $\max L$  and  $\min L$  to denote the maximum and the minimum elements, respectively, of the set  $L$ ).

We now introduce two technical definitions related to manipulating intervals and their lengths. These definitions are needed in the next section. If  $L$  is a set of reals, then  $L^{(k)}$  is the set of all integer linear combinations of numbers from  $L$  with the sum of the absolute values of their coefficients bounded by  $k$ . For instance,  $L^{(0)} = \{0\}$  and  $L^{(1)} = L \cup \{0\}$ . An  $L$ -distance of two intervals  $[a, b)$  and  $[c, d)$  is the smallest  $k$  such that  $c - a \in L^{(k)}$ . If no such  $k$  exists, then the  $L$ -distance of two intervals is defined to be  $\infty$ .

Since we do not restrict our attention to  $L$ -interval graphs where  $L$  is a set of rationals, we should specify the computational model considered. We use the standard RAM model with infinite arithmetic precision and unit cost of all arithmetic operation, but we refrain from trying to exploit the power of this computational model by encoding any data in the numbers we store. In particular, we only store the end points of the intervals in the considered representations of graphs in numerical variables with infinite precision.

## 2.1 Clique-width

We now briefly present the notion of clique-width introduced in [3]. Our results on interval graphs related to this notion are given in Section 4. A  $k$ -labeled graph is a graph with vertices that are assigned integers (called labels) from 1 to  $k$ . The *clique-width* of a graph  $G$  equals the minimum  $k$  such that  $G$  can be obtained from single vertex graphs with label 1 using the following four operations: relabeling all vertices with label  $i$  to  $j$ , adding all edges between the vertices with label  $i$  and the vertices with label  $j$  ( $i$  and  $j$  can be the same), creating a vertex labeled with 1, and taking a disjoint union of graphs obtained using these operations.

## 2.2 First Order Properties

In this subsection, we introduce concepts from logic and model theory which we use. A *first order (FO) sentence* is a formula with no free variables with the usual logical connectives and quantification allowed only over variables for elements (vertices in the case of graphs). A *monadic second order (MSO) sentence* is a formula with no free variables with the usual logical connectives where, unlike in FO sentences, quantification over subsets of elements is allowed. An FO *property* is a property expressible by an FO sentence; similarly, an *MSO property* is a property expressible by an MSO sentence. Finally, the *quantifier rank* of a formula is the maximum number of nested quantifiers in it.

FO sentences are closely related to the so-called Ehrenfeucht-Fraïssé games. The  $d$ -round *Ehrenfeucht-Fraïssé game* is played on two relational structures  $R_1$  and  $R_2$  by two players referred to as the *spoiler* and the *duplicator*. In each of the  $d$  rounds, the spoiler chooses an element in one of the structures and the duplicator chooses an element in the other. Let  $x_i$  be the element of  $R_1$  chosen in the  $i$ -th round and  $y_i$  be the element of  $R_2$ . We say that the duplicator *wins* the game if there is a strategy for the duplicator such that the substructure of  $R_1$  induced by the elements  $x_1, \dots, x_d$  is always isomorphic to the substructure of  $R_2$  induced by the elements  $y_1, \dots, y_d$ , with the isomorphism mapping each  $x_i$  to  $y_i$ .

The following theorem [8,10] relates this notion to FO sentences of quantifier rank at most  $d$ .

**Theorem 2.1.** *Let  $d$  be an integer. The following statements are equivalent for any two structures  $R$  and  $R'$ :*

- The structures  $R$  and  $R'$  satisfy the same FO sentences of quantifier rank at most  $d$ .
- The duplicator wins the  $d$ -round Ehrenfeucht-Fraïssé game for  $R$  and  $R'$ .

We describe possible courses of the  $d$ -round Ehrenfeucht-Fraïssé game on a single relational structure by a rooted tree, which we call a  $d$ -EF-tree. All the leaves of an  $d$ -EF-tree are at depth  $d$  and each of them is associated with a relational structure with elements labeled from 1 to  $d$ . The  $d$ -EF-tree  $\mathcal{T}$  for the game played on a relational structure  $R$  is obtained as follows. The number of children of every internal node of  $\mathcal{T}$  is the number of elements of  $R$  and the edges leaving the internal node to its children are associated in the one-to-one way with the elements of  $R$ . So, every path from the root of  $\mathcal{T}$  to a leaf  $u$  of  $\mathcal{T}$  yields a sequence  $x_1, \dots, x_d$  of the elements of  $R$  (those associated with the edges of that path) and the substructure of  $R$  induced by  $x_1, \dots, x_d$  with  $x_i$  labeled with  $i$  is the one associated with the leaf  $u$ .

A mapping  $f$  from a  $d$ -EF-tree  $\mathcal{T}$  to another  $d$ -EF-tree  $\mathcal{T}'$  is an EF-homomorphism if the following three conditions hold:

1. if  $u$  is a parent of  $v$  in  $\mathcal{T}$ , then  $f(u)$  is a parent of  $f(v)$  in  $\mathcal{T}'$ , and
2. if  $u$  is a leaf of  $\mathcal{T}$ , then  $f(u)$  is a leaf of  $\mathcal{T}'$ , and
3. the structures associated with  $u$  and  $f(u)$  are isomorphic through the bijection given by the labelings.

Two trees  $\mathcal{T}$  and  $\mathcal{T}'$  are *EF-equivalent* if there exist an EF-homomorphism from  $\mathcal{T}$  to  $\mathcal{T}'$  and an EF-homomorphism from  $\mathcal{T}'$  to  $\mathcal{T}$ .

Let  $\mathcal{T}$  and  $\mathcal{T}'$  be the  $d$ -EF-trees for the game played on relational structures  $R$  and  $R'$ , respectively. Suppose that  $\mathcal{T}$  and  $\mathcal{T}'$  are EF-equivalent and let  $f_1$  and  $f_2$  be the EF-homomorphism from  $\mathcal{T}$  to  $\mathcal{T}'$  and  $\mathcal{T}'$  to  $\mathcal{T}$ , respectively. We claim that the duplicator wins the  $d$ -round Ehrenfeucht-Fraïssé game for  $R$  and  $R'$ . Let us describe a possible strategy for the duplicator. We restrict to the case when the spoiler chooses an element  $x_i$  of  $R$  in the  $i$ -th step, assuming elements  $x_1, \dots, x_{i-1}$  in  $R$  and elements  $y_1, \dots, y_{i-1}$  in  $R'$  have been chosen in the previous rounds. Let  $u_0, \dots, u_i$  be the path in  $\mathcal{T}$  corresponding to  $x_1, \dots, x_i$ . The duplicator chooses the element  $y_i$  in  $R'$  that corresponds to the edge  $f_1(u_{i-1})f_1(u_i)$  in  $\mathcal{T}'$ . It can be verified that if the duplicator follows this strategy, then the substructures of  $R$  and  $R'$  induced by  $x_1, \dots, x_d$  and  $y_1, \dots, y_d$ , respectively, are isomorphic.

Let us summarize our findings from the previous paragraph.

**Theorem 2.2.** *Let  $d$  be an integer. If the  $d$ -EF-trees for the game played on two relational structure  $R$  and  $R'$  are EF-equivalent, then the duplicator wins the  $d$ -round Ehrenfeucht-Fraïssé game for  $R$  and  $R'$ .*

The converse implication, i.e., that if the duplicator can win the  $d$ -round Ehrenfeucht-Fraïssé game for  $R$  and  $R'$ , then the  $d$ -EF-trees for the game played on relational structures  $R$  and  $R'$  are EF-equivalent, is also true, but we omit further details as we only need the implication given in Theorem 2.2 in our considerations.

We finish this section with some observations on minimal  $d$ -EF-trees in EF-equivalence classes. Let  $\mathcal{T}$  be a  $d$ -EF-tree. Suppose that an internal node at level  $d - 1$  has two children, which are leaves, associated with the same labeled structure. Observe that deleting one of them yields an EF-equivalent  $d$ -EF-tree. Suppose that we have deleted all such leaves and an internal node at level  $d - 2$  has two children with their subtrees isomorphic (in the usual sense). Again, deleting one of them (together with its subtree) yields an EF-equivalent  $d$ -EF-tree. So, if  $\mathcal{T}'$  is a minimal subtree of  $\mathcal{T}$  that is EF-equivalent to  $\mathcal{T}$  and  $K$  is the number of non-isomorphic  $d$ -labeled structures, then the degree of nodes at depth  $d - 1$  does not exceed  $K$ , those at depth  $d - 2$  does not exceed  $2^K$ , those at depth  $d - 3$  does not exceed  $2^{2^K}$ , etc. We conclude that the size of a minimal subtree of  $\mathcal{T}$  that is EF-equivalent to  $\mathcal{T}$  is bounded by a function of  $d$  and the type of relational structures considered only.

### 3 FO Model Checking

Using Theorems 2.1 and 2.2, we prove the following kernelization result for  $L$ -interval graphs.

**Theorem 3.1.** *For every finite subset  $L$  of reals and every  $d$ , there exists an integer  $K_0$  and an algorithm  $\mathcal{A}$  with the following properties. The input of  $\mathcal{A}$  is an  $L$ -representation of an  $n$ -vertex  $L$ -interval graph  $G$  and  $\mathcal{A}$  outputs in time  $O(n \log n)$  an  $L$ -representation of an induced subgraph  $G'$  of  $G$  such that*

- every unit interval contains at most  $K_0$  left end points of the intervals corresponding to vertices of  $G'$ , and
- $G$  and  $G'$  satisfy the same FO sentences with quantifier rank at most  $d$ .

*Proof.* We first focus on proving the existence of the number  $K_0$  and the subgraph  $G'$  and we postpone the algorithmic considerations to the end of the proof.

As the first step, we show that we can assume that all the left end points are distinct. Choose  $\delta$  to be the minimum distance between distinct end points of intervals in the representation. Suppose that the intervals are sorted by their left end points (resolving ties arbitrarily). Shifting the  $i$ -th interval by  $i\delta/2n$ , for  $i = 1, \dots, n$ , to the right does not change the graph represented by the intervals and all the end points become distinct.

Choose  $\varepsilon$  to be the minimum positive element of  $L^{(2^{d+1})}$ . Fix any real  $a$  and let  $\mathcal{I}$  be the set of all intervals  $[x, x + \varepsilon)$  such that  $x - a \in L^{(2^{d+1})}$ . By the choice of  $\varepsilon$ , the intervals of  $\mathcal{I}$  are disjoint. In addition, the set  $\mathcal{I}$  is finite (since  $L$  is finite). Let  $W$  be the set of vertices  $w$  of  $G$  such that  $\ell(w)$  lies in an interval from  $\mathcal{I}$ , and for such a vertex  $w$ , let  $i(w)$  be the left end point of that interval from  $\mathcal{I}$ . Define a linear order on  $W$  such that  $w \leq w'$  for  $w$  and  $w'$  from  $W$  iff  $\ell(w) - i(w) \leq \ell(w') - i(w')$  and resolve the cases of equality for distinct vertices  $w$  and  $w'$  arbitrarily.

We view  $W$  as a linearly ordered set with elements associated with intervals from  $\mathcal{I}$  (specifically the interval with the left end point at  $i(w)$ ) as well as associated with the lengths of their corresponding intervals in the representation of  $G$ . Let us establish the following claim.

*Claim.* There exists a number  $K$  depending on only  $|\mathcal{I}|$  and  $d$  such that if  $W$  contains more than  $K$  elements associated with the interval  $[a, a + \varepsilon)$ , then there exists an element  $w \in W$  associated with  $[a, a + \varepsilon)$  such that the  $d$ -EF-trees for the game played on  $W$  and  $W \setminus \{w\}$  are EF-equivalent.

Indeed, let  $\mathcal{T}$  be the  $d$ -EF-tree for the game played on  $W$  and let  $\mathcal{T}'$  a minimal subtree of  $\mathcal{T}$  that is EF-equivalent to  $\mathcal{T}$ . Recall that the size of  $\mathcal{T}'$  does not exceed a number  $K$  depending on only  $|\mathcal{I}|$  and  $d$ . If  $W$  contains more than  $K$  elements associated with  $[a, a + \varepsilon)$ , then one of them is not associated with edges that are present in  $\mathcal{T}'$ . We set  $w$  to be this element. This finishes the proof of the claim.

Since the  $d$ -EF-trees for the game played on  $W$  and  $W \setminus \{w\}$  are EF-equivalent, the duplicator wins the  $d$ -round Ehrenfeucht-Fraïssé game for  $W$  and  $W \setminus \{w\}$  by Theorems 2.1 and 2.2.

We now describe a strategy for the duplicator to win the  $d$ -round Ehrenfeucht-Fraïssé game for the graphs  $G$  and  $G \setminus w$ . During the game, some intervals from  $\mathcal{I}$  will be marked as *altered*. At the beginning, the only altered interval is the interval  $[a, a + \varepsilon)$ .

The duplicator strategy in the  $i$ -th round of the game is the following.

- If the spoiler chooses a vertex  $w'$  with  $\ell(w')$  in an interval of  $\mathcal{I}$  at  $L$ -distance at most  $2^{d+1-i}$  from an altered interval, then the duplicator follows its winning strategy for the  $d$ -round Ehrenfeucht-Fraïssé game for  $W$  and  $W \setminus \{w\}$  which gives the vertex to choose in the other graph. In addition, the duplicator marks the interval of  $\mathcal{I}$  that contains  $\ell(w')$  as altered. Note that the interval of the chosen vertex has its left end point in the same interval of  $\mathcal{I}$  as  $w'$ .
- Otherwise, the duplicator chooses the same vertex in the other graph. No new intervals are marked as altered.

It remains to argue that the subgraphs of  $G$  and  $G \setminus w$  obtained in this way are isomorphic. Let  $w_1, \dots, w_d$  be the chosen vertices of  $G$  and  $w'_1, \dots, w'_d$  the chosen vertices of  $G'$ . For brevity, let us refer to vertices corresponding to the intervals with left end points in the altered intervals as to *altered vertices*. If  $w_i$  is not altered, then  $w_i = w'_i$ . If  $w_i$  is altered, then  $\ell(w_i)$  and  $\ell(w'_i)$  are in the same interval  $J \in \mathcal{I}$  and the only intervals that might intersect the intervals corresponding to  $w_i$  and  $w'_i$  differently are those with left end points in the intervals of  $\mathcal{I}$  at  $L$ -distance at most two from  $J$ . However, if some chosen vertices have their left end points in such intervals, then these intervals must also be altered and these chosen vertices are altered. Since we have followed a winning strategy for the duplicator for  $W$  and  $W \setminus \{w\}$  when choosing altered vertices, the subgraphs of  $G$  and  $G \setminus w$  induced by the altered vertices are isomorphic. We

conclude that the subgraphs of  $G$  and  $G \setminus w$  induced by the vertices  $w_1, \dots, w_d$  and  $w'_1, \dots, w'_d$ , respectively, are isomorphic. So, the duplicator wins the game.

Let us summarize our findings. If an interval of length  $\varepsilon$  contains more than  $K$  left end points of intervals in the given  $L$ -representation of  $G$ , then one of the vertices corresponding to these intervals can be removed from  $G$  without changing the set of FO sentences with rank at most  $d$  that are satisfied by  $G$ . So, the statement of the theorem is true with  $K_0$  set to  $K \lceil \varepsilon^{-1} \rceil$ .

It remains to consider the algorithmic aspects of the theorem. The values of  $\varepsilon$  and  $K_0$  are determined by  $L$  and  $d$ . The algorithm sorts the left end points of all the intervals (this requires  $O(n \log n)$  time) and for each of these points computes the distance to the left end of the interval that is  $K$  positions later in the obtained order. If all these distances are at least  $\varepsilon$ , then every interval of length at most  $\varepsilon$  contains at most  $K_0$  left end points of the intervals and the representation is of the desired form.

Otherwise, we choose the smallest of these distances and consider the corresponding interval  $[a, b)$ ,  $b - a < \varepsilon$ , containing at least  $K_0$  left end points of the intervals from the representation. By the choice of this interval, any interval of length  $b - a$  at  $L$ -distance at most  $2^{d+1}$  from  $[a, b]$  contains at most  $K_0 + 1$  left end points of the intervals from the representation. So, the size of the  $d$ -EF-tree for the game played on the vertices  $v$  with  $\ell(v)$  in such intervals is bounded by a function of  $K_0$ ,  $d$  and  $|L|$ . Since this quantity is independent of the input graph, we can identify in constant time a vertex  $w$  with  $\ell(w) \in [a, b)$  whose removal from  $G$  does not change the set of FO sentences with quantifier rank  $d$  satisfied by  $G$ .

We then update the order of the left end points and the at most  $K_0$  computed distances affected by removing  $w$ , and iterate the whole process. Since at each step we alter at most  $K_0$  distances, using a heap to store the computed distances and choose the smallest of them requires  $O(\log n)$  time per vertex removal. So, the running time of the algorithm is bounded by  $O(n \log n)$ .  $\square$

It is possible to think of several strategies to efficiently decide FO properties of  $L$ -interval graphs given Theorem 3.1. We present one of them. Fix an FO sentence  $\Phi$  with quantifier rank  $d$  and apply the algorithm from Theorem 3.1 to get an  $L$ -interval graph and a representation of this graph such that every unit interval contains at most  $K$  left end points of the intervals of the representation. After this preprocessing step, every vertex of the new graph has at most  $K(|L| + 1) \lceil \max L \rceil$  neighbors. In particular, the maximum degree of the new graph is bounded. The result of Seese [20] asserts that every FO property can be decided in linear time for graphs with bounded maximum degree, and so we conclude:

**Theorem 3.2.** *For every finite subset  $L$  of reals and every FO sentence  $\Phi$ , there exists an algorithm running in time  $O(n \log n)$  that decides whether an input  $n$ -vertex  $L$ -interval graph  $G$  given by its  $L$ -representation satisfies  $\Phi$ .*

## 4 Clique-width of Interval Graphs

Unit interval graphs can have unbounded clique-width [12], but Lozin [15] noted that every proper hereditary subclass of unit interval graphs has bounded clique-width. In particular, the class of  $(\{1\}, d)$ -interval graphs has bounded clique-width for every  $d > 0$ . Using Gaifman's theorem, it follows that testing FO properties of unit interval graphs can be performed in linear time if the input graph is given by its  $\{1\}$ -representation with the left end points of the intervals sorted. We provide an easy extension of this, and outline how it can be used to prove the special case of our main result for FO model checking when  $L$  is a finite set of rational numbers

**Lemma 4.1.** *Let  $L$  be a finite set of positive rational numbers. For any  $d > 0$ , the class of  $(L, d)$ -interval graphs has bounded clique-width.*

*Proof.* Let  $a$  be the largest rational number such that every element of  $L$  is an integer multiple of  $a$ . Without loss of generality, we can assume that  $d$  is not a multiple of  $a$  (otherwise, we slightly increase  $d$ ). We show that the clique-width of any  $(L, d)$ -interval graph is at most  $K := \lceil d/a \rceil + 1$ .

Let  $G$  be an  $(L, d)$ -interval graph with vertices  $v_1, \dots, v_n$ . Consider an  $(L, d)$ -representation of  $G$ . Let  $b_i$  be the smallest non-negative real such that  $\ell(v_i) - b_i$  is a multiple of  $a$ . By a slight perturbation, we may assume that all the numbers  $b_i$  are distinct. Without loss of generality, we can also assume that  $0 < b_1 < \dots < b_n < a$ .

We will now proceed in several steps. After the  $i$ -th step, we will have constructed the subgraph of  $G$  induced by the vertices  $v_1, \dots, v_i$  such that the label of the vertex  $v_i$  is  $\lceil \ell(v_i)/a \rceil$ . In the first step, we insert the vertex  $v_1$  with label  $\lceil \ell(v_1)/a \rceil$ . In the  $i$ -th step, we insert the vertex  $v_i$  with label  $K$ , join it by edges to all vertices with labels between  $\lceil \ell(v_i)/a \rceil$  and  $\lceil r(v_i)/a \rceil$ , and relabel it to  $\lceil \ell(v_i)/a \rceil$ . By the choice of  $a$  and the assumption that  $b_1 < \dots < b_n$ , the vertex  $v_i$  is adjacent to its neighbors among  $v_1, \dots, v_{i-1}$ .  $\square$

From Lemma 4.1 and Gaifman's theorem, one can approach the FO model checking problem on  $L$ -interval graphs with  $L$  containing rational numbers only as follows.  $L$ -interval graphs with radius  $d$  are  $(L, (2d+1) \max L)$ -interval graphs. By Gaifman's theorem, every FO model checking instance can be reduced to model checking of basic local FO sentences, i.e., to FO model checking on  $L$ -interval graphs with bounded radius. Since such graphs have bounded clique-width, the latter can be solved in linear time by [2]. Combining this with the covering technique from [11], which can be adapted to run in linear time in the case of  $L$ -interval graphs, we obtain the following.

**Corollary 4.2.** *Let  $L$  be a finite set of positive rational numbers. The FO model checking problem can be solved in linear time on the class of  $L$ -interval graphs if the input graph is given by its  $L$ -representation with the left end points of the intervals sorted.*

However, Corollary 4.2 is just a fortunate special case, since aside of rational lengths one can prove the following.

**Lemma 4.3.** *For any irrational  $q > 0$  there is  $d$  such that the class of  $(\{1, q\}, d)$ -interval graphs has unbounded clique-width.*

*Proof.* This proof is in a sense complementary to that of Lemma 4.1. We may assume  $q > 1$  (otherwise, we rescale and consider the set  $\{1, 1/q\}$ ). So, fix  $L = \{1, q\}$ ,  $d = q + 3$  and an integer  $n$  (to be specified later).

Our task is to construct a  $(\{1, q\}, d)$ -interval representation of a graph  $G$  with large clique-width. Since  $q$  is irrational, for every  $\ell$  we can find  $n$  such that  $L^{(n)} \cap [0, d - q)$  contains more than  $\ell$  points. We actually construct an arbitrarily long sequence  $P = (a_1, a_2, \dots, a_n)$  of such points as follows:  $a_1 = 0, a_2 = 1$ , and for  $i > 2$  set

- $a_i = a_{i-1} + 1$ , provided that  $|a_{i-2} - a_{i-1}| = 1$  and  $a_{i-1} < d - 2$ ,
- $a_i = a_{i-1} - 1$ , provided that  $a_{i-2} - a_{i-1} = q$ , and
- $a_i = a_{i-1} - q$  otherwise (we call this  $a_i$  a  $q$ -element of  $P$ ).

Informally, we are “folding” a long sequence with differences from  $L^{(n)}$  into a bounded length interval, avoiding as much collisions of points as possible.

Let  $\delta > 0$  be such that  $n\delta$  is smaller than the smallest number in  $L^{(n)} \cap (0, d - q)$ . Let us introduce the following shorthand notation: if  $J$  is an interval and  $r$  a real, then  $J + r$  is the interval  $J$  shifted by  $r$  to the right. Similarly, if  $\mathcal{I}$  is a set of intervals, then  $\mathcal{I} + r$  is the set of the intervals from  $\mathcal{I}$  shifted by  $r$  to the right. We define sets of intervals  $\mathcal{U}_1 := \{[i\delta, 1 + i\delta) : i = 0, \dots, n - 1\}$  and  $\mathcal{U}_q := \{[i\delta, q + i\delta) : i = 0, \dots, n - 1\}$ . For further reference we say that intervals  $[i\delta, 1 + i\delta)$  or  $[i\delta, q + i\delta)$  are *at level  $i$* .

For  $i = 1, \dots, n$ , we set  $\mathcal{W}_i = \mathcal{U}_q + a_i$  if  $a_i$  is a  $q$ -element of  $P$ , and  $\mathcal{W}_i = \mathcal{U}_1 + a_i$  otherwise. Then every interval of  $\mathcal{W}_i$  is a subinterval of  $[0, d)$ . Let  $G$  be a graph on  $n^2$  vertices represented by the union of the interval sets  $\mathcal{W}_1 \cup \mathcal{W}_2 \cup \dots \cup \mathcal{W}_n$ . Let  $W_i$ ,  $i = 1, \dots, n$ , be the vertices represented by  $\mathcal{W}_i$ . We claim that the clique-width of  $G$  exceeds any fixed number  $k \in \mathbb{N}$  when  $n$  sufficiently large.

Assume, for a contradiction, that the clique-width of  $G$  is at most  $k$ . We can view the construction of  $G$  as a binary tree and conclude a  $k$ -labeled subgraph  $G_1$  of  $G$  with  $\frac{1}{3}n^2 \leq |V(G_1)| \leq \frac{2}{3}n^2$  appeared during the construction of  $G$ . However, this implies that vertices of  $G_1$  have at most  $k$  different neighborhoods in  $G \setminus V(G_1)$ . We will show that this is not possible (assuming that  $n$  is large).

For  $2 \leq i \leq n$ , vertices  $x \in W_{i-1}$  and  $y \in W_i$  are *mates* if they are represented by copies of the same-level intervals from  $\mathcal{U}_1$  or  $\mathcal{U}_q$  above. Our first observation is that, up to symmetry between  $i-1$  and  $i$ ,  $0 \leq |W_{i-1} \cap V(G_1)| - |W_i \cap V(G_1)| \leq k$ . Suppose not. Then there exist  $k + 1$  vertices in  $W_{i-1} \cap V(G_1)$  whose mates are in  $W_i \setminus V(G_1)$ , and thus certify pairwise distinct neighborhoods of the former ones in  $G \setminus V(G_1)$ .

A set  $W_i$  is *crossing*  $G_1$  if  $\emptyset \neq W_i \cap V(G_1) \neq W_i$ . The arguments given in the previous paragraph and  $\frac{1}{3}n^2 \leq |V(G_1)| \leq \frac{2}{3}n^2$  imply that for any  $m$ , if  $n$  is large, there exist sets  $W_{i_0}, W_{i_0+1}, \dots, W_{i_0+m}$  in  $G$  all crossing  $G_1$ . So, we can

select an arbitrarily large index set  $I \subseteq \{i_0, \dots, i_0 + m - 1\}$ ,  $|I| = \ell$ , such that for each  $i \in I$  the element  $a_{i+1}$  is to the right of  $a_i$ , and that all intervals in  $\bigcup_{i \in I} \mathcal{W}_i$  share a common point. In particular,  $a_i$  is not a  $q$ -element and so both  $\mathcal{W}_i$  and  $\mathcal{W}_{i+1}$  are shifted copies of  $\mathcal{U}_1$ . Let  $i_1, \dots, i_\ell$  be the elements of  $\mathcal{I}$  ordered according to the (strictly) increasing values of  $a_i$ , i.e.,  $a_{i_1} < \dots < a_{i_\ell}$ .

Finally, for any  $j, j' \in \{1, \dots, \ell\}$  such that  $j' > j + 1$ , we see that each vertex of  $\mathcal{W}_{i_j} \cap V(G_1)$  cannot have the same neighborhood as any vertex of  $\mathcal{W}_{i_{j'}} \cap V(G_1)$ : this is witnessed by the non-empty set  $\mathcal{W}_{i_{j+1}+1} \setminus V(G_1)$  (represented to the right of the intervals from  $\mathcal{W}_{i_j}$  while intersecting every interval from  $\mathcal{W}_{i_{j'}}$ ). Therefore, the vertices of  $G_1$  have at least  $\ell/2 > k$  distinct neighborhoods in  $G \setminus V(G_1)$ , which contradicts the fact that the clique-width of  $G$  is at most  $k$ .  $\square$

## 5 Graph Interpretation in Interval Graphs

A useful tool when solving the model checking problem on a class of structures is the ability to “efficiently translate” an instance of the problem to a different class of structures, for which we already may have an efficient model checking algorithm. To this end we introduce simple FO graph interpretation, which is an instance of the general concept of interpretability of logic theories [19] restricted to simple graphs with vertices represented by singletons.

An FO *graph interpretation* is a pair  $\mathcal{I} = (\nu, \mu)$  of FO formulae (with 1 and 2 free variables respectively) where  $\mu$  is symmetric, i.e.,  $G \models \mu(x, y) \leftrightarrow \mu(y, x)$  in every graph  $G$ . If  $G$  is a graph, then  $\mathcal{I}(G)$  is the graph defined as follows:

- The vertex set of  $\mathcal{I}(G)$  is the set of all vertices  $v$  of  $G$  such that  $G \models \nu(v)$ , and
- the edge set of  $\mathcal{I}(G)$  is the set of all the pairs  $\{u, v\}$  of vertices of  $G$  such that  $G \models \nu(u) \wedge \nu(v) \wedge \mu(u, v)$ .

We say that a class  $\mathcal{C}_1$  of graphs has an FO interpretation in a class  $\mathcal{C}_2$  if there exists an FO graph interpretation  $\mathcal{I}$  such that every graph from  $\mathcal{C}_1$  is isomorphic to  $\mathcal{I}(G)$  for some  $G \in \mathcal{C}_2$ .

**Lemma 5.1.** *If  $L$  is a subset of non-negative reals that is dense in some non-empty open set, then there exists a polynomially bounded simple FO interpretation of the class of all graphs in the class of  $L$ -interval graphs.*

*Proof.* By scaling, we can assume that  $L$  is dense in  $[1, 1 + \varepsilon]$  for some  $\varepsilon > 0$ . To simplify our considerations, we also assume that  $L = [1, 1 + \varepsilon]$ ; it can be routinely verified that the lengths of intervals appearing in the representation of the graph  $H$  we construct can be slightly perturbed, preserving the structure of the graph.


Let  $G$  be a simple graph with  $n \geq 2$  vertices (the case  $n = 1$  can be handled separately) and let  $v_1, \dots, v_n$  be its vertices. Further, let  $\delta = \frac{\varepsilon}{n+1}$ . We are going to construct an  $L$ -interval graph  $H$  with  $3n+5+|E(G)|$  vertices, and FO formulas  $\mu$  and  $\nu$  such that  $(\mu, \nu)$  is a simple FO interpretation of  $G$  in  $H$ . The formulas  $\mu$  and  $\nu$  will be independent of  $G$  and  $H$ .

The vertex set of  $H$  will be formed by sets  $V_1$ ,  $V_2$  and  $V_3$  (each of them containing  $n + 1$  vertices), a set  $W$  (containing  $|E(G)|$  vertices) and two special vertices  $a$  and  $b$ . Specifically, the set  $V_i$ ,  $i = 1, 2, 3$ , contains vertices  $t_{i,j}$ ,  $j = 0, \dots, n$ , and the set  $W$  contains vertices  $e_{j,j'}$  for all pairs  $1 \leq j < j' \leq n$  such that  $v_j v_{j'} \in E(G)$ .

We now describe an  $L$ -representation of  $H$ . To simplify our description, we use closed intervals but an appropriate perturbation can translate the representation to the model of half-open intervals considered elsewhere in the paper.

- For  $i = 1, 2, 3$  and  $j = 0, 1, \dots, n$ , the interval corresponding to  $t_{i,j}$  is the unit interval  $[i - 1 + (i + j)\delta, i + (i + j)\delta]$ .
- The interval corresponding to  $a$  is the unit interval  $[0, 1]$ .
- The interval corresponding to  $b$  is the unit interval  $[(n + 2)\delta, 1 + (n + 2)\delta]$ .
- The interval corresponding to a vertex  $e_{j,k}$  is the interval  $[1 + (1 + j)\delta, 2 + (k + 3)\delta]$ .

An illustration of the construction can be found in Figure 1. Observe that the vertices  $a$  and  $t_{1,0} \in V_1$  are *twins*, i.e., they have the same neighbors in  $H$ . The vertex  $b$  is adjacent to every vertex in  $V_1 \cup V_2 \cup \{a\} \cup W$ .



**Fig. 1.** Construction of the interval representation of the graph  $H$  in the proof of Lemma 5.1.

Our aim is to interpret the vertices of  $G$  in the set  $V'_1 = \{t_{1,1}, \dots, t_{1,n}\}$ . Since the vertices  $a$  and  $t_{1,0}$  are the only twins in the constructed graph  $H$ , we can identify any one of them by the following formula:

$$\text{anchor}(x) \equiv \exists y (x \neq y \wedge \text{edge}(x, y) \wedge \forall z \neq x, y. \text{edge}(x, z) \leftrightarrow \text{edge}(y, z)).$$

In what follows, we refer to these two vertices as *anchors*.

Let  $\text{dist}(x, y) = c$  for an integer  $c$  stand for an FO formula expressing that the distance of  $x$  and  $y$  is  $c$ , and let  $\text{adist}(x) = c$  stand for an FO formula expressing that the distance of  $x$  from an anchor is  $c$ , i.e.,

$$\text{adist}(x) = c \equiv \exists y (\text{anchor}(y) \wedge \text{dist}(x, y) = c).$$

Using the just defined notation, the following formula is true for exactly the vertices of  $V_1'$ :

$$\mu(x) \equiv \neg \text{anchor}(x) \wedge \text{adist}(x) = 1 \wedge \exists y (\text{adist}(y) = 2 \wedge \neg \text{edge}(x, y)).$$

Note that the latter part makes  $\mu(x)$  false for  $x = b$ . So,  $H \models \mu(t)$  iff  $t \in V_1'$  as needed.

Notice that the set  $V_3$  contains exactly the vertices at distance three or four from the anchors. We say that a vertex  $x' = t_{3,j'} \in V_3$  is the *mate* of a vertex  $x = t_{1,j} \in V_1'$  if  $j = j'$ . The following formula is true iff  $x'$  is the mate of  $x$ .

$$\begin{aligned} \text{mates}(x, x') &\equiv \mu(x) \wedge (\text{adist}(x') = 3 \wedge \text{adist}(x') = 4) \wedge \\ &\quad \exists! y (\text{adist}(y) = 2 \wedge \neg \text{edge}(x, y) \wedge \neg \text{edge}(x', y)) \end{aligned}$$

where  $\exists!$  is a shortcut for “there exists unique”. Note that if  $x = t_{1,j}$  and  $x' = t_{3,j'}$ , then no  $y$  as above exists if  $j' < j$ , and there are several choices of  $y$ , in particular,  $t_{2,j}, \dots, t_{2,j'}$ , if  $j' > j$ .

The main trick of our construction is the ability to “determine the position” of each of the intervals corresponding to  $e_{j,k}$  with respect to the intervals corresponding to the vertices in  $V_1$  and  $V_3$ . To do so, we use the orderings on the intervals corresponding to the vertices of  $V_1$  and  $V_3$  given by their left end points. This ordering is reflected by their neighborhoods in  $V_2$ .

Let us make this more precise. A vertex  $x$  *dominates* a vertex  $y$  if  $y$  and all its neighbors are also neighbors of  $x$ . This property can be expressed using an FO formula as follows:

$$\text{domin}(x, y) \equiv x \neq y \wedge \text{edge}(x, y) \wedge \forall z (\text{edge}(y, z) \rightarrow \text{edge}(x, z)).$$

Using this formula, we can write down the FO formula for  $\nu$  as follows:

$$\begin{aligned} \nu(x, y) &\equiv \nu'(x, y) \vee \nu'(y, x), \quad \text{where} \\ \nu'(x, y) &\equiv \mu(x) \wedge \mu(y) \wedge x \neq y \wedge \exists y', z [\text{mates}(y, y') \wedge \\ &\quad \text{edge}(x, z) \wedge \forall t (\text{domin}(x, t) \rightarrow \neg \text{edge}(t, z)) \wedge \\ &\quad \text{edge}(y', z) \wedge \forall t (\text{domin}(y', t) \rightarrow \neg \text{edge}(t, z))] . \end{aligned}$$

Note that  $\nu'(x, y)$  is true if and only if there is an interval that which intersects both the intervals corresponding to  $x \in V_1'$  and  $y' \in V_3$ , where  $y'$  is the mate of  $y$ , but not the intervals to the left from the interval corresponding for  $x$  and to the right from the interval corresponding for  $y'$ . If such an interval exists, then it corresponds to the vertex of  $W$  associated with the edge  $xy$ . So, the graph  $H$  is indeed an interpretation of  $G$ .  $\square$

Since many FO properties are  $W[2]$ -hard for general graphs, we can immediately conclude the following.

**Corollary 5.2.** *If  $L$  is a subset of non-negative reals that is dense in some non-empty open set, then FO model checking is  $W[2]$ -hard on  $L$ -interval graphs.*

We now turn our attention to interpretation in unit interval graphs. The price we pay for restricting to a smaller class of interval graphs is the strength of the interpretation language used, namely that of MSO logic. At this point we remark that there exist two commonly used MSO frameworks for graphs; the  $\text{MSO}_1$  language which is allowed to quantify over vertices and vertex sets only, and  $\text{MSO}_2$  which is in addition allowed to quantify over edges and edge sets. We stay with the former weaker one in this paper.

An  $\text{MSO}_1$  graph interpretation is defined in the analogous way to former FO interpretation with the formulas  $\mu$  and  $\nu$  being  $\text{MSO}_1$  formulas.

**Lemma 5.3.** *There is a polynomially bounded simple  $\text{MSO}_1$  interpretation of the class of all graphs into the class of unit interval graphs.*

*Proof.* Suppose that we are given an  $n$ -vertex  $G$  with  $n \geq 4$  (the cases with  $n = 1, 2, 3$  could be held separately). Our aim is to find a unit interval graph  $H$  and  $\text{MSO}_1$  formulas  $\mu$  and  $\nu$  such that  $(\mu, \nu)$  is an interpretation of  $G$  in  $H$ . Let  $v_1, \dots, v_n$  be the vertices of  $G$  and  $m$  the number of its edges.

As in the proof of Lemma 5.1, we will describe  $H$  by giving its interval representation and we again use the closed intervals in the representation. Let us introduce the following shorthand notation: if  $J$  is an interval and  $x$  a real, then  $J + x$  is the interval  $J$  shifted by  $x$  to the right. Similarly, if  $\mathcal{I}$  is a set of intervals, then  $\mathcal{I} + x$  is the set of the intervals from  $\mathcal{I}$  shifted by  $x$  to the right.

Choose  $\delta > 0$  such that  $\delta n \ll 1$  and  $\mathcal{U} = \{[i\delta, 1 + i\delta] : i = 0, 1, \dots, n - 1\}$ . The graph  $H$  will contain  $n(3m + 1)$  vertices corresponding to the intervals from the sets  $\mathcal{U} + k(1 + \delta)$  for  $k = 1, \dots, 3m + 1$ . For further reference, we say that the intervals  $[i\delta, 1 + i\delta]$  and  $[i\delta, 1 + i\delta] + k(1 + \delta)$  are at the *same level*  $i$ . Let  $W_\ell$ ,  $\ell = 0, \dots, m$ , be the set of the  $n$  vertices corresponding to the intervals from  $\mathcal{U} + (3\ell + 1)(1 + \delta)$ .


Let  $A$  be the multiset containing three copies of the interval  $[0, 1]$ . The graph  $H$  further contains three vertices corresponding to the intervals in  $A$  and  $3m$  vertices corresponding to the intervals in  $A + (3i - 1/2)(1 + \delta)$ ,  $i = 1, \dots, m$ . The vertices of  $H$  corresponding to these  $3m + 3$  intervals will be referred to as anchors and they will be the only vertices of  $H$  with two twins.

Fix an ordering of the  $m$  edges of  $G$ . If the  $\ell$ -th edge joins vertices  $v_j$  and  $v_k$ , include an additional pair of vertices corresponding to the intervals  $[j\delta, 1 + j\delta] + (3\ell + 1)(1 + \delta)$  and  $[k\delta, 1 + k\delta] + (3\ell + 1)(1 + \delta)$  to  $H$ . The vertices included in this step are the only vertices of  $H$  that have unique twins.

Finally, the graph  $H$  also contains a vertex corresponding to the interval  $[1/2, 3/2]$ . Observe that the three vertices of  $H$  corresponding to the vertices of  $A$  are the only anchors in  $H$  with degree four. The whole construction of  $H$  is illustrated in Figure 5.

We now give the  $\text{MSO}_1$  formulas  $\mu$  and  $\nu$  such that  $(\mu, \nu)$  is a simple  $\text{MSO}_1$  interpretation of  $G$  in  $H$ . First, recall the FO property  $\text{twin}(x, y)$  from the proof of Lemma 5.1 which expresses that the vertices  $x$  and  $y$  are twins;

$$\text{twin}(x, y) \equiv (x \neq y \wedge \forall z \neq x, y. \text{edge}(x, z) \leftrightarrow \text{edge}(y, z)).$$



**Fig. 2.** An illustration of the interval representation  $Y$  (of the graph  $H$ ) which shows the interpretation of an edge  $e_1 = v_j v_k$  (of the graph  $G$ ).

Using this formula, we can give formulas to identify anchors in  $H$  and the formula to identify vertices not adjacent to any anchors (these are the vertices in the sets  $W_1, \dots, W_m$  and the  $2m$  twins of them included because of the edges of  $G$ ):

$$\begin{aligned} \text{anchor}(x) &\equiv \exists y, z (z \neq y \wedge \text{edge}(z, y) \wedge \text{twin}(x, y) \wedge \text{twin}(x, z)), \\ \text{noanch}(x) &\equiv \forall z (\text{anchor}(z) \rightarrow \neg \text{edge}(x, z)). \end{aligned}$$

The vertices of  $H$  used to model the vertices of  $G$  are those in the set  $W_0$ . These are precisely the vertices not adjacent to any anchor with distance two from the three anchors of degree four. So, they can be identified using the following FO formula.

$$\mu(x) \equiv \text{noanch}(x) \wedge \exists t (\text{anchor}(t) \wedge \text{deg}(t) = 4 \wedge \text{dist}(t, x) = 2).$$

We now introduce a notion of mates. Two vertices are mates if they belong to the sets  $W_{p-1}$  and  $W_p$  for  $p = 1, \dots, 3m$  and the intervals corresponding to them are at the same level. Hence, two vertices  $x$  and  $x'$  are mates iff they satisfy the following formula.

$$\begin{aligned} \text{mates}(x, x') &\equiv \text{noanch}(x) \wedge \text{noanch}(x') \wedge \text{dist}(x, x') = 4 \wedge \exists t \\ &\quad [\text{anchor}(t) \wedge \exists! y \exists! z (\neg \text{edge}(y, z) \wedge \text{edge}(y, t) \wedge \text{edge}(z, t) \wedge \\ &\quad \text{dist}(x, y) = 2 \wedge \text{dist}(x', y) > 2 \wedge \text{dist}(x', z) = 2 \wedge \text{dist}(x, z) > 2)]. \end{aligned}$$

Let  $\text{mates}^*(x, y)$  be the formula describing the transitive closure of the relation “being mates” (this is the only place in the proof where we need the expressive power of an  $\text{MSO}_1$  formula). The edges of  $G$  can now be interpreted in  $H$  using the following formula.

$$\begin{aligned} \nu(x, y) &\equiv \mu(x) \wedge \mu(y) \wedge x \neq y \wedge \exists x_1, x_2, y_1, y_2 (\text{edge}(x_1, y_1) \wedge \\ &\quad \text{mates}^*(x, x_1) \wedge \text{mates}^*(y, y_1) \wedge \text{twin}(x_1, x_2) \wedge \text{twin}(y_1, y_2)). \end{aligned}$$

Indeed, if  $x$  and  $y$  belong to  $W_0$ , then  $\nu(x, y)$  is true only if there exist vertices  $x_1$  and  $y_1$  at the same level as  $x$  and  $y$ , respectively, that are adjacent (and therefore, in the same set  $W_p$ ) and both  $x_1$  and  $y_1$  have twins. However, this happens only if  $xy$  should interpret an edge  $e_p$  of  $G$ .  $\square$

Again, we can immediately conclude the following.

**Corollary 5.4.** *MSO<sub>1</sub> model checking is W[2]-hard on unit interval graphs.*

This corollary is rather tight since the aforementioned result of Lozin [15] claims that every proper hereditary subclass of unit interval graphs has bounded clique-width, and hence MSO<sub>1</sub> model checking on this class is in linear time [2].

Lastly, we remark that Fellows et al [9] have shown that testing FO properties on unit two-interval graphs (i.e., such that each vertex corresponds to a pair of intervals, each on a distinct line) is W[1]-hard.

## References

1. B. Courcelle. The monadic second order logic of graphs I: Recognizable sets of finite graphs. *Inform. and Comput.*, 85:12–75, 1990.
2. B. Courcelle, J. A. Makowsky, and U. Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory Comput. Syst.*, 33(2):125–150, 2000.
3. B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. *Discrete Appl. Math.*, 101(1-3):77–114, 2000.
4. A. Dawar, M. Grohe, and S. Kreutzer. Locally excluding a minor. In *LICS'07*, pages 270–279. IEEE Computer Society, 2007.
5. A. Dawar and S. Kreutzer. Parameterized complexity of first-order logic. *Electronic Colloquium on Computational Complexity (ECCC)*, TR09-131, 2009.
6. R. Downey and M. Fellows. *Parameterized complexity*. Monographs in Computer Science. Springer, 1999.
7. Z. Dvořák, D. Král', and R. Thomas. Deciding first-order properties for sparse graphs. In *FOCS'10*, pages 133–142. IEEE Computer Society, 2010.
8. A. Ehrenfeucht. An application of games to the completeness problem for formalized theories. *Fund. Math.*, 49:129–141, 1961.
9. M. Fellows, D. Hermelin, F. Rosamond, and S. Vialette. On the parameterized complexity of multiple-interval graph problems. *Theoret. Comput. Sci.*, 410(1):53–61, 2009.
10. R. Fraïssé. Sur quelques classifications des systèmes de relations. *Université d'Alger, Publications Scientifiques, Série A*, 1:35–182, 1954.
11. M. Frick and M. Grohe. Deciding first-order properties of locally tree-decomposable structures. *J. ACM*, 48(6):1184–1206, 2001.
12. M. Golumbic and U. Rotics. On the clique-width of some perfect graph classes. *Int. J. Found. Comput. Sci.*, 11(3):423–443, 2000.
13. M. Grohe and S. Kreutzer. Methods for algorithmic meta theorems. In *Model Theoretic Methods in Finite Combinatorics: AMS-ASL Special Session, January 5-8, 2009*, Contemporary Mathematics, pages 181–206. AMS, 2011.
14. S. Kreutzer. Algorithmic meta-theorems. *Electronic Colloquium on Computational Complexity (ECCC)*, TR09-147, 2009.
15. V. Lozin. From tree-width to clique-width: Excluding a unit interval graph. In *ISAAC'08*, volume 5369 of *LNCS*, pages 871–882. Springer, 2008.
16. J. Nešetřil and P. O. de Mendez. Grad and classes with bounded expansion I. Decompositions. *European J. Combin*, 29(3):760–776, 2008.
17. J. Nešetřil and P. O. de Mendez. Grad and classes with bounded expansion II. Algorithmic aspects. *European J. Combin*, 29(3):777–791, 2008.

18. J. Nešetřil and P. O. de Mendez. Grad and classes with bounded expansion III. Restricted graph homomorphism dualities. *European J. Combin.*, 29(4):1012–1024, 2008.
19. M. O. Rabin. A simple method for undecidability proofs and some applications. In Y. Bar-Hillel, editor, *Logic, Methodology and Philosophy of Sciences*, volume 1, pages 58–68. North-Holland, Amsterdam, 1964.
20. D. Seese. Linear time computable problems and first-order descriptions. *Math. Structures Comput. Sci.*, 6(6):505–526, 1996.