



FI MU

**Faculty of Informatics
Masaryk University**

Complexity Issues of the Pattern Equations in Idempotent Semigroups

by

**Ondřej Klíma
Jiří Srba**

FI MU Report Series

FIMU-RS-99-02

Copyright © 1999, FI MU

August 1999

Complexity Issues of the Pattern Equations in Idempotent Semigroups^{*}

Ondřej Klíma¹ and Jiří Srba²

¹ Faculty of Science MU, Dept. of Mathematics, Janáčkovo nám. 2a, 662 95 Brno, Czech Republic, klima@math.muni.cz

² Faculty of Informatics MU, Botanická 68a, 602 00 Brno, Czech Republic, srba@fi.muni.cz

Abstract. A pattern equation is a word equation of the form $X = A$ where X is a sequence of variables and A is a sequence of constants. The problem whether $X = A$ has a solution in a free idempotent semigroup (free band) is shown to be NP-complete.

1 Introduction

The main topic of interest in several fields of computer science is solving equations. Many areas such as logic programming, automated theorem proving and pattern matching exploit solving equations, and unification is a typical example of it. An important role is played also by semantic unification, which is in fact solving word equations in some variety. Makanin (see [10]) shows that the question whether an equation in a free monoid has a solution is decidable. It can be even generalized in the way that existential first-order theory of equations over free monoid is decidable including additional regular constraints on the variables [13].

In this paper we focus on a certain subclass of equations which we call pattern equations. Pattern equation is a word equation where we have on the left-hand side just variables and on the righthand side only constants. Many practical problems such as pattern matching and speech recognition/synthesis lead to this kind of equations. This work has been inspired by the papers [5] and [6] where the basic approach – syllable-based speech synthesis – is in assigning prosody attributes to a given text and segmentation into syllable segments. We examine the solvability of word equations in the variety of idempotent semigroups (bands) which we call stuttering equations. Their name comes from practical motivation. For example in speech recognition the speaker sometimes stutters some words and we would like to eliminate this effect and enable the correct variables assignation even in the case of stuttering. Therefore we allow to eliminate multiple occurrences of the same constant into only one occurrence, which can be modelled by the identity $x^2 = x$. The decidability of the

* The paper is supported by the Grant Agency of the Czech Republic, grant No. 201/97/0456 and by grant FRVŠ 409/1999

satisfiability problem (even in the general case) is a consequence of the local finiteness of the free idempotent semigroup and an exponential upper bound on the length of a minimal solution can be given [3]. We can also establish a polynomial time decision procedure for the word problem in a free idempotent semigroup.

In this paper we prove that there is a polynomial bound on the length of a minimal solution in the case of stuttering pattern equations and thus show that the satisfiability problem is in NP. The proof exploits the confluent and terminating word rewriting system for bands by Siekmann and Szabo (see [14]). This means that the identity $p = q$ holds in a free idempotent semigroup if and only if the words p and q have the same normal form w.r.t. the rewriting system $\{xx \rightarrow x \mid c(x) \neq \emptyset\} \cup \{uvw \rightarrow uw \mid \emptyset \neq c(v) \subseteq c(u) = c(w)\}$.

Moreover using the reduction from 3-SAT to our problem, we get its NP-completeness. This shows that the problem is unfortunately intractable. One of the ways how to solve the problem is to use heuristic algorithms. They are the current field of interest in speech recognition. Another approach that could be used for solving the problem is *Concurrent Constraint Programming*. For the background see [2].

2 Basic definitions

An idempotent semigroup (also called a band) is a semigroup where the identity $x^2 = x$ is satisfied. Let \mathcal{C} be a finite set and we define a binary relation $\rightarrow \subseteq \mathcal{C}^* \times \mathcal{C}^*$ such that $uvvw \rightarrow uvw$ for $u, v, w \in \mathcal{C}^*$ and let \sim be its symmetric and transitive closure, i.e. $\sim := (\rightarrow \cup \rightarrow^{-1})^*$. Then the identity $p = q$ holds in a free band over \mathcal{C} if and only if $p \sim q$ (completeness of the equational logic).

Let \mathcal{C} be a finite set of *constants* and \mathcal{V} be a finite set of *variables* such that $\mathcal{C} \cap \mathcal{V} = \emptyset$. A *word equation* $L = R$ is a pair $(L, R) \in (\mathcal{C} \cup \mathcal{V})^* \times (\mathcal{C} \cup \mathcal{V})^*$. A *system of word equations* is a finite set of equations of the form $\{L_1 = R_1, \dots, L_n = R_n\}$ for $n > 0$. A *solution (in a free idempotent semigroup)* of such a system is a homomorphism $\alpha : (\mathcal{C} \cup \mathcal{V})^* \rightarrow \mathcal{C}^*$ which behaves as an identity on the letters from \mathcal{C} and equates all the equations of the system, i.e. $\alpha(L_i) \sim \alpha(R_i)$ for all $1 \leq i \leq n$. Such a homomorphism is then fully established by a mapping $\alpha : \mathcal{V} \rightarrow \mathcal{C}^*$. A solution is called *non-singular*, if $\alpha(x) \neq \epsilon$ for all $x \in \mathcal{V}$. Otherwise we will call it *singular*. We say that the system of word equations (in a free idempotent semigroup) is *satisfiable* whenever it has a solution. For the introduction into word equations and combinatorics on words we refer to [7], [8] and [12].

In what follows we will use a uniform notation. The set $\mathcal{C} = \{a, b, c, \dots\}$ denotes the alphabet of constants and $\mathcal{V} = \{x, y, z, \dots\}$ stands for variables (unknowns) with the assumption that $\mathcal{C} \cap \mathcal{V} = \emptyset$. We will use the same symbol α for the mapping $\alpha : \mathcal{V} \rightarrow \mathcal{C}^*$ and its unique extension to a homomorphism $\alpha : (\mathcal{C} \cup \mathcal{V})^* \rightarrow \mathcal{C}^*$. The empty word will be denoted as ϵ and the length of a word w as $|w|$.

We exploit the fact that the word problem in a free band is decidable (see [4] and its generalization [9]), which is a consequence of the next lemma. Let $w \in \mathcal{C}^*$. We define

- $c(w)$ – the set of all letters that occur in w ,
- $0(w)$ – the longest prefix of w in $\text{card}(c(w)) - 1$ letters,
- $1(w)$ – the longest suffix of w in $\text{card}(c(w)) - 1$ letters.

Let also $\bar{0}(w)$ resp. $\bar{1}(w)$ be the letter that immediately succeeds $0(w)$ resp. precedes $1(w)$.

Lemma 1 ([4]). *Let $p, q \in \mathcal{C}^*$. Then $p \sim q$ if and only if $c(p) = c(q)$, $0(p) \sim 0(q)$ and $1(p) \sim 1(q)$.*

It is obvious that if a stuttering equation system has a solution then it has always infinitely many solutions, which we show in the following lemma.

Lemma 2. *Let $\{L_1 = R_1, \dots, L_n = R_n\}$ be a stuttering equation system and α its solution. Then also any β that satisfies $\alpha(x) \sim \beta(x)$ for all $x \in \mathcal{V}$ (we simply write $\alpha \sim \beta$) is a solution.*

Proof. Immediate. □

This gives an idea that we should look just for the solutions where $\alpha(x)$ is the shortest word in the \sim class for each variable x . We introduce the size of the solution α as $\text{size}(\alpha) := \max_{x \in \mathcal{V}} |\alpha(x)|$ and say that α is *minimal* iff for any solution β of the system we have $\text{size}(\alpha) \leq \text{size}(\beta)$. Given a stuttering equation system it is decidable whether the system is satisfiable because of the local finiteness of free idempotent semigroups. The following lemma just gives a precise exponential upper bound on the size of a minimal solution.

Lemma 3 ([3]). *Let $k = \text{card}(\mathcal{C}) \geq 2$ and let $\{L_1 = R_1, \dots, L_n = R_n\}$ be a stuttering equation system. If the system is satisfiable then there exists a solution α such that $\text{size}(\alpha) \leq 2^k + 2^{k-2} - 2$.*

In general it can be shown that there are stuttering equation systems such that all their solutions are at least exponentially large w.r.t. the cardinality of the set \mathcal{C} . Consider the following sequence of equations: $x_1 = a_1$ and $x_{i+1} = x_i a_{i+1} x_i$ for a sequence of pairwise different constants a_1, a_2, \dots . For any solution α of the system we have that $|\alpha(x_i)| \geq 2^i - 1$.

In this paper we focus on a special kind of word equations which we call *pattern equations*.

Definition 1. *A pattern equation system is a set $\{X_1 = A_1, \dots, X_n = A_n\}$ where $X_i \in \mathcal{V}^*$ and $A_i \in \mathcal{C}^*$ for all $1 \leq i \leq n$. The solution of the pattern equation system is defined as in the general case.*

Two natural decidability problems (PATTERN-EQUATION and NON-SINGULAR-PATTERN-EQUATION problem) appear in this context and are defined below.

Definition 2. *Given a pattern equation system $\{X_1 = A_1, \dots, X_n = A_n\}$ as an instance of the PATTERN-EQUATION problem, the task is to decide whether this system has a solution. If we require the solution to be non-singular we call the problem NON-SINGULAR-PATTERN-EQUATION .*

The PATTERN-EQUATION problem for a single stuttering pattern equation $X = A$ is trivial since it is always solvable: $\alpha(x) = A$ for all $x \in \mathcal{V}$. On the other hand a system is not always solvable: e.g. $\{x = a, x = b\}$ has no solution.

We give an example of a pattern equation system and demonstrate its solutions.

Example 1. Let us have the following system where $\mathcal{C} = \{a, b\}$, $\mathcal{V} = \{x, y, z\}$ and the pattern equations are $\{yxy = aba, yz = a\}$. A singular solution exists $\alpha(x) = aba, \alpha(y) = \epsilon, \alpha(z) = a$, however, there is also a non-singular solution $\beta(x) = bab, \beta(y) = a, \beta(z) = a$ since $ababa \sim aba$ and $aa \sim a$. We have also another non-singular solution $\gamma(x) = ab, \gamma(y) = a, \gamma(z) = a$.

Our goal is to show that a minimal solution of a stuttering pattern equation system is of a polynomial length. This implies that the problem of deciding whether a stuttering pattern equation system is satisfiable is in NP.

3 Rewriting system for idempotent semigroups

In this section we summarize several properties of the rewriting system by Siekmann and Szabo in [14] and prove some technical lemmas. First of all we have to give some definitions and results concerning rewriting systems as it can be found e.g. in [1].

A *rewriting system* R over \mathcal{C} is a subset of $\mathcal{C}^* \times \mathcal{C}^*$. The elements of R will be called *rules*. Having such a system R we can define a *rewrite relation* $\rightarrow \subseteq \mathcal{C}^* \times \mathcal{C}^*$ in the following way:

$$\forall p, q \in \mathcal{C}^* : p \rightarrow q \text{ iff } \exists (u, v) \in R, s, t \in \mathcal{C}^* : p = sut, q = svt.$$

The elements (u, v) of R will be often written as $u \rightarrow v$. For a word $q \in \mathcal{C}^*$ we write $q \not\rightarrow$ iff there is no q' such that $q \rightarrow q'$ and we say that q is in a *normal form*. We define the set of normal forms of $p \in \mathcal{C}^*$ as $\langle p \rangle = \{q \mid p \rightarrow^* q \not\rightarrow\}$. We say that R (resp. the relation \rightarrow) is *terminating* iff there is no infinite sequence $p_1, p_2, p_3 \dots \in \mathcal{C}^*$ such that $p_1 \rightarrow p_2 \rightarrow p_3 \rightarrow \dots$. The system R (resp. the relation \rightarrow) is *confluent* iff

$$\forall p, p_1, p_2 \in \mathcal{C}^* \exists q \in \mathcal{C}^* : \text{if } (p \rightarrow^* p_1 \text{ and } p \rightarrow^* p_2) \text{ then } (p_1 \rightarrow^* q \text{ and } p_2 \rightarrow^* q).$$

The system R (resp. the relation \rightarrow) is *locally confluent* iff

$$\forall p, p_1, p_2 \in \mathcal{C}^* \exists q \in \mathcal{C}^* : \text{if } (p \rightarrow p_1 \text{ and } p \rightarrow p_2) \text{ then } (p_1 \rightarrow^* q \text{ and } p_2 \rightarrow^* q).$$

Following lemma shows the relationship between confluence and local confluence.

Lemma 4 ([1]). *Let R be a terminating rewriting system. Then R is confluent if and only if R is locally confluent.*

It is easy to see that if R is confluent and terminating rewriting system, then a word $p \in \mathcal{C}^*$ has exactly one normal form, i.e. $\langle p \rangle = \{q\}$ for some q , and in such a case we simply write $\langle p \rangle = q$.

Example 2. Let $\{xx \rightarrow x \mid x \in \mathcal{C}^*, c(x) \neq \emptyset\}$ be a rewriting system over \mathcal{C} . Then this system is terminating but it is not confluent. For $p = ababcbabc$ we have $p = (ab)(ab)cbabc \rightarrow abcbabc$ and $p = a(babc)(babc) \rightarrow (ab)(ab)c \rightarrow abc$ where $abcbabc$ and abc are in normal forms. It is easy to see that $\langle p \rangle = \{abc, abcbabc\}$.

In this paper we will exploit the rewriting system by Siekmann and Szabo in [14].

Lemma 5 ([14]). *The rewriting system $\{xx \rightarrow x \mid x \in \mathcal{C}^*, c(x) \neq \emptyset\} \cup \{uvw \rightarrow uv \mid u, v, w \in \mathcal{C}^*, \emptyset \neq c(v) \subseteq c(u) = c(w)\}$ is confluent and terminating. Moreover for $p, q \in \mathcal{C}^*$ we have $p \sim q$ if and only if p and q have the same normal form w.r.t the system.*

We will refer the rewriting system $\{xx \rightarrow x \mid c(x) \neq \emptyset\} \cup \{uvw \rightarrow uv \mid \emptyset \neq c(v) \subseteq c(u) = c(w)\}$ as RS . Since RS contains two different types of rewriting rules we denote RS_1 the rewriting system $\{xx \rightarrow x \mid c(x) \neq \emptyset\}$ and RS_2 the rewriting system $\{uvw \rightarrow uv \mid \emptyset \neq c(v) \subseteq c(u) = c(w)\}$. The corresponding rewrite relations are denoted $\rightarrow, \rightarrow_1$ resp. \rightarrow_2 and for a word $p \in \mathcal{C}^*$ the set of its normal forms is denoted as $\langle p \rangle, \langle p \rangle_1$ resp. $\langle p \rangle_2$.

If we want to investigate the complexity issues for stuttering equations, the first question we have to answer is the complexity of checking whether some identity holds in a free band. We will show that the word problem (i.e. the problem whether $p \sim q$) can be decided in a polynomial time by using the rewriting system RS . If we note that a string of the length k contains $O(k^2)$ substrings (each substring is identified by its beginning and its length) we get that each reduction of RS can be done in a polynomial time. Since every reduction decreases the length of the word, we have a polynomial time decision algorithm for the word problem in a free band.

We know that RS is confluent and terminating. Our goal in this section is to show that RS_2 is also confluent and terminating rewriting system and that $\langle p \rangle = \langle \langle p \rangle_2 \rangle_1$.

We define a rewrite relation $\rightarrow_{2l} \subset \rightarrow_2$ such that $suvt \rightarrow_{2l} suvt$ if and only if $|v| = 1$ and $c(v) \subseteq c(u) = c(w)$. It is easy to see that $\rightarrow_2 \subseteq \rightarrow_{2l}^*$ and hence $\rightarrow_{2l}^* = \rightarrow_2^*$. The last relation we will use is $\rightarrow_{2m} \subset \rightarrow_2$, consisting of all rules that leave out the maximal number of letters in the following sense. Let $\text{head}(w)$ resp. $\text{tail}(w)$ mean the first resp. the last letter of the word w . We write $suvt \rightarrow_{2m} suvt$ if and only if $\emptyset \neq c(v) \subseteq c(u) = c(w)$ and for any suffix s_1 of s and any prefix t_1 of t (including empty s_1 or t_1) the following conditions hold:

- (i) $c(s_1u) \neq c(wt_1)$
- (ii) $c(u) - c(\text{tail}(u)) \neq c(u)$

(iii) $c(w) - c(\text{head}(w)) \neq c(w)$

Note that if $suvt \rightarrow_{2m} suwt$ then $u = 0(u)\bar{0}(u)$, $w = \bar{1}(w)1(w)$ and the last letter of s and the first letter of t (if they exist) are new and different letters¹. We show that $\langle p \rangle_{2m} = \langle p \rangle_2$. The inclusion $\langle p \rangle_{2m} \subseteq \langle p \rangle_2$ is obvious and the rest is the content of the following lemmas. For more transparent proofs we use the notation $\underline{suvt} \rightarrow_2 suwt$ in the sense that $suvt \rightarrow_2 suwt$ where $\emptyset \neq c(v) \subseteq c(u) = c(w)$ (and the same for \rightarrow_{2l} , \rightarrow_{2m}). In the following, whenever we say that u is a subword of sut , we always refer to the concrete (and obvious) occurrence of the subword u in sut .

Lemma 6. *The relation \rightarrow_{2m} is confluent and terminating.*

Proof. The termination is obvious. Let p be a word and suppose that we can apply two different rules of \rightarrow_{2m} on p , say $p = s_1\underline{u_1}v_1\underline{w_1}t_1 \rightarrow_{2m} s_1u_1w_1t_1$ and $p = s_2\underline{u_2}v_2\underline{w_2}t_2 \rightarrow_{2m} s_2u_2w_2t_2$.

Let us suppose that u_1 is a subword of $u_2v_2w_2$. Then the whole $u_1v_1w_1$ is a subword of $u_2v_2w_2$, because $u_2v_2w_2$ is followed by a new letter (if it exists), which is not contained in u_1 . If $u_1v_1w_1$ is a subword of u_2 resp. w_2 then our two rules commute. In the other case we will show that v_1 is a subword of v_2 , which is a contradiction with the maximality of the rule $s_1\underline{u_1}v_1\underline{w_1}t_1 \rightarrow_{2m} s_1u_1w_1t_1$. Suppose that the occurrence of $\bar{0}(u_2)$ (the last letter of u_2) is in $u_1v_1w_1$, then it is surely in u_1 . Similarly if the occurrence of $\bar{1}(w_2)$ is in $u_1v_1w_1$, then it is in w_1 . This constraints that v_1 is a subword of v_2 .

If u_1 resp. w_1 is not a subword of $u_2v_2w_2$, and u_2 resp. w_2 is not a subword of $u_1v_1w_1$, then our two rules commute. \square

Remark 1. From the previous proof we can see that arbitrary two applications of \rightarrow_{2m} , say $p = s_1\underline{u_1}v_1\underline{w_1}t_1 \rightarrow_{2m} s_1u_1w_1t_1$ and $p = s_2\underline{u_2}v_2\underline{w_2}t_2 \rightarrow_{2m} s_2u_2w_2t_2$, commute and they can be nested exactly in one of the following ways (up to symmetry):

1. $w_1 = w'_1q, u_2 = qu'_2$ and $p = s_1u_1v_1w'_1qu'_2v_2w_2t_2$
2. $u_1v_1w_1$ is a subword of u_2

Lemma 7. *RS_2 is a confluent and terminating rewriting system and $\langle p \rangle_{2m} = \langle p \rangle_2$ for any $p \in \mathcal{C}^*$.*

Proof. The termination is clear. If we have $u \rightarrow_2 v, u \rightarrow_2 w$ then there is v_1 and w_1 such that $v \rightarrow_2 v_1, w \rightarrow_2 w_1, u \rightarrow_{2m} v_1$ and $u \rightarrow_{2m} w_1$. Since \rightarrow_{2m} is confluent, we have the confluence of RS_2 . The equality $\langle p \rangle_{2m} = \langle p \rangle_2$ is a trivial consequence. \square

Lemma 8. *For any $p, q \in \mathcal{C}^*$ such that $p = \langle p \rangle_2$ and $p \rightarrow_1 q$ it holds that $\langle q \rangle_2 = q$. In particular for a word $p \in \mathcal{C}^*$ we have $\langle \langle p \rangle_2 \rangle_1 = \langle p \rangle$.*

¹ Observe that it doesn't hold that if $p \rightarrow_{2m} q$ then $spt \rightarrow_{2m} sqt$ for $s, t \in \mathcal{C}^*$. This means that \rightarrow_{2m} is not a rewriting relation in the previously introduced sense.

Proof. Assume for the moment that $\langle q \rangle_2 \neq q$, which means that $q = suawt$ where $s, u, w, t \in \mathcal{C}^*$, $a \in \mathcal{C}$, $a \in c(u) = c(w)$, i.e. $q = \underline{suawt} \rightarrow_{2l} suwt$. Then (w.l.o.g.) $p = su_1xu_2awt$ where $u_1, x, u_2 \in \mathcal{C}^*$, $u_1u_2 = u$ and su_1 has a suffix x or u_2awt has a prefix x . We discuss four different cases.

- 1) x is a suffix of u_1
- 2) x is a suffix of su_1 and $|x| > |u_1|$
- 3) x is a prefix of u_2aw
- 4) x is a prefix of u_2awt and $|x| > |u_2aw|$

In the case 1) we get $c(x) \subseteq c(u) = c(w)$ and we could also use the reduction $p = \underline{su_1xu_2awt} \rightarrow_{2l} su_1xu_2wt$ since $a \in c(u_1xu_2) = c(u) = c(w)$. In the case 2) we may write $x = x_1u_1$ and then $p = su_1x_1u_1u_2awt = su_1x_1\underline{uawt} \rightarrow_{2l} su_1x_1uwt$. Cases 3) and 4) are similar and all the four cases lead to a contradiction. \square

4 Upper bound for the size of the solution

This section aims to prove that the PATTERN-EQUATION problem is in NP by giving a polynomial upper bound on the size of a minimal solution. In the following we assume implicitly that $A, B \in \mathcal{C}^*$. Realise that each reduction of RS just leaves out some subword, the case $uvw \rightarrow uw$ is clear, and in the case $xx \rightarrow x$ we leave out the right occurrence of x in the square. If we have a word uAv , we can speak about the *residual* of A in the sense that the residual consists of all letter occurrences of A that were not left out during the sequence of reductions. Moreover if we use two different sequences of reductions by \rightarrow_{2m} , which give normal form w.r.t. \rightarrow_2 , then the residuals are the same after the both reduction sequences, since any two applications of \rightarrow_{2m} commute by the Remark 1.

Lemma 9. *Let A and B be in normal forms and $AB \rightarrow_{2m} AB' \not\rightarrow_{2m}$ where B' is the residual of B . Then the word B' contains at most one square x^2 which arises from xvx where v is the word left out by the reduction rule $\underline{uvw} \rightarrow_{2m} uw$, and x is both a suffix of u and a prefix of w . Moreover in the case when B' contains a square we have $B' \rightarrow_1 \langle B' \rangle$.*

Proof. Assume that we have used $AB = \underline{suvw}t \rightarrow_{2m} suwt = AB'$ and B' contains a square x^2 . Since B is in normal form, x^2 contains "the space" of the cancelled v , i.e. $xx = u_1w_1$ where u_1 is a suffix of u (u starts in A) and w_1 is a prefix of wt .

We show that w_1 is a prefix of w . In the case when $|w_1| > |w|$ we can deduce that occurrences of $\text{tail}(u_1)$ and $\text{head}(t)$ must lie in the left x since they are the first occurrences of the constants $\text{tail}(u_1)$ and $\text{head}(t)$ in B (from the maximality of \rightarrow_{2m}). It means that $x = u_1wz$ where z is a prefix of t . Since $c(u_1) \subseteq c(u) = c(w) \subseteq c(wz)$, we can reduce $xx = u_1\underline{wz}u_1\underline{wz} \rightarrow_2 u_1wzww$ and this is a contradiction with $B' \not\rightarrow_2$.

So, w_1 is a prefix of w . The last letter of u_1 is in the left x and the first letter of w_1 is in the right x and we see that x^2 arises from xvx and x is a suffix of u and a prefix of w (i.e. $u = u_0x$, $w = xw_0$).

We have $uwt = u_0xxw_0t \rightarrow_1 u_0xw_0t$. It is enough to show that the word u_0xw_0t does not contain a square and in such a case we get $B' \rightarrow_1 \langle B' \rangle_1$. Assume that u_0xw_0t contains a square y^2 . Recall $xw_0t = wt$ is a suffix of B . Thus y^2 is a subword of u_0xw_0 since $\text{head}(t) \notin c(u_0xw_0)$ and because of the similar arguments as in the second paragraph. Now $u_0x = u$ and $xw_0 = w$ imply that y^2 contains both $\text{tail}(u_0)$ and $\text{head}(w_0)$. However, $\text{tail}(x) = \text{tail}(u)$ is the first occurrence of this letter in u_0xw_0 and it must be in the left y and from the same reason $\text{head}(x) = \text{head}(w)$ is in the right y . This is impossible. So u_0xw_0t contains no square.

The case when B' contains two or more squares is a contradiction with $B' \rightarrow_1 \langle B' \rangle_1$. \square

Remark 2. The same arguments as in the proof of the Lemma 9 give the following analogue.

If A_1, B, A_2 are in normal forms and $A_1BA_2 \rightarrow_{2m} A_1B'A_2 \not\rightarrow_{2m}$ and if the residual B' of B contain a square x^2 , then $B' \rightarrow_1 \langle B' \rangle_1$ and x has the same properties as in the Lemma 9.

Proposition 1. *Let A and B be in normal forms such that $\langle AB \rangle_2 = AB'$ where B' is the residual of B , then $|B'| \leq |\langle B' \rangle|^2$.*

Proof. By the Lemma 7 we have $\langle AB \rangle_2 = \langle AB \rangle_{2m}$ and we can use the maximal reductions. W.l.o.g. assume that the reductions \rightarrow_{2m} did not leave out some prefix B_1 of the word B , otherwise we can start with the words A and B_2 where $B = B_1B_2$. The Remark 1 shows how two applications of \rightarrow_{2m} can be nested. Since A and B are in normal forms, we can see that any reduction \rightarrow_{2m} uses some letters from both A and B . This means that we can write $A = s_{n+1}s_n \dots s_1$, $B = u_1v_1w_1 \dots u_nv_nw_nu_{n+1}$ where $s_i, u_i, v_i, w_i \in \mathcal{C}^*$ for all possible i and we have n reductions of the form

$$s_{n+1} \dots \underline{s_i \dots s_1 u_1 w_1 \dots u_i v_i w_i} \dots u_{n+1} \rightarrow_{2m} s_{n+1} \dots s_1 u_1 w_1 \dots u_i w_i \dots u_{n+1}$$

where $c(v_i) \subseteq c(s_i \dots s_1 u_1 w_1 \dots u_i) = c(w_i)$ and $B' = u_1 w_1 \dots u_n w_n u_{n+1}$.

Since each step of the maximal reduction needs a new letter (the letter that immediately succeeds w_i), we get an upper bound for n (the number of steps in \rightarrow_{2m}^*), $n + 1 \leq \text{card}(c(B))$. Let us denote $B'' = \langle B' \rangle_1 = \langle B' \rangle$ and $w_0 = \epsilon$. By induction (where $i = 1 \dots n$) and by the Lemma 9 applied on A and $\langle w_0 u_1 \dots w_{i-1} u_i \rangle v_i w_i u_{i+1}$ we can see that $|B''| \geq \max_{i=1}^{n+1} \{|w_{i-1} u_i|\}$ since after every application $xx \rightarrow x$ we can found each $w_{i-1} u_i$ as a subword in the residual of B . Hence we get that $|B''| \geq \max_{i=1}^{n+1} \{|w_{i-1} u_i|\} \geq \frac{1}{n+1} \sum_{i=1}^{n+1} |w_{i-1} u_i| = \frac{|B'|}{n+1}$ and from the fact $n + 1 \leq \text{card}(c(B)) = \text{card}(c(B''))$ we can deduce that $|\langle B' \rangle|^2 = |B''|^2 \geq \text{card}(c(B'')) \cdot |B''| \geq (n + 1) \frac{|B'|}{n+1} = |B'|$. \square

The previous proposition can be generalized in the following way.

Corollary 1. *Let A_1, B and A_2 be in normal forms and $\langle A_1 B A_2 \rangle_2 = A'_1 B' A'_2$ where A'_1, B', A'_2 are the residuals of A_1, B, A_2 . Then $|B'| \leq 2 \cdot |\langle B' \rangle|^2$.*

Proof. We prove the assertion for $A'_1 = A_1$ and $A'_2 = A_2$, because in the case when we leave out some occurrences of letters in A_1 and A_2 , we can reduce $A'_i \rightarrow_1^* \langle A'_i \rangle$ and start with these new surroundings $\langle A'_1 \rangle$ and $\langle A'_2 \rangle$ of the word B , since $\langle \langle A'_1 \rangle B \langle A'_2 \rangle \rangle_2 = \langle A'_1 \rangle B' \langle A'_2 \rangle$. So, $\langle A_1 B A_2 \rangle_2 = A_1 B' A_2$ where B' is the residual of B .

We will use the maximal reduction again and for an arbitrary word $B_i \in \mathcal{C}^*$ we denote B'_i its residual (after the applications of \rightarrow_{2m}). Three different cases must be discussed.

- 1) There is a reduction \rightarrow_{2m} using letters from both A_1 and A_2 .
- 2) There is a letter in B which is not involved any reduction \rightarrow_{2m} .
- 3) Otherwise.

In the case 1) we can write $B = B_1 v B_2$, $A_1 = s u_1$ and $A_2 = w_2 t$ where $s u_1 B_1 v B_2 w_2 t \rightarrow_{2m} A_1 B_1 B_2 A_2$. We apply twice the Proposition 1 on the words $A_1 B'_1$ and $B'_2 A_2$. We can deduce $|B'| = |B'_1 B'_2| = |B'_1| + |B'_2| \leq |\langle B'_1 \rangle|^2 + |\langle B'_2 \rangle|^2 \leq 2 \cdot |\langle B'_1 B'_2 \rangle|^2 = 2 \cdot |\langle B' \rangle|^2$ where the last inequality holds, because by the Remark 2 we have $\langle B'_1 \rangle \langle B'_2 \rangle \rightarrow_1 \langle B' \rangle$ (in the case when $\langle B'_1 \rangle \langle B'_2 \rangle$ contain a square) and so $|\langle B'_1 B'_2 \rangle| \geq \max\{|\langle B'_1 \rangle|, |\langle B'_2 \rangle|\}$, which implies that $2 \cdot |\langle B'_1 B'_2 \rangle|^2 \geq |\langle B'_1 \rangle|^2 + |\langle B'_2 \rangle|^2$.

In the case 2) we can write $B = B_1 B_2 B_3$ where B_2 is not involved in any reduction by \rightarrow_{2m} . Then we have $\langle B' \rangle = \langle B'_1 \rangle B_2 \langle B'_3 \rangle$ and so by the Proposition 1 we get $|B'| = |B'_1 B_2 B'_3| = |B'_1| + |B_2| + |B'_3| \leq |\langle B'_1 \rangle|^2 + |B_2| + |\langle B'_3 \rangle|^2 \leq |\langle B'_1 \rangle B_2 \langle B'_3 \rangle|^2 = |\langle B' \rangle|^2$.

In the case 3) we can write $B = B_1 v_1 B_2 B_3 B_4 v_2 B_5$, $A_1 = s u_1$, $A_2 = w_2 t$ where

$$A_1 B A_2 = s u_1 B_1 v_1 B_2 B_3 B_4 v_2 B_5 w_2 t \rightarrow_{2m} A_1 B_1 B_2 B_3 B_4 v_2 B_5 A_2,$$

$$A_1 B A_2 = s u_1 B_1 v_1 B_2 B_3 B_4 v_2 B_5 w_2 t \rightarrow_{2m} A_1 B_1 v_1 B_2 B_3 B_4 B_5 A_2$$

are the unique overlapping reductions. We have $B' = B'_1 B_2 B_3 B_4 B'_5$ and $B_0 = \langle B'_1 B_2 B_3 \rangle$ is a prefix of $\langle B' \rangle = \langle B_0 B_4 B'_5 \rangle$, and so $|B_0| \leq |\langle B' \rangle|$. We can see that $|B'| = |B'_1 B_2 B_3 B_4 B'_5| = |B'_1 B_2 B_3| + |B_4 B'_5| \leq |\langle B'_1 B_2 B_3 \rangle|^2 + |B_0 B_4 B'_5| \leq |B_0|^2 + |\langle B_0 B_4 B'_5 \rangle|^2 \leq 2 \cdot |\langle B' \rangle|^2$. \square

Lemma 10. *Let $sxxt$ be a word such that $\langle sxxt \rangle_2 = sxxt$ and $sxxt$ contains another square y^2 ($|y| \leq |x|$) such that one of these occurrences of y lies inside the x^2 . Then one of the following conditions holds:*

1. y is a suffix of s and a prefix of x
2. y is a prefix of t and a suffix of x
3. y^2 is a subword of x

Proof. Since y is a subword of x^2 , we have $c(y) \subseteq c(x)$ and let us suppose that conditions 1. and 2. do not hold. If $|x| = |y|$ then we get $c(x) = c(y)$ and we can apply \rightarrow_2 (w.l.o.g. $\underline{x}v\underline{y} \rightarrow_2 xy$ where v is both a prefix of x and a suffix of y), which is a contradiction. Assume that $|y| < |x|$. Notice that the first and the last

letter of x are unique occurrences of these constants in the word x because in another case we can apply \rightarrow_{2l} on x^2 . This implies that y^2 does not contain the first letter of the right x and y^2 also does not contain the last letter of the left x , constraining that y^2 is a subword of x . \square

Remark 3. The previous lemma shows that for two applications of the rules $xx \rightarrow_1 x$ and $yy \rightarrow_1 y$ on a word p in a normal form w.r.t. \rightarrow_2 , one of the following conditions holds (up to symmetry):

1. xx and yy do not overlap
2. yy is a subword of x
3. $x = x'z, y = zy'$ and $xx'zy'y$ is a subword of p

Lemma 11. *If $\langle sxxt \rangle_2 = sxxt$ and sxt contains a square y^2 which is not in $sxxt$ then $y = s_1xt_1$ where $|s_1|, |t_1| \geq 1$.*

Proof. Since y^2 is not in $sxxt$, we have $y^2 = s_0xt_0$ where s_0 is a suffix of s and t_0 is a prefix of t . If x is not a subword of y then $y = s_0x_1, y = x_2t_0$ where $x = x_1x_2$. Hence $s_0xxt_0 = \underline{s_0x_1x_2x_1x_2t_0} \rightarrow_2 s_0x_1x_2t_0$ since $c(x_2x_1) \subseteq c(y) = c(s_0x_1) = c(x_2t_0)$. The case when x_1 or x_2 is an empty word (i.e. x is a prefix or a suffix of y) is also included and we can conclude that $y = s_1xt_1$ and $|s_1|, |t_1| \geq 1$. \square

Proposition 2. *Let A and B be in normal forms, $\text{card}(c(AB)) \geq 2$ and $\langle AB \rangle_2 = AB$. Then $|AB| \leq |\langle AB \rangle|^2$.*

Proof. We denote k the length of $\langle AB \rangle_1 = \langle AB \rangle$. The case $k = 2$ is trivial. Now we assume that $k \geq 3$.

At first we have a look at the squares in AB . Since A and B are in normal forms, each square has got some letters from A and some from B . By the Remark 3 we have that AB contains at most two squares because the cases 1. and 2. of the Remark 3 are impossible.

By the Lemma 8, anytime during any reduction sequence by \rightarrow_1 the residuals of A and B remain in normal forms and their concatenations are in a normal form w.r.t. \rightarrow_2 . So by the previous arguments there are at most two squares and by the Lemma 11, if the reduction \rightarrow_1 introduces a new square then it is larger at least by two letters than the previous one. We use the sequence of reductions which in each step reduces the smallest square. Then the last reduction in this sequence reduces a square x^2 with the property $|x| \leq k$. The previous reduction reduces a square y^2 with the property $|y| \leq k - 1$ (the case $|y| = |x| = k = |\langle AB \rangle|$ is a contradiction with AB is in normal form w.r.t. \rightarrow_2). Thus we can see that for the length of AB we have an upper bound $k + k + (k - 1) + (k - 2) + \dots \leq k^2$. \square

Proposition 3. *There is a polynomial $p : \mathbb{N} \rightarrow \mathbb{N}$, such that for an arbitrary $A_1, B, A_2 \in \mathcal{C}^*$ in normal forms and $\langle A_1BA_2 \rangle_2 = A'_1B'A'_2$ where $A'_i, 1 \leq i \leq 2$, is the residual of A_i and B' is the residual of B , we have $|B'| \leq p(|\langle A_1BA_2 \rangle|)$.*

Proof. We may assume that $\text{card}(c(B)) \geq 2$. By the Corollary 1 we know that $|B'| \leq 2 \cdot |\langle B' \rangle|^2$, which is of course less or equal to $2 \cdot |\langle A'_1 \rangle \langle B' \rangle|^2$. Since

$\langle\langle A'_1 \rangle\langle B' \rangle\rangle_2 = \langle A'_1 \rangle\langle B' \rangle$ by the Lemma 8, we can use the Proposition 2 and we get that $2 \cdot |\langle A'_1 \rangle\langle B' \rangle|^2 \leq 2 \cdot |\langle A'_1 B' \rangle|^4$, which is again less or equal to $2 \cdot |\langle A'_1 B' \rangle\langle A'_2 \rangle|^4$. Analogously we have that $2 \cdot |\langle A'_1 B' \rangle\langle A'_2 \rangle|^4 \leq 2 \cdot |\langle A'_1 B' A'_2 \rangle|^8$. Thus we have $|B'| \leq p(|\langle A_1 B A_2 \rangle|)$ for the polynomial $p(n) = 2 \cdot n^8$. \square

Proposition 4. *Let p be a polynomial that satisfies the condition from the Proposition 3. If a stuttering pattern equation system $\{X_1 = A_1, \dots, X_n = A_n\}$ is satisfiable then there exists a solution α with $\text{size}(\alpha) \leq \sum_{i=1}^n |X_i| \cdot p(|A_i|)$.*

Proof. Of course, we can assume that all A_i 's are in normal forms. Let α be a solution of the stuttering pattern equation system $\{X_1 = A_1, \dots, X_n = A_n\}$ which minimizes both $\text{size}(\alpha)$ and the number of variables x such that $|\alpha(x)| = \text{size}(\alpha)$. Assume for the moment that there is some x such that $\text{size}(\alpha) = |\alpha(x)| > \sum_{i=1}^n |X_i| \cdot p(|A_i|)$. We may assume that $\alpha(x)$ is in normal form, otherwise we have a smaller solution.

We now reduce $\alpha(X_i) \rightarrow_{2m}^* \langle \alpha(X_i) \rangle_2$. If we look at an arbitrary residual B' of an occurrence of $\alpha(x)$ in $\langle \alpha(X_i) \rangle_2$, we see that $|B'| \leq p(|A_i|)$ by the Proposition 3. This means that there are at most $\sum_{i=1}^n |X_i| \cdot p(|A_i|)$ letter's occurrences in the residuals of all occurrences of $\alpha(x)$ in all $\langle \alpha(X_i) \rangle_2$. By the assumption $|\alpha(x)| > \sum_{i=1}^n |X_i| \cdot p(|A_i|)$ we get that there is an occurrence of a letter a in $\alpha(x)$ that has been left out from all the occurrences of $\alpha(x)$ by the rule \rightarrow_{2m} . We can erase this occurrence of the letter a from $\alpha(x)$ and we get a smaller solution, which is a contradiction. \square

The previous considerations lead to the following corollary.

Corollary 2. *The PATTERN-EQUATION problem is in NP.*

Proof. We can guess the solution α and by the Proposition 4, if the system is satisfiable then there is a solution of a polynomial length. Checking whether α solves all the equations takes also a polynomial time and so the problem is in NP. \square

5 NP-hardness of the PATTERN-EQUATION problem

In this section we show that the PATTERN-EQUATION problem in a free idempotent semigroup is NP-hard. We use a reduction from the NP-complete problem 3-SAT (see [11]).

Proposition 5. *The PATTERN-EQUATION problem is NP-hard.*

Proof. Suppose we have an instance of 3-SAT, i.e.

$$C \equiv C_1 \wedge C_2 \wedge \dots \wedge C_n$$

is a conjunction of clauses and each clause C_i , $1 \leq i \leq n$, is of the form

$$l_1 \vee l_2 \vee l_3$$

where l_j , $1 \leq j \leq 3$, is a literal (l_j is a variable from the set Var , eventually negated – we call it positive resp. negative literal). A valuation is a mapping $v : \text{Var} \rightarrow \{T, F\}$. This valuation extends naturally to C and we say that C is satisfiable if and only if there exists a valuation v such that $v(C) = T$.

We construct a stuttering pattern equation system such that the system is satisfiable if and only if C is satisfiable. The system will consist of the following equations (1) – (6) and $\mathcal{C} = \{a, b, c\}$, $\mathcal{V} = \{x, s_1^x, t_1^x, s_2^x, t_2^x \mid x \in \text{Var} \cup \overline{\text{Var}}\} \cup \{y_a, y_b, y_c\}$ where $\overline{\text{Var}} = \{\bar{x} \mid x \in \text{Var}\}$ is a disjunct copy of Var .

$$y_a = a, \quad y_b = b, \quad y_c = c \quad (1)$$

We define $\tilde{x} = x$ if x is a positive literal, $\widetilde{\bar{x}} = \bar{x}$ if $\neg x$ is a negative literal and for all clauses $C_i \equiv l_1 \vee l_2 \vee l_3$ we have the equation

$$y_a \tilde{l}_1 \widetilde{\tilde{l}_2} \widetilde{\tilde{l}_3} y_a = aba \quad (2)$$

for each $x \in \text{Var}$ we add the equations

$$y_b x \bar{x} y_b = bab \quad (3)$$

$$y_a x \bar{x} y_a = aba \quad (4)$$

and finally for each $x \in \text{Var} \cup \overline{\text{Var}}$ we have the following equations.

$$s_1^x x t_1^x = acb, \quad s_1^x y_c = ac \quad (5)$$

$$s_2^x x t_2^x = bca, \quad s_2^x y_c = bc \quad (6)$$

The intuition behind the construction is following. If a variable x is true then $x = b$ and if x is false then $x = a$.

Suppose that C is satisfiable, i.e. there is a valuation v such that $v(C) = T$. Then we show that α defined below is a solution of our system. Let us state

$$\alpha(y_a) = a, \quad \alpha(y_b) = b, \quad \alpha(y_c) = c$$

and for all $x \in \text{Var}$ such that $v(x) = T$ let

$$\alpha(x) = b, \quad \alpha(\bar{x}) = a$$

$$\alpha(s_1^x) = ac, \quad \alpha(t_1^x) = b, \quad \alpha(s_1^{\bar{x}}) = a, \quad \alpha(t_1^{\bar{x}}) = cb$$

$$\alpha(s_2^x) = b, \quad \alpha(t_2^x) = ca, \quad \alpha(s_2^{\bar{x}}) = bc, \quad \alpha(t_2^{\bar{x}}) = a$$

and if $v(x) = F$ then

$$\alpha(x) = a, \quad \alpha(\bar{x}) = b$$

$$\alpha(s_1^x) = a, \quad \alpha(t_1^x) = cb, \quad \alpha(s_1^{\bar{x}}) = ac, \quad \alpha(t_1^{\bar{x}}) = b$$

$$\alpha(s_2^x) = bc, \quad \alpha(t_2^x) = a, \quad \alpha(s_2^{\bar{x}}) = b, \quad \alpha(t_2^{\bar{x}}) = ca.$$

Checking that α is a solution (even non-singular) is a routine. The only interesting equation is (2). This equation is also satisfied by α since we have the assumption that under the valuation v there is at least one true literal in each clause.

Let us suppose that α is an arbitrary solution of our system and we find a valuation that satisfies C . The equation (3) constraints that $c(\alpha(x)) \subseteq \{a, b\}$ for all $x \in \text{Var} \cup \overline{\text{Var}}$. We will conclude that it is not possible that $c(\alpha(x)) = \{a, b\}$.

Suppose that it is the case and using the equations (5) we get that $\alpha(x)$ does not begin with the constant a . For the moment assume that $\alpha(x)$ begins with a . We have $ac = 0(acb) \sim 0(\alpha(s_1^x x t_1^x))$. If $c(\alpha(s_1^x)) = \{a\}$ then $c(0(\alpha(s_1^x x t_1^x))) = \{a, b\}$ whereas $c(0(acb)) = \{a, c\}$, which is a contradiction. Otherwise we have $c(\alpha(s_1^x)) = \{a, c\}$ (thanks to the equation $s_1^x y_c = ac$) and we get $0(\alpha(s_1^x x t_1^x)) \sim aca \not\sim ac \sim 0(acb)$.

By the similar arguments and using the equations (6) we get that $\alpha(x)$ does not begin with the constant b . This yields that there are just three possibilities for $\alpha(x)$, namely $\alpha(x) = a$, $\alpha(x) = b$ or $\alpha(x) = \epsilon$.

By the equations (3) and (1) we know that for all $x \in \text{Var}$ at least $\alpha(x) = a$ or $\alpha(\bar{x}) = a$. The equation (4) constraints that either $\alpha(x) = b$ or $\alpha(\bar{x}) = b$. Similarly for each clause, the equation (2) with (1) gives that there is j , $1 \leq j \leq 3$, such that $\alpha(\tilde{l}_j) = b$. Let us finally define the valuation v as $v(x) = T$ if $\alpha(x) = b$ and $v(x) = F$ if $\alpha(x) = a$ for each $x \in \text{Var}$. The valuation is consistent and it holds that $v(C) = T$.

This is enough to demonstrate that the PATTERN-EQUATION problem is NP-hard since the reduction can be done in a polynomial time and is effectively constructable. \square

It is not difficult to see that the same reduction as above would also work for the NON-SINGULAR-PATTERN-EQUATION problem, which is consequently also NP-hard. We can now formulate the main result of this paper.

Theorem 1. *The PATTERN-EQUATION problem and NON-SINGULAR-PATTERN-EQUATION problem are NP-complete.*

Acknowledgements We would like to thank Ivana Černá and Michal Kunc for their comments and suggestions.

References

- [1] Book R., Otto F.: String-Rewriting Systems (1993) Springer-Verlag
- [2] Brim L., Gilbert D.R., Jacquet J-M, Křetínský M.: A Process Algebra for Synchronous Concurrent Constraint Programming, Proceedings of ALP96: Fifth International Conference on Algebraic and Logic Programming, Springer LNCS volume 1139 (1996) 165–178, Springer-Verlag.
- [3] Černá I., Klíma O., Srba J.: Pattern Equations and Equations with Stuttering, To appear in Proceedings of SOFSEM'99, the 26th Seminar on Current Trends in Theory and Practice of Informatics (1999) Springer-Verlag.

- [4] Green J.A., Rees D.: On semigroups in which $x^r = x$, Proc. Camb. Phil. Soc. **48** (1952) 35–40.
- [5] Kopeček I.: Automatic Segmentation into Syllable Segments, Proceedings of First International Conference on Language Resources and Evaluation (1998) 1275–1279.
- [6] Kopeček I., Pala K.: Prosody Modelling for Syllable-Based Speech Synthesis, Proceedings of the IASTED International Conference on Artificial Intelligence and Soft Computing, Cancun (1998) 134–137.
- [7] Lothaire M.: Algebraic Combinatorics on Words, Preliminary version available at <http://www-igm.univ-mlv.fr/~berstel/Lothaire/index.html>
- [8] Lothaire, M.: Combinatorics on Words, Volume **17** of Encyclopedia of Mathematics and its Applications (1983) Addison-Wesley.
- [9] Kađourek J., Polák L.: On free semigroups satisfying $x^r = x$, Simon Stevin **64**, No.1 (1990) 3–19.
- [10] Makanin, G. S.: The Problem of Solvability of Equations in a Free Semigroup, Mat. Sbornik. **103(2)** (1977) 147–236. (In Russian) English translation in: Math. USSR Sbornik **32** (1977) 129–198.
- [11] Papadimitriou, C.H.: Computational Complexity, Addison-Wesley Publishing Company (1994), Reading, Mass.
- [12] Perrin D.: Equations in Words, In H. Ait-Kaci and M. Nivat, editors, Resolution of Equations in Algebraic Structures, Vol. **2** (1989) 275–298, Academic Press.
- [13] Schulz, K. U.: Makanin's Algorithm for Word Equations: Two Improvements and a Generalization, In Schulz, K.–U. (Ed.), Proceedings of Word Equations and Related Topics, 1st International Workshop, IWW-ERT'90, Tübingen, Germany, Vol. **572** of LNCS (1992) 85–150, Berlin-Heidelberg-New York, Springer Verlag.
- [14] Siekmann J., Szabó P.: A Noetherian and Confluent Rewrite System for Idempotent Semigroups, Semigroup Forum **25** (1982)

**Copyright © 1999, Faculty of Informatics, Masaryk University.
All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**

**Publications in the FI MU Report Series are in general accessible
via WWW and anonymous FTP:**

`http://www.fi.muni.cz/informatics/reports/
ftp ftp.fi.muni.cz (cd pub/reports)`

Copies may be also obtained by contacting:

**Faculty of Informatics
Masaryk University
Botanická 68a
602 00 Brno
Czech Republic**