

Achim Blumensath
blumens@fi.muni.cz

This document was last updated 2018-07-15.
The latest version can be found at

www.fi.muni.cz/~blumens

COPYRIGHT 2018 Achim Blumensath



This work is licensed under the *Creative Commons Attribution 4.0 International License*. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Contents

A. Set Theory	1
<i>A1 Basic set theory</i>	3
1 Sets and classes	3
2 Stages and histories	11
3 The cumulative hierarchy	18
<i>A2 Relations</i>	27
1 Relations and functions	27
2 Products and unions	36
3 Graphs and partial orders	39
4 Fixed points and closure operators	47
<i>A3 Ordinals</i>	57
1 Well-orders	57
2 Ordinals	64
3 Induction and fixed points	74
4 Ordinal arithmetic	85
<i>A4 Zermelo-Fraenkel set theory</i>	105
1 The Axiom of Choice	105
2 Cardinals	113
3 Cardinal arithmetic	116
4 Cofinality	122

5	The Axiom of Replacement	131
6	Stationary sets	134
7	Conclusion	145
B. General Algebra		147
<i>B1 Structures and homomorphisms</i>		149
1	Structures	149
2	Homomorphisms	156
3	Categories	162
4	Congruences and quotients	175
<i>B2 Trees and lattices</i>		189
1	Trees	189
2	Lattices	197
3	Ideals and filters	205
4	Prime ideals and ultrafilters	209
5	Atomic lattices and partition rank	217
<i>B3 Universal constructions</i>		231
1	Terms and term algebras	231
2	Direct and reduced products	242
3	Directed limits and colimits	250
4	Equivalent diagrams	262
5	Links and dense functors	275
<i>B4 Accessible categories</i>		289
1	Filtered limits and inductive completions	289
2	Extensions of diagrams	304
3	Presentable objects	320
4	Accessible categories	333

<i>B5 Topology</i>	345	
1	Open and closed sets	345
2	Continuous functions	350
3	Hausdorff spaces and compactness	354
4	The Product topology	361
5	Dense sets and isolated points	365
6	Spectra and Stone duality	374
7	Stone spaces and Cantor-Bendixson rank	381
<i>B6 Classical Algebra</i>		389
1	Groups	389
2	Group actions	393
3	Rings	401
4	Modules	407
5	Fields	414
6	Ordered fields	429
C. First-Order Logic and its Extensions		445
<i>C1 First-order logic</i>		447
1	Infinitary first-order logic	447
2	Axiomatisations	458
3	Theories	464
4	Normal forms	469
5	Translations	476
6	Extensions of first-order logic	485
<i>C2 Elementary substructures and embeddings</i>		497
1	Homomorphisms and embeddings	497
2	Elementary embeddings	502
3	The Theorem of Löwenheim and Skolem	508

4 The Compactness Theorem 515

5 Amalgamation 525

c3 *Types and type spaces* 531

1 Types 531

2 Type spaces 537

3 Retracts 550

4 Local type spaces 561

5 Stable theories 566

c4 *Back-and-forth equivalence* 581

1 Partial isomorphisms 581

2 Hintikka formulae 590

3 Ehrenfeucht-Fraïssé games 593

4 κ -complete back-and-forth systems 602

5 The theorems of Hanf and Gaifman 609

c5 *General model theory* 617

1 Classifying logical systems 617

2 Hanf and Löwenheim numbers 621

3 The Theorem of Lindström 628

4 Projective classes 640

5 Interpolation 651

6 Fixed-point logics 661

D. Axiomatisation and Definability 687

D1 *Quantifier elimination* 689

1 Preservation theorems 689

2 Quantifier elimination 693

3 Existentially closed structures 703

4 Abelian groups 709

5 Fields 714

D2 *Products and varieties* 721

1 Ultraproducts 721

2 The theorem of Keisler and Shelah 726

3 Reduced products and Horn formulae 739

4 Quasivarieties 744

5 The Theorem of Feferman and Vaught 756

D3 *O-minimal structures* 761

1 Ordered topological structures 761

2 O-minimal groups and rings 767

3 Cell decompositions 769

E. Classical Model Theory 789

E1 *Saturation* 791

1 Homogeneous structures 791

2 Saturated structures 797

3 Projectively saturated structures 808

4 Pseudo-saturated structures 811

E2 *Definability and automorphisms* 819

1 Definability in projectively saturated models 819

2 Imaginary elements and canonical parameters 830

3 Galois bases 838

4 Elimination of imaginaries 844

5 Weak elimination of imaginaries 850

E3 Prime models 859

1 Isolated types 859

2 The Omitting Types Theorem 861

3 Prime and atomic models 869

4 Constructible models 873

E4 \aleph_0 -categorical theories 881

1 \aleph_0 -categorical theories and automorphisms 881

2 Back-and-forth arguments in accessible categories 897

3 Fraïssé limits 909

4 Zero-one laws 921

E5 Indiscernible sequences 929

1 Ramsey Theory 929

2 Ramsey Theory for trees 934

3 Indiscernible sequences 945

4 The independence and strict order properties 956

E6 Functors and embeddings 969

1 Local functors 969

2 Word constructions 976

3 Ehrenfeucht-Mostowski models 985

E7 Abstract elementary classes 999

1 Abstract elementary classes 999

2 Amalgamation and saturation 1008

3 Limits of chains 1021

4 Categoricity and stability 1025

F. Independence and Forking 1033

F1 Geometries 1035

1 Dependence relations 1035

2 Matroids and geometries 1040

3 Modular geometries 1047

4 Strongly minimal sets 1053

5 Vaughtian pairs and the Theorem of Morley 1061

F2 Ranks and forking 1073

1 Morley rank and Δ -rank 1073

2 Independence relations 1087

3 Preforking relations 1101

4 Forking relations 1118

F3 Simple theories 1131

1 Dividing and forking 1131

2 Simple theories and the tree property 1141

F4 Theories without the independence property 1159

1 Honest definitions 1159

2 Lascar invariant types 1174

3 \sqrt{i} -Morley sequences 1200

4 Dp-rank 1213

F5 Theories without the array property 1225

1 The array property 1225

2 Forking and dividing 1234

3 The Independence Theorem 1253

G. Geometric Model Theory 1267

Contents

<i>G1 Stable theories</i>	1269
1 Definable types	1269
2 Forking in stable theories	1274
3 Stationary types	1278
4 The multiplicity of a type	1284
5 Morley sequences in stable theories	1291
6 The stability spectrum	1296
<i>G2 Models of stable theories</i>	1303
1 Isolation relations	1303
2 Constructions	1312
3 Prime models	1320
4 $\text{at}\sqrt{\text{}}$ -constructible models	1325
5 Strongly independent stratifications	1334
6 Representations	1343
<i>Recommended Literature</i>	1355
<i>Symbol Index</i>	1357
<i>Index</i>	1369

Part A.
Set Theory

f5. Theories without the array property

(EXT) Suppose that $\bar{a} \downarrow_U^\circ B_0$ and let $B_0 \subseteq B_1$. Using the extension axioms, we can find a tuple \bar{a}' such that

$$\text{atp}(\bar{a}'/UB_0) = \text{atp}(\bar{a}/UB_0) \quad \text{and} \quad (\bar{a}' \setminus U) \cap B_1 = \emptyset.$$

By ultrahomogeneity, there exists an automorphism $\pi \in \text{Aut } \mathbb{M}_{UB_0}$ mapping \bar{a} to \bar{a}' . Hence, $\bar{a}' \equiv_{UB_0} \bar{a}$ and $\bar{a}' \downarrow_U^\circ B_1$.

(IND) We prove that \downarrow° satisfies the Independence Theorem for the class of all subsets of \mathbb{M} . Suppose that

$$\bar{a} \equiv_U \bar{b}, \quad \bar{a} \downarrow_U^\circ A, \quad \bar{b} \downarrow_U^\circ B, \quad \text{and} \quad A \downarrow_U^\circ B.$$

Replacing A and B by, respectively, $A \setminus U$ and $B \setminus U$, we may assume that $A \cap U = \emptyset$ and $B \cap U = \emptyset$. Let

$$\bar{d} := \bar{a} \cap U, \quad \bar{a}' := \bar{a} \setminus U, \quad \text{and} \quad \bar{b}' := \bar{b} \setminus U.$$

Note that $\bar{a}' \cap (U \cup A) = \emptyset$ and $\bar{b}' \cap (U \cup B) = \emptyset$. Since U, A, B are disjoint, we can use the extension axioms to find a tuple \bar{c}' disjoint from $U \cup A \cup B$ such that

$$\text{atp}(\bar{c}'/UA) = \text{atp}(\bar{a}'/UA) \quad \text{and} \quad \text{atp}(\bar{c}'/UB) = \text{atp}(\bar{b}'/UB).$$

It follows that

$$\bar{c}'\bar{d} \equiv_{UA} \bar{a}'\bar{d}, \quad \bar{c}'\bar{d} \equiv_{UB} \bar{b}'\bar{d}, \quad \text{and} \quad \bar{c}'\bar{d} \downarrow_U^\circ AB. \quad \square$$

Part G.

Geometric Model Theory

G1. Stable theories

1. Definable types

A key property of stable theories is the definability of types. The relation $\overset{\text{df}}{\vee}$ will thus play a major role in this chapter. In the next section, we will study its properties in the context of stable theories. But first, we consider the relation $\overset{\text{df}}{\vee}$ in an arbitrary theory, where it is usually much less well-behaved: if U and B are small enough it might happen that $\text{tp}(\bar{a}/UB)$ is definable over U just because some formula ‘accidentally’ is a φ -definition, although it ceases to be a definition for every extension $\text{tp}(\bar{a}/UB')$ with $B' \supseteq B$. Therefore, when investigating a statement of the form $A \overset{\text{df}}{\vee} B$ we usually assume that one of the sets B and U is large. In particular, there is hope that the derived relation $^*(\overset{\text{df}}{\vee})$ is much better behaved. We start with relating $\overset{\text{df}}{\vee}$ to the relation $\overset{\forall}{\vee}$.

Lemma 1.1. *Let $A, A', B, U \subseteq \mathbb{M}$.*

- (a) $A \overset{\text{df}}{\vee} B$ and $B \overset{\forall}{\vee} U$ implies $B \overset{\forall}{\vee} A$.
- (b) $A \overset{\text{df}}{\vee} B$, $B \overset{\forall}{\vee} A'$, and $A \equiv_U A'$ implies $A \equiv_{UB} A'$.

Proof. (a) Let $A \overset{\text{df}}{\vee} B$ and $B \overset{\forall}{\vee} U$. To show that $B \overset{\forall}{\vee} A$, suppose that $\mathbb{M} \models \varphi(\bar{a}; \bar{b})$, where $\bar{a} \subseteq A$, $\bar{b} \subseteq B$, and $\varphi(\bar{x}; \bar{y})$ is a formula over U . We have to find a tuple $\bar{c} \subseteq U$ such that $\mathbb{M} \models \varphi(\bar{a}; \bar{c})$. Since $\bar{a} \overset{\text{df}}{\vee} \bar{b}$, the type $\text{tp}(\bar{a}/U\bar{b})$ has a φ -definition δ over U . Hence,

$$\mathbb{M} \models \varphi(\bar{a}; \bar{b}) \text{ implies } \mathbb{M} \models \delta(\bar{b}).$$

Since $\bar{b} \overset{\forall}{\vee} U$, there is some $\bar{c} \subseteq U$ with $\mathbb{M} \models \delta(\bar{c})$. Consequently, $\mathbb{M} \models \varphi(\bar{a}; \bar{c})$.

(b) Let \bar{a} be an enumeration of A and \bar{a}' the corresponding enumeration of A' . For every formula $\varphi(\bar{x}; \bar{y})$ over U , we fix a φ -definition $\delta_\varphi(\bar{y})$ of $\text{tp}(\bar{a}/UB)$ over U . It is sufficient to prove that δ_φ is also a φ -definition of $\text{tp}(\bar{a}'/UB)$ since this implies that

$$\mathbb{M} \models \varphi(\bar{a}; \bar{b}) \quad \text{iff} \quad \mathbb{M} \models \delta(\bar{b}) \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}'; \bar{b}),$$

for all $\bar{b} \subseteq U \cup B$.

For a contradiction, suppose that the formula δ_φ is not a φ -definition of $\text{tp}(\bar{a}'/UB)$. Then there exists a tuple $\bar{b} \subseteq U \cup B$ such that

$$\mathbb{M} \models \neg(\varphi(\bar{a}'; \bar{b}) \leftrightarrow \delta_\varphi(\bar{b})).$$

Since $B \forall_U \bar{a}'$, there is a tuple $\bar{c} \subseteq U$ such that

$$\mathbb{M} \models \neg(\varphi(\bar{a}'; \bar{c}) \leftrightarrow \delta_\varphi(\bar{c})).$$

As $\bar{a} \equiv_U \bar{a}'$, this implies that

$$\mathbb{M} \models \neg(\varphi(\bar{a}; \bar{c}) \leftrightarrow \delta_\varphi(\bar{c})).$$

Consequently, δ_φ is no φ -definition of $\text{tp}(\bar{a}/UB)$. A contradiction. \square

The relation $\overset{\text{df}}{\forall}$ is particularly well-behaved if the base set is a model.

Lemma 1.2. *Let T be a complete first-order theory, \mathfrak{M} a model of T , and $A \subseteq \mathbb{M}$ a set such that $A \overset{\text{df}}{\forall}_M M$.*

(a) $A \overset{*}{\overset{\text{df}}{\forall}}_M M$.

(b) For every set $B \subseteq \mathbb{M}$, there exists a set $A' \equiv_M A$ such that $B \overset{\forall}{\forall}_M A'$.

(c) If $A', B \subseteq \mathbb{M}$ are sets such that

$$B \overset{\forall}{\forall}_M A, \quad B \overset{\forall}{\forall}_M A' \quad \text{and} \quad A \equiv_M A', \quad \text{then} \quad A \equiv_{\mathbb{M}} \overset{\forall}{\forall}_M A'.$$

Proof. (a) Suppose that $\bar{a} \overset{\text{df}}{\forall}_M M$. To show that $\bar{a} \overset{*}{\overset{\text{df}}{\forall}}_M M$, consider a set $C \supseteq M$. For every formula $\varphi(\bar{x}; \bar{y})$ fix a φ -definition $\delta_\varphi(\bar{y})$ of $\text{tp}(\bar{a}/M)$ over M and set

$$\Phi(\bar{x}) := \{ \varphi(\bar{x}; \bar{c}) \mid \bar{c} \subseteq C, \mathbb{M} \models \delta_\varphi(\bar{c}) \}.$$

Note that $\text{tp}(\bar{a}/M) \subseteq \Phi$. Hence, if \bar{a}' is a tuple satisfying Φ , then $\bar{a}' \equiv_M \bar{a}$ and $\bar{a}' \overset{\text{df}}{\forall}_M C$ since each formula δ_φ is a φ -definition of $\text{tp}(\bar{a}'/C)$.

Thus, it remains to prove that Φ is satisfiable. Consider a finite subset $\Phi_o = \{ \varphi_o(\bar{x}; \bar{c}_o), \dots, \varphi_n(\bar{x}; \bar{c}_n) \} \subseteq \Phi$. By definition of Φ we have

$$\mathbb{M} \models \delta_{\varphi_o}(\bar{c}_o) \wedge \dots \wedge \delta_{\varphi_n}(\bar{c}_n).$$

We have seen in Lemma F2.3.15 that $C \overset{\forall}{\forall}_M M$. Hence, we can find tuples $\bar{b}_o, \dots, \bar{b}_n \subseteq M$ such that

$$\mathbb{M} \models \delta_{\varphi_o}(\bar{b}_o) \wedge \dots \wedge \delta_{\varphi_n}(\bar{b}_n).$$

This implies that

$$\mathbb{M} \models \varphi_o(\bar{a}; \bar{b}_o) \wedge \dots \wedge \varphi_n(\bar{a}; \bar{b}_n).$$

Consequently, the model \mathbb{M} satisfies Φ_o if we interpret the variables \bar{x} by \bar{a} and the constants \bar{c}_i by \bar{b}_i .

(b) Given $B \subseteq \mathbb{M}$, we can use (a) to find a set $A' \equiv_M A$ such that $A' \overset{\text{df}}{\forall}_M B$. Furthermore, we have $B \overset{\forall}{\forall}_M M$, by Lemma F2.3.15. Hence, Lemma 1.1 (a) implies that $B \overset{\forall}{\forall}_M A'$.

(c) Let \bar{a} be an enumeration of A and let \bar{a}' be the corresponding enumeration of A' . By (a), we can find a tuple $\bar{a}'' \equiv_M \bar{a}$ such that $\bar{a}'' \overset{\text{df}}{\forall}_M B$. By Lemma 1.1 (b), $B \overset{\forall}{\forall}_M \bar{a}$ implies that $\bar{a}'' \equiv_{MB} \bar{a}$. Since $\bar{a}'' \equiv_M \bar{a}'$ and $B \overset{\forall}{\forall}_M \bar{a}'$, it follows in the same way that $\bar{a}'' \equiv_{MB} \bar{a}'$. Hence, $\bar{a} \equiv_{MB} \bar{a}'' \equiv_{MB} \bar{a}'$. \square

Lemma 1.3 (Harrington). *Let \mathfrak{M} be an \aleph_o -saturated model, $\mathfrak{p}(\bar{x}), \mathfrak{q}(\bar{y}) \in S^{<\omega}(M)$, and let $\varphi(\bar{x}; \bar{y})$ be a formula over M that does not have the order property. If $\delta(\bar{y})$ is a φ -definition of \mathfrak{p} and $\varepsilon(\bar{x})$ a φ -definition of \mathfrak{q} , then*

$$\varepsilon(\bar{x}) \in \mathfrak{p}(\bar{x}) \quad \text{iff} \quad \delta(\bar{y}) \in \mathfrak{q}(\bar{y}).$$

Proof. Let $\bar{c} \subseteq M$ be the parameters occurring in φ . By induction on n , we construct two sequences $(\bar{a}_n)_{n < \omega}$ and $(\bar{b}_n)_{n < \omega}$ in M as follows. Suppose that we have already defined $\bar{a}_0, \dots, \bar{a}_{n-1}$ and $\bar{b}_0, \dots, \bar{b}_{n-1}$. As \mathfrak{M} is \aleph_0 -saturated, we can choose a tuple $\bar{b}_n \subseteq M$ realising $q \upharpoonright \bar{c}\bar{a}_0 \dots \bar{a}_{n-1}$ and a tuple $\bar{a}_n \subseteq M$ realising $p \upharpoonright \bar{c}\bar{b}_0 \dots \bar{b}_n$.

Having constructed $(\bar{a}_n)_{n < \omega}$ and $(\bar{b}_n)_{n < \omega}$ it follows, for $i < k$, that

$$\begin{aligned} \mathfrak{M} \models \varphi(\bar{a}_i; \bar{b}_k) & \text{ iff } \varphi(\bar{a}_i; \bar{y}) \in q \upharpoonright \bar{c}\bar{a}_0 \dots \bar{a}_{k-1} \\ & \text{ iff } \mathfrak{M} \models \varepsilon(\bar{a}_i) \\ & \text{ iff } \varepsilon(\bar{x}) \in p, \end{aligned}$$

and, for $i \geq k$,

$$\begin{aligned} \mathfrak{M} \models \varphi(\bar{a}_i; \bar{b}_k) & \text{ iff } \varphi(\bar{x}; \bar{b}_k) \in p \upharpoonright \bar{c}\bar{b}_0 \dots \bar{b}_i \\ & \text{ iff } \mathfrak{M} \models \delta(\bar{b}_k) \\ & \text{ iff } \delta(\bar{x}) \in q. \end{aligned}$$

Hence, if $\delta \notin q$ and $\varepsilon \in p$, then

$$\mathfrak{M} \models \varphi(\bar{a}_i; \bar{b}_k) \quad \text{iff} \quad i < k,$$

and φ has the order property. A contradiction. In the same way we obtain a contradiction if we assume that $\delta \in q$ and $\varepsilon \notin p$. \square

We have already seen in Lemma F2.3.3 (a) that $\overset{\text{df}}{\surd} \subseteq \overset{s}{\surd}$. The converse holds only in special circumstances.

Lemma 1.4. *Suppose that \mathfrak{M} is a κ -saturated and strongly κ -homogeneous model and let $U \subseteq M$ be a set of size $|U| < \kappa$. If $A \subseteq \mathbb{M}$ is a set such that $A \overset{\text{df}}{\surd}_M M$, then*

$$A \overset{\text{df}}{\surd}_U M \quad \text{iff} \quad A \overset{s}{\surd}_U M.$$

Proof. (\Rightarrow) We have seen in Lemma F2.3.3 (a) that $\overset{\text{df}}{\surd} \subseteq \overset{s}{\surd}$.

(\Leftarrow) Let $\bar{a} \subseteq A$, let $\varphi(\bar{x}; \bar{y})$ be a formula, and let $\delta(\bar{y})$ be a φ -definition of $\text{tp}(\bar{a}/M)$ over M . It is sufficient to show that the relation $\delta^{\mathfrak{M}}$ is definable over U . By definition of $\overset{s}{\surd}$,

$$\bar{b} \equiv_U \bar{b}' \quad \text{implies} \quad \bar{b} \equiv_{U\bar{a}} \bar{b}', \quad \text{for all } \bar{b}, \bar{b}' \subseteq M.$$

Hence, if $\bar{b}, \bar{b}' \subseteq M$ are tuples such that $\bar{b} \equiv_U \bar{b}'$, then

$$\begin{aligned} \mathbb{M} \models \delta(\bar{b}) & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}; \bar{b}) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}; \bar{b}') \quad \text{iff} \quad \mathbb{M} \models \delta(\bar{b}'). \end{aligned}$$

Consequently, we have

$$\mathfrak{M} \models \delta(\bar{b}) \quad \text{iff} \quad \mathfrak{M} \models \delta(\pi(\bar{b})), \quad \text{for all } \pi \in \text{Aut } \mathfrak{M}_U,$$

and it follows by Lemma E2.1.10 that $\delta^{\mathfrak{M}}$ is definable over U . \square

Another immediate consequence of the inclusion $\overset{\text{df}}{\surd} \subseteq \overset{s}{\surd}$ is the corresponding inclusion between the starred relations.

Proposition 1.5. $*(\overset{\text{df}}{\surd}) \subseteq \overset{i}{\surd}$

Proof. We have seen in Lemma F2.3.3 (a) that $\overset{\text{df}}{\surd} \subseteq \overset{s}{\surd}$. This implies that $*(\overset{\text{df}}{\surd}) \subseteq *(\overset{s}{\surd}) = \overset{i}{\surd}$. \square

We conclude this section with a comparison of $\overset{\text{df}}{\surd}$ with $\overset{d}{\surd}$.

Lemma 1.6. *Let \mathfrak{M} be a κ -saturated model and let $U \subseteq M$ be a set of size $|U| < \kappa$. Then*

$$A \overset{\text{df}}{\surd}_U M \quad \text{implies} \quad A \overset{d}{\surd}_U M, \quad \text{for all } A \subseteq \mathbb{M}.$$

Proof. Suppose that $\bar{a} \overset{\text{df}}{\surd}_U M$. To show that $\bar{a} \overset{d}{\surd}_U M$ it is sufficient, by Lemma F3.1.3 and (DEF), to prove that, for every indiscernible sequence

$(\bar{b}_n)_{n < \omega}$ over U with $\bar{b}_0 \subseteq M$ and $|\bar{b}_0| < \aleph_0$, there exists a tuple $\bar{a}' \equiv_{U\bar{b}_0} \bar{a}$ such that

$$\bar{b}_m \equiv_{U\bar{a}'} \bar{b}_n, \quad \text{for all } m, n < \omega.$$

Hence, let $(\bar{b}_n)_{n < \omega}$ be such a sequence. As M is κ -saturated, we can find tuples $\bar{c}_n \subseteq M$, $n < \omega$, such that

$$(\bar{c}_n)_n \equiv_{U\bar{b}_0} (\bar{b}_n)_n.$$

Note that, by Lemma F2.3.3 (a),

$$\bar{a} \stackrel{\text{df}}{\bigvee}_U M \text{ implies } \bar{a} \stackrel{s}{\bigvee}_U M.$$

Consequently,

$$\bar{c}_m \equiv_U \bar{c}_n \text{ implies } \bar{c}_m \equiv_{U\bar{a}} \bar{c}_n, \quad \text{for } m, n < \omega.$$

Fixing a tuple \bar{a}' such that

$$\bar{a}(\bar{c}_n)_n \equiv_{U\bar{b}_0} \bar{a}'(\bar{b}_n)_n,$$

it follows that

$$\bar{c}_m \equiv_{U\bar{a}} \bar{c}_n \text{ implies } \bar{b}_m \equiv_{U\bar{a}'} \bar{b}_n, \quad \text{for } m, n < \omega. \quad \square$$

2. Forking in stable theories

In this section we collect properties of preforking relations in stable theories. First, note that we have seen in Corollary F3.2.19 that every stable theory is simple. Hence, in stable theories there is no difference between dividing and forking and both relations are symmetric.

Furthermore, in stable theories the relation $\stackrel{\text{df}}{\bigvee}$ is much better behaved than usual. For instance, we have already seen in Theorem C3.5.17 that $\stackrel{\text{df}}{\bigvee}$ is right reflexive.

Theorem 2.1. *Let T be a stable theory. Then*

$$A \stackrel{\text{df}}{\bigvee}_U U, \quad \text{for all } A, U \subseteq \mathbb{M}.$$

Let us consider the case where the base set is a model. In this case it will turn out that most preforking relations coincide.

Lemma 2.2. *Let T be a stable theory and \mathfrak{M} a model of T . Then*

$$A \downarrow_M^d B \text{ implies } A \bigvee_M B, \quad \text{for all } A, B \subseteq \mathbb{M}.$$

Proof. Suppose that $\bar{a} \downarrow_M^d \bar{b}$ and $\mathbb{M} \models \varphi(\bar{a}; \bar{b})$. We have to find a tuple $\bar{a}' \subseteq M$ such that $\mathbb{M} \models \varphi(\bar{a}'; \bar{b})$.

By Lemma F2.3.15, we have $\bar{a} \bigvee_M M$. Hence, we can use Proposition F2.4.10 to construct a \bigvee -Morley sequence $(\bar{a}^n)_{n < \omega}$ for $\text{tp}(\bar{a}/M)$ over M . Since $\bar{b} \downarrow_M^d \bar{a}$, it follows by Lemma F3.1.3 that there exists a tuple $\bar{b}' \equiv_{M\bar{a}} \bar{b}$ such that $(\bar{a}^n)_{n < \omega}$ is indiscernible over $M \cup \bar{b}'$. Consequently, $\mathbb{M} \models \varphi(\bar{a}^n; \bar{b}')$, for all $n < \omega$. By Theorem 2.1, we have

$$\bar{b}' \stackrel{\text{df}}{\bigvee}_{M \cup \bigcup_{n < \omega} \bar{a}^n} M \cup \bigcup_{n < \omega} \bar{a}^n.$$

Let $\delta(\bar{y})$ be a φ -definition of $\text{tp}(\bar{b}'/M \cup \bigcup_n \bar{a}^n)$ over $M \cup \bigcup_n \bar{a}^n$ and choose $n < \omega$ such that δ is a formula over $M \cup \bar{a}^0 \dots \bar{a}^{n-1}$. Since

$$\bar{a}^n \bigvee_M M \bar{a}^0 \dots \bar{a}^{n-1} \text{ and } \mathbb{M} \models \delta(\bar{a}^n),$$

there exists a tuple $\bar{a}' \subseteq M$ such that $\mathbb{M} \models \delta(\bar{a}')$. Hence, $\mathbb{M} \models \varphi(\bar{a}'; \bar{b})$, as desired. \square

Corollary 2.3. *Let T be a stable theory and \mathfrak{M} a model of T . Then*

$$A \downarrow_M^d B \text{ implies } A \stackrel{\text{df}}{\bigvee}_M B, \quad \text{for all } A, B \subseteq \mathbb{M}.$$

Proof. According to Theorem 2.1, we have $A \stackrel{\text{df}}{\bigvee}_M M$, which implies that $A \stackrel{*}{\bigvee}_M M$, by Lemma 1.2 (a). Hence, we can find a set $A' \equiv_M A$

with $A' \overset{\text{df}}{\vee}_M B$. Furthermore, $B \downarrow_M^d A$ implies $B \overset{\vee}{\vee}_M A$, by Lemma 2.2. Consequently, it follows by Lemma 1.1 (b) that $A' \equiv_{MB} A$. Therefore, $A' \overset{\text{df}}{\vee}_M B$ implies that $A \overset{\text{df}}{\vee}_M B$. \square

The previous results imply that, in a stable theory, the relations $\overset{\vee}{\vee}$, $\overset{i}{\vee}$, $\overset{\text{df}}{\vee}$, \downarrow^d , and \downarrow^f are all equivalent, at least over models.

Theorem 2.4. *Let T be a stable theory and \mathfrak{M} a model of T . Then*

$$\begin{aligned} A \overset{\vee}{\vee}_M B &\text{ iff } A \overset{i}{\vee}_M B &\text{ iff } A \overset{\text{df}}{\vee}_M B \\ &\text{ iff } A \downarrow_M^f B &\text{ iff } A \downarrow_M^d B. \end{aligned}$$

Proof. We have already seen in Proposition F3.1.12 that

$$A \overset{\vee}{\vee}_M B \Rightarrow A \overset{i}{\vee}_M B \Rightarrow A \downarrow_M^f B \Rightarrow A \downarrow_M^d B.$$

For stable theories, the implication $A \downarrow_M^d B \Rightarrow A \overset{\vee}{\vee}_M B$ is provided by Lemma 2.2. Furthermore, we can use Corollary 2.3 to show that

$$A \downarrow_M^d B \Rightarrow A \overset{\text{df}}{\vee}_M B,$$

while Lemma 1.1 (a) implies that

$$A \overset{\text{df}}{\vee}_M B \Rightarrow B \overset{\vee}{\vee}_M A.$$

Since we have already proved that, over models, $\overset{\vee}{\vee}$ is equivalent to \downarrow^d , it follows by symmetry of \downarrow^d that

$$A \overset{\text{df}}{\vee}_M B \Rightarrow B \overset{\vee}{\vee}_M A \Leftrightarrow A \overset{\vee}{\vee}_M B. \quad \square$$

There is an even closer connection between $\overset{i}{\vee}$, $\ast(\overset{\text{df}}{\vee})$, and \downarrow^f .

Proposition 2.5. *Let T be a stable theory.*

$$(a) \quad \overset{i}{\vee} = \ast(\overset{\text{df}}{\vee}).$$

$$(b) \quad A \downarrow_U^f B \quad \text{iff} \quad A \overset{i}{\vee}_{\text{acl}^{\text{eq}}(U)} B.$$

Proof. (a) (\supseteq) was already proved in Proposition 1.5. For (\subseteq), suppose that $\bar{a} \overset{i}{\vee}_U B$. To show that $\bar{a} \ast(\overset{\text{df}}{\vee})_U B$, we consider a set $C \supseteq B$. We have to find a tuple $\bar{a}' \equiv_{UB} \bar{a}$ with $\bar{a}' \overset{\text{df}}{\vee}_U C$. Let \mathfrak{M} be a $|U|^+$ -saturated and strongly $|U|^+$ -homogeneous model of T that contains $U \cup C$. Since $\bar{a} \overset{i}{\vee}_U B$, there exists a tuple $\bar{a}' \equiv_{UB} \bar{a}$ with $\bar{a}' \overset{s}{\vee}_U M$. Furthermore, we have seen in Theorem 2.1 that $\bar{a}' \overset{\text{df}}{\vee}_M M$. Consequently, it follows by Lemma 1.4 that $\bar{a}' \overset{\text{df}}{\vee}_U M$. In particular, $\bar{a}' \overset{\text{df}}{\vee}_U C$.

(b) (\Leftarrow) According to Proposition F3.1.12, $A \overset{i}{\vee}_{\text{acl}^{\text{eq}}(U)} B$ implies that $A \downarrow_{\text{acl}^{\text{eq}}(U)}^f B$. Moreover, we have $\text{acl}^{\text{eq}}(U) \downarrow_U^f A$, by Corollary F2.2.12 and Lemma F3.1.8. By symmetry and transitivity, it therefore follows that $A \downarrow_U^f B$.

(\Rightarrow) Suppose that $\bar{a} \downarrow_U^f B$. We will prove that $\bar{a} \ast(\overset{\text{df}}{\vee})_{\text{acl}^{\text{eq}}(U)} B$. Then the claim follows by (a). Hence, consider a set $C \supseteq \text{acl}^{\text{eq}}(U) \cup B$. We fix a $(|T| \oplus |U|)^+$ -saturated and strongly $(|T| \oplus |U|)^+$ -homogeneous model \mathfrak{N} containing C and a tuple $\bar{a}' \equiv_{UB} \bar{a}$ with $\bar{a}' \downarrow_U^f N$. We will prove that

$$\text{Gb}(\text{tp}(\bar{a}'/N)) \subseteq M^{\text{eq}}, \quad \text{for every model } \mathfrak{M} \text{ with } U \subseteq M \subseteq N.$$

By Lemma E2.1.9, this implies that $\text{Gb}(\text{tp}(\bar{a}'/N)) \subseteq \text{acl}^{\text{eq}}(U)$. Consequently, we can use Lemma E2.3.10 to show that $\text{tp}(\bar{a}'/N)$ is definable over $\text{acl}^{\text{eq}}(U)$. In particular, $\bar{a}' \overset{\text{df}}{\vee}_{\text{acl}^{\text{eq}}(U)} C$.

It remains to prove the above claim. Consider a model \mathfrak{M} with $U \subseteq M \subseteq N$. Then $\bar{a}' \downarrow_M^f N$ implies that $\bar{a}' \overset{\text{df}}{\vee}_M N$, by Theorem 2.4. Therefore, it follows by Lemma E2.3.8 that

$$\text{Gb}(\text{tp}(\bar{a}'/N)) \subseteq \text{dcl}^{\text{eq}}(M) = M^{\text{eq}}. \quad \square$$

Corollary 2.6. *In a stable theory,*

$$A \downarrow_U^f B \quad \text{implies} \quad A \overset{\text{df}}{\vee}_{\text{acl}^{\text{eq}}(U)} B.$$

3. Stationary types

In this section we study types with a unique free extension over every set of parameters. Such types are called *stationary*.

Definition 3.1. A type \mathfrak{p} over U is *stationary* if, for every set $C \subseteq \mathbb{M}$, \mathfrak{p} has a unique free extension to a complete type over $U \cup C$.

We start by proving that stationary types exist.

Proposition 3.2. *In a stable theory, every type over a set of the form $\text{acl}^{\text{eq}}(U)$ is stationary.*

Proof. Note that a type $\mathfrak{p}(\bar{x})$ is stationary if, and only if, for every finite tuple $\bar{x}' \subseteq \bar{x}$ of variables, the restriction $\mathfrak{p} \upharpoonright \bar{x}'$ is stationary. Hence, it is sufficient to consider types $\mathfrak{p} \in S^{<\omega}(\text{acl}^{\text{eq}}(U))$. Let $C \supseteq \text{acl}^{\text{eq}}(U)$ be a set and suppose that \bar{a} and \bar{a}' are two realisations of \mathfrak{p} with

$$\bar{a} \downarrow_{\text{acl}^{\text{eq}}(U)}^f C \quad \text{and} \quad \bar{a}' \downarrow_{\text{acl}^{\text{eq}}(U)}^f C.$$

We have to show that $\bar{a} \equiv_C \bar{a}'$. Hence, consider a formula $\varphi(\bar{x}; \bar{c})$ with $\bar{c} \subseteq C$. We choose an \aleph_0 -saturated model \mathfrak{M} containing $C \cup \bar{a}\bar{a}'$. There are tuples $\bar{a}_* \equiv_C \bar{a}$ and $\bar{a}'_* \equiv_C \bar{a}'$ with

$$\bar{a}_* \downarrow_{\text{acl}^{\text{eq}}(U)}^f M \quad \text{and} \quad \bar{a}'_* \downarrow_{\text{acl}^{\text{eq}}(U)}^f M.$$

Since $\bar{c} \downarrow_{\text{acl}^{\text{eq}}(U)}^f \bar{a}$, there is a tuple $\bar{c}_* \equiv_{\text{acl}^{\text{eq}}(U) \cup \bar{a}} \bar{c}$ with

$$\bar{c}_* \downarrow_{\text{acl}^{\text{eq}}(U)}^f M.$$

By Corollary 2.6, the types

$$\mathfrak{q} := \text{tp}(\bar{a}_*/M), \quad \mathfrak{q}' := \text{tp}(\bar{a}'_*/M), \quad \text{and} \quad \mathfrak{r} := \text{tp}(\bar{c}_*/M)$$

are definable over $\text{acl}^{\text{eq}}(U)$. Let $\delta(\bar{y})$, $\delta'(\bar{y})$, and $\varepsilon(\bar{x})$ be the corresponding φ -definitions. By Lemma 1.3, it follows that

$$\begin{aligned} \mathbb{M} \models \varphi(\bar{a}; \bar{c}) & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}_*; \bar{c}) \\ & \quad \text{iff} \quad \mathbb{M} \models \delta(\bar{c}) \\ & \quad \text{iff} \quad \mathbb{M} \models \delta(\bar{c}_*) \\ & \quad \text{iff} \quad \delta(\bar{y}) \in \mathfrak{r} \\ & \quad \text{iff} \quad \varepsilon(\bar{y}) \in \mathfrak{q} \\ & \quad \text{iff} \quad \varepsilon(\bar{y}) \in \mathfrak{q}' \\ & \quad \text{iff} \quad \delta'(\bar{y}) \in \mathfrak{r} \\ & \quad \text{iff} \quad \mathbb{M} \models \delta'(\bar{c}_*) \\ & \quad \text{iff} \quad \mathbb{M} \models \delta'(\bar{c}) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}'_*; \bar{c}) \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}'; \bar{c}). \quad \square \end{aligned}$$

Corollary 3.3. *In a stable theory types over models are stationary.*

Proof. Note that every type over a model M has a unique extension to a type over $M^{\text{eq}} = \text{dcl}^{\text{eq}}(M)$, which is an algebraically closed set. Hence, the claim follows by Proposition 3.2. \square

In Proposition 3.7 below we will present a characterisation of stationary types in terms of the relation \downarrow^f . We start with two technical lemmas. In the first one, we prove that all free extensions of a given type are conjugate.

Lemma 3.4. *Let T be a stable theory, $\kappa > |T|$ a cardinal, \mathfrak{M} a strongly κ -homogeneous model of T , and $U \subseteq M$ a set of size $|U| < \kappa$. If*

$$\bar{a} \equiv_U \bar{a}', \quad \bar{a} \downarrow_U^f M, \quad \text{and} \quad \bar{a}' \downarrow_U^f M,$$

then there exists an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ such that $\pi(\bar{a}') = \bar{a}$ and $\pi[M] = M$.

Proof. Since $\bar{a} \equiv_U \bar{a}'$, there exists an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ with $\pi(\bar{a}') = \bar{a}$. By Corollary E2.1.7, we have

$$\pi[\text{acl}^{\text{eq}}(U)] = \text{acl}^{\text{eq}}(U).$$

As \mathfrak{M} is strongly $|\text{acl}^{\text{eq}}(U)|^+$ -homogeneous, we can find an automorphism $\sigma_o \in \text{Aut } \mathfrak{M}$ with

$$\sigma_o \upharpoonright \text{acl}^{\text{eq}}(U) = \pi \upharpoonright \text{acl}^{\text{eq}}(U).$$

Let $\sigma \in \text{Aut } \mathbb{M}$ be an extension of σ_o . For every formula $\varphi(\bar{x}; \bar{c})$ with parameters $\bar{c} \subseteq \text{acl}^{\text{eq}}(U)$, it follows that

$$\begin{aligned} \mathbb{M} \models \varphi(\sigma(\bar{a}'); \bar{c}) & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}'; \sigma^{-1}(\bar{c})) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\pi(\bar{a}'); \pi(\sigma^{-1}(\bar{c}))) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}; \bar{c}). \end{aligned}$$

Hence, $\sigma(\bar{a}') \equiv_{\text{acl}^{\text{eq}}(U)} \bar{a}$. By invariance, we have

$$\sigma(\bar{a}') \downarrow_{\text{acl}^{\text{eq}}(U)}^f M \quad \text{and} \quad \bar{a} \downarrow_{\text{acl}^{\text{eq}}(U)}^f M.$$

Moreover, $\text{tp}(\bar{a}/\text{acl}^{\text{eq}}(U))$ is stationary according to Proposition 3.2. Therefore, $\sigma(\bar{a}') \equiv_M \bar{a}$ and there exists an automorphism $\rho \in \text{Aut } \mathbb{M}_M$ mapping $\sigma(\bar{a}')$ to \bar{a} . Since

$$\rho[\sigma[M]] = \rho[\sigma_o[M]] = \rho[M] = M,$$

the composition $\rho \circ \sigma \in \text{Aut } \mathbb{M}_U$ is the desired automorphism mapping \bar{a}' to \bar{a} . \square

The second lemma characterises those free extensions that are unique.

Definition 3.5. We write

$$\bar{a} \downarrow_U^! B \quad : \text{iff} \quad \text{tp}(\bar{a}/UB) \text{ is the unique free extension of } \text{tp}(\bar{a}/U) \text{ over } U \cup B.$$

Lemma 3.6. Let T be a stable theory, $\kappa > |T|$ a cardinal, \mathfrak{M} a strongly κ -homogeneous model of T , and $U \subseteq M$ a set of size $|U| < \kappa$. Then

$$\bar{a} \downarrow_U^! M \quad \text{iff} \quad \bar{a} \downarrow_U^f M \quad \text{and} \quad \bar{a} \not\downarrow_U^s M.$$

Proof. (\Leftarrow) As $\text{tp}(\bar{a}/M)$ is a free extension of $\text{tp}(\bar{a}/U)$, we only need to prove uniqueness. If $\text{tp}(\bar{a}/M)$ and $\text{tp}(\bar{a}'/M)$ are two free extensions of $\text{tp}(\bar{a}/U)$, we can use Lemma 3.4 to find an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ with $\pi(\bar{a}') = \bar{a}$ and $\pi[M] = M$. Hence, for every formula $\varphi(\bar{x}; \bar{y})$ over U and every $\bar{b} \subseteq M$, $\bar{a} \not\downarrow_U^s M$ implies that

$$\mathbb{M} \models \varphi(\bar{a}'; \bar{b}) \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}; \pi(\bar{b})) \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}; \bar{b}).$$

Consequently, $\bar{a} \equiv_M \bar{a}'$.

(\Rightarrow) If $\bar{a} \not\downarrow_U^f M$, the type $\text{tp}(\bar{a}/M)$ is not a free extension of $\text{tp}(\bar{a}/U)$ and we are done. Hence, suppose that $\bar{a} \downarrow_U^f M$ and $\bar{a} \not\downarrow_U^s M$. We claim that $\text{tp}(\bar{a}/U)$ has at least two free extensions over M . By assumption, we can find finite tuples $\bar{b}, \bar{b}' \subseteq M$ with $\bar{b} \equiv_U \bar{b}'$ and $\bar{b} \not\equiv_{U\bar{a}} \bar{b}'$. Let $\varphi(\bar{x}; \bar{y})$ be a formula over U such that

$$\mathbb{M} \models \varphi(\bar{a}; \bar{b}) \wedge \neg \varphi(\bar{a}; \bar{b}').$$

Since $\bar{b} \equiv_U \bar{b}'$ and \mathfrak{M} is strongly κ -homogeneous, there exists an automorphism $\pi_o \in \text{Aut } \mathfrak{M}_U$ mapping \bar{b} to \bar{b}' . Let π be an automorphism of \mathbb{M} extending π_o . Then

$$\mathbb{M} \models \varphi(\bar{a}; \bar{b}) \quad \text{implies} \quad \mathbb{M} \models \varphi(\pi(\bar{a}); \bar{b}').$$

Hence, $\pi(\bar{a}) \not\equiv_M \bar{a}$. Furthermore,

$$\bar{a} \downarrow_U^f M \quad \text{implies} \quad \pi(\bar{a}) \downarrow_U^f \pi[M].$$

Since $\pi[M] = \pi_o[M] = M$, it follows that $\text{tp}(\bar{a}/M)$ and $\text{tp}(\pi(\bar{a})/M)$ are two different free extensions of $\text{tp}(\bar{a}/U)$. \square

Proposition 3.7. Let T be stable. Then

$$\text{tp}(\bar{a}/U) \text{ is stationary} \quad \text{iff} \quad \bar{a} \downarrow_U^! U.$$

Proof. (\Rightarrow) Suppose that $\mathfrak{p} := \text{tp}(\bar{a}/U)$ is stationary. Since $\checkmark = {}^*(\checkmark)$, it is sufficient to show that $\bar{a} \checkmark ({}^*(\checkmark))_U U$. Hence, consider a set $B \supseteq U$. We fix a strongly $(|T|^+ \oplus |U|^+)$ -homogeneous model \mathfrak{M} containing B . Let \mathfrak{q} be the unique free extension of \mathfrak{p} to M and let \bar{a}' be a realisation of \mathfrak{q} . Then $\bar{a}' \equiv_U \bar{a}$ and it follows by Lemma 3.6 that $\bar{a}' \checkmark_U M$. In particular, we have $\bar{a}' \checkmark_U B$.

(\Leftarrow) Suppose that $\bar{a} \checkmark_U U$. To show that $\mathfrak{p} := \text{tp}(\bar{a}/U)$ is stationary, consider a set $B \supseteq U$. We fix a strongly $(|T|^+ \oplus |U|^+)$ -homogeneous model \mathfrak{M} containing B . Let $\bar{a}' \equiv_U \bar{a}$ be a tuple with $\bar{a}' \checkmark_U M$. By Proposition F3.1.12, it follows that $\bar{a}' \downarrow_U^f M$ and $\bar{a}' \checkmark_U M$. Therefore, Lemma 3.6 implies that $\bar{a}' \downarrow_U^! M$. In particular, $\text{tp}(\bar{a}'/B)$ is the unique free extension of \mathfrak{p} over B . \square

As an application of stationary types, we present the following topological characterisation of the set of free extensions of a type.

Theorem 3.8 (Open Mapping Theorem). *Let T be a stable theory, $U \subseteq A$ sets, and let $\mathfrak{F}^{\bar{s}}(A/U)$ denote the subspace of $\mathfrak{S}^{\bar{s}}(A)$ consisting of all types that do not fork over U .*

- (a) $F^{\bar{s}}(A/U)$ is a closed subset of $\mathfrak{S}^{\bar{s}}(A)$.
- (b) The restriction map $\rho : \mathfrak{F}^{\bar{s}}(A/U) \rightarrow \mathfrak{S}^{\bar{s}}(U) : \mathfrak{p} \mapsto \mathfrak{p}|_U$ is continuous, closed, open, and surjective.

Proof. (a) We use (DEF) to fix, for every type $\mathfrak{p} \in S^{\bar{s}}(A) \setminus F^{\bar{s}}(A/U)$, a formula $\varphi_{\mathfrak{p}}(\bar{x}) \in \mathfrak{p}$ such that

$$\mathbb{M} \models \varphi_{\mathfrak{p}}(\bar{c}) \quad \text{implies} \quad \bar{c} \not\downarrow_U^f A.$$

Setting

$$\Phi := \{ \neg\varphi_{\mathfrak{p}} \mid \mathfrak{p} \in S^{\bar{s}}(A) \setminus F^{\bar{s}}(A/U) \}$$

it follows that $\Phi \subseteq \mathfrak{p}$, for every $\mathfrak{p} \in F^{\bar{s}}(A/U)$, while $\neg\varphi_{\mathfrak{p}} \in \Phi \setminus \mathfrak{p}$, for every $\mathfrak{p} \notin F^{\bar{s}}(A/U)$. Hence,

$$F^{\bar{s}}(A/U) = \langle \Phi \rangle_{\mathfrak{S}^{\bar{s}}(A)},$$

which is a closed set.

(b) We have seen in Corollary c3.2.22 that the restriction map

$$\rho^+ : \mathfrak{S}^{\bar{s}}(A) \rightarrow \mathfrak{S}^{\bar{s}}(U) : \mathfrak{p} \mapsto \mathfrak{p}|_U$$

is continuous, closed, and surjective. By (a) it follows that the restriction

$$\rho = \rho^+ \upharpoonright F^{\bar{s}}(A/U) : \mathfrak{F}^{\bar{s}}(A/U) \rightarrow \mathfrak{S}^{\bar{s}}(U)$$

is also continuous and closed. Furthermore, it follows by (EXT) that every type over U has a free extension to a type over A . Hence, ρ is surjective and it remains to prove that it is open.

First, we consider the case where A is a strongly $|U|^+$ -homogeneous model. Every open set $O \subseteq F^{\bar{s}}(A/U)$ is a union of basic open sets of the form

$$\langle \varphi \rangle_{\mathfrak{F}^{\bar{s}}(A/U)} := \{ \mathfrak{p} \in F^{\bar{s}}(A/U) \mid \varphi \in \mathfrak{p} \}.$$

Therefore it is sufficient to prove that the image of a basic open set is open. Let $\varphi(\bar{x}; \bar{y})$ be a formula over U , $\bar{c} \subseteq A$ parameters, and let

$$\overline{\langle \varphi(\bar{x}; \bar{c}) \rangle}_{\mathfrak{F}^{\bar{s}}(A/U)} := \bigcup \{ \langle \varphi(\bar{x}; \pi(\bar{c})) \rangle_{\mathfrak{F}^{\bar{s}}(A/U)} \mid \pi \in \text{Aut } \mathfrak{M}_U \}$$

be the closure of $\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^{\bar{s}}(A/U)}$ under conjugates. Being a union of open sets, this set is also open. Furthermore,

$$\rho^{-1}[\rho[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^{\bar{s}}(A/U)}]] = \overline{\langle \varphi(\bar{x}; \bar{c}) \rangle}_{\mathfrak{F}^{\bar{s}}(A/U)},$$

since, for types $\mathfrak{p}, \mathfrak{q} \in F^{\bar{s}}(A/U)$ with $\mathfrak{p}|_U = \mathfrak{q}|_U$, we can use Lemma 3.4 to find an automorphism $\pi \in \text{Aut } \mathfrak{M}_U$ with $\pi(\mathfrak{p}) = \mathfrak{q}$.

For a type $\mathfrak{p}_0 \in S^{\bar{s}}(U)$, it follows that

$$\begin{aligned} & \mathfrak{p}_0 \notin \rho[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^{\bar{s}}(A/U)}] \\ \text{iff} & \quad \rho^{-1}(\mathfrak{p}_0) \cap \langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^{\bar{s}}(A/U)} = \emptyset \\ \text{iff} & \quad \rho^{-1}(\mathfrak{p}_0) \cap \overline{\langle \varphi(\bar{x}; \bar{c}) \rangle}_{\mathfrak{F}^{\bar{s}}(A/U)} = \emptyset \end{aligned}$$

$$\begin{aligned} \text{iff } & \rho^{-1}(\mathfrak{p}_o) \setminus \overline{\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}} \neq \emptyset \\ \text{iff } & \mathfrak{p}_o \in \rho[F^s(A/U) \setminus \overline{\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}}]. \end{aligned}$$

Hence, the complement of $\rho[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}]$ is the image of a closed set. As we have shown above that the map ρ is closed, it follows that the complement is closed and the image $\rho[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}]$ is open.

It remains to prove the general case. We fix a strongly $|U|^+$ -homogeneous model \mathfrak{M} containing A and let

$$\rho' : \mathfrak{F}^s(M/U) \rightarrow \mathfrak{F}^s(A/U)$$

be the corresponding restriction map. Consider a basic open set

$$\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}$$

in $\mathfrak{F}^s(A/U)$. Then $\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(M/U)}$ is basic open in $\mathfrak{F}^s(M/U)$ and

$$\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(M/U)} = (\rho')^{-1}[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}].$$

As ρ' is surjective, we have

$$\begin{aligned} \rho[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}] &= \rho[\rho'[(\rho')^{-1}[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(A/U)}]]] \\ &= (\rho \circ \rho')[\langle \varphi(\bar{x}; \bar{c}) \rangle_{\mathfrak{F}^s(M/U)}] \end{aligned}$$

This set is open, since we have shown above that the composition $\rho \circ \rho'$ is an open map. \square

4. The multiplicity of a type

Most types have several free extensions. In this section we study their number. We will prove in Theorem 4.6 below that a theory is stable if, and only if, the number of such extensions is bounded.

Definition 4.1. Let T be a theory and \surd a forking relation.

(a) The \surd -multiplicity $\text{mult}_{\surd}(\mathfrak{p})$ of a type \mathfrak{p} over A is the minimal cardinal κ such that, for every set $B \supseteq A$, there are at most κ complete types over B that are \surd -free extensions of \mathfrak{p} . If such a cardinal κ does not exist, we set $\text{mult}_{\surd}(\mathfrak{p}) := \infty$. For $\surd = \downarrow^f$, we drop the subscript and simply write $\text{mult}(\mathfrak{p})$.

(b) The multiplicity $\text{mult}(\surd)$ of \surd is the maximal \surd -multiplicity of some complete type (with finitely many variables). If there is no maximum, we set $\text{mult}(\surd) := \infty$.

We start by proving that, in a stable theory, every type has bounded multiplicity.

Lemma 4.2. Let T be a stable theory.

(a) For every type $\mathfrak{p} \in S^s(A)$ with $|\bar{s}| < \omega$, there exists some model \mathfrak{M} of T of size $|M| \leq |T|$ such that

$$\text{mult}_{\downarrow^f}(\mathfrak{p}) \leq |S^s(M)|.$$

(b) $\text{mult}(\downarrow^f) \leq \sup \{ |S^{<\omega}(U)| \mid |U| \leq |T| \} \leq 2^{|T|}$

Proof. (a) For every type $\mathfrak{p} \in S^s(A)$, there exists a set $U \subseteq A$ of size

$$|U| < \text{loc}_o(\downarrow^f) \leq \text{fc}(\downarrow^f) \leq |T|^+$$

such that \mathfrak{p} does not fork over U . Since $|U| \leq |T|$, we can find a model \mathfrak{M} of size $|M| = |T|$ containing U . Since every type has at least one free extension over any given set, it is sufficient to bound the number of free extensions of \mathfrak{p} over sets B containing M . Hence, let $\mathfrak{q} \in S^s(B)$ be an extension of \mathfrak{p} with $B \supseteq M$. We have seen in Corollary 3.3 that types over models are stationary. Hence, \mathfrak{q} is the unique free extension of $\mathfrak{q}|_M$. Consequently, if $\mathfrak{q}, \mathfrak{q}' \in S^s(B)$ are distinct free extensions of \mathfrak{p} , then $\mathfrak{q}|_M \neq \mathfrak{q}'|_M$. Therefore, \mathfrak{p} has at most $|S^s(M)|$ free extensions.

(b) The first inequality follows immediately from (a). The second one follows from the fact that there are at most $2^{|T|}$ types over a set of size $|T|$. \square

A characterisation of stable theories

Recall that we write $\mathfrak{p} \leq_{\sqrt{}} \mathfrak{q}$ if \mathfrak{q} is a $\sqrt{\quad}$ -free extension of \mathfrak{p} . This is a definition of $\leq_{\sqrt{}}$ in terms of a given preforking relation $\sqrt{\quad}$. Conversely, given an extension relation \leq we can recover a corresponding preforking relation $\sqrt{\quad}$. In the remainder of this section we present two characterisations of stable theories one in terms of the extension relation \leq and one in terms of the parameter $\text{mult}(\downarrow^f)$.

Proposition 4.3. *If \downarrow is a symmetric forking relation with $\text{mult}(\downarrow) < \infty$, then \leq_{\downarrow} satisfies the following conditions:*

- (INV) *Invariance.* $\mathfrak{p} \leq \mathfrak{q}$ implies $\pi(\mathfrak{p}) \leq \pi(\mathfrak{q})$, for every automorphism $\pi \in \text{Aut}(\mathbb{M})$.
- (LC) *Local Character.* There exists a cardinal κ such that, for every set U and every type $\mathfrak{p} \in S^{<\omega}(U)$, there exists a subset $U_o \subseteq U$ of size $|U_o| < \kappa$ such that $\mathfrak{p} \upharpoonright U_o \leq \mathfrak{p}$.
- (BND) *Boundedness.* For every type $\mathfrak{p} \in S^{<\omega}(U)$, there exists a cardinal μ such that, for every set $C \subseteq \mathbb{M}$, \mathfrak{p} has at most μ extensions $\mathfrak{q} \in S^{<\omega}(U \cup C)$ with $\mathfrak{p} \leq \mathfrak{q}$.
- (EXT) *Extension.* For every $\mathfrak{p} \in S^{<\omega}(U)$ and every set $C \subseteq \mathbb{M}$, there exists some type $\mathfrak{q} \in S^{<\omega}(U \cup C)$ with $\mathfrak{p} \leq \mathfrak{q}$.
- (TR) *Transitivity.* $\mathfrak{p} \leq \mathfrak{q} \leq \mathfrak{r}$ implies $\mathfrak{p} \leq \mathfrak{r}$.
- (MON) *Monotonicity.* $\mathfrak{p} \leq \mathfrak{r}$ implies $\mathfrak{p} \leq \mathfrak{q}$, for all $\mathfrak{p} \subseteq \mathfrak{q} \subseteq \mathfrak{r}$.

Proof. (BND) holds since $\text{mult}(\downarrow) < \infty$. The other axioms follow from the fact that \downarrow is a symmetric forking relation: (INV) follows by invariance; (LC) follows by right locality; (EXT) follows by the extension axiom; (TR) follows by left transitivity and symmetry; and (MON) follows by monotonicity. \square

For the converse statement, we need a technical lemma.

Lemma 4.4. *Let \mathfrak{M} be a κ^+ -saturated, strongly κ^+ -homogeneous model, $U \subseteq M$ a set of size $|U| < \kappa$, and $\bar{a} \in \mathbb{M}^{<\omega}$. If $\bar{a} \not\equiv_U^{\downarrow} M$, then $\text{tp}(\bar{a}/M)$ has at least κ conjugates over U .*

Proof. If $\bar{a} \not\equiv_U^{\downarrow} M$, there exists a finite tuple $\bar{b} \subseteq M$ such that $\bar{a} \not\equiv_U^{\downarrow} \bar{b}$. By Lemma F3.1.3, we can find an indiscernible sequence $(\bar{b}_n)_{n < \omega}$ over U with $\bar{b} = \bar{b}_0$ such that, for each tuple $\bar{a}' \equiv_{U\bar{b}} \bar{a}$, there are indices $m, n < \omega$ with

$$\bar{b}_m \not\equiv_{U\bar{a}'} \bar{b}_n.$$

Fix a formula $\varphi(\bar{x}; \bar{y})$ over U such that

$$\mathbb{M} \models \varphi(\bar{a}; \bar{b}_m) \wedge \neg \varphi(\bar{a}; \bar{b}_n), \quad \text{for some } m, n < \omega.$$

Replacing φ be its negation, if necessary, we may assume that there are infinitely many indices n such that $\mathbb{M} \models \neg \varphi(\bar{a}; \bar{b}_n)$. By compactness, it follows that there exists a tuple $\bar{a}' \equiv_U \bar{a}$ and an indiscernible sequence $(\bar{b}'_\alpha)_{\alpha < \kappa}$ of length κ such that

$$\mathbb{M} \models \varphi(\bar{a}'; \bar{b}'_\alpha) \quad \text{iff} \quad \alpha = 0.$$

As \mathfrak{M} is κ^+ -saturated, we may choose the sequence $(\bar{b}'_\alpha)_{\alpha < \kappa}$ to be in M . By strong κ^+ -homogeneity we can find, for every $\alpha < \kappa$, an automorphism $\sigma_\alpha \in \text{Aut } \mathfrak{M}_U$ such that

$$\sigma_\alpha(\bar{b}'_\beta) = \bar{b}'_{\alpha+\beta}, \quad \text{for all } \beta < \kappa.$$

Let $\pi_\alpha \in \text{Aut } \mathbb{M}$ be an extension of σ_α and set

$$\bar{a}_\alpha := \pi_\alpha(\bar{a}'), \quad \text{for } \alpha < \kappa.$$

Then

$$\begin{aligned} \mathbb{M} \models \varphi(\bar{a}_\alpha; \bar{b}'_{\alpha+\beta}) & \quad \text{iff} \quad \mathbb{M} \models \varphi(\pi_\alpha(\bar{a}'); \pi_\alpha(\bar{b}'_\beta)) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}'; \bar{b}'_\beta) \\ & \quad \text{iff} \quad \beta = 0. \end{aligned}$$

Consequently, $\text{tp}(\bar{a}_\alpha/M) \neq \text{tp}(\bar{a}_\beta/M)$, for $\alpha \neq \beta$. Since these types are extensions of $\text{tp}(\bar{a}/U)$ and they are conjugate over U , the claim follows. \square

Theorem 4.5. *Let T be a complete first-order theory.*

- (a) *T is stable if, and only if, there exists a relation \leq on complete types satisfying (INV), (LC), and (BND).*
- (b) *If \leq is an extension relation satisfying (INV), (LC), (BND), (EXT), (TR), and (MON), then $\leq = \leq_{\text{f}}$.*

Proof. (a) (\Rightarrow) If T is stable, the relation \leq_{f} has the desired properties by Proposition 4.3. For (\Leftarrow) , suppose that \leq satisfies (INV), (LC), and (BND). Let κ be the cardinal from (LC) and fix a κ -saturated model \mathfrak{M} . For $U \subseteq M$ and $\mathfrak{p} \in S^{<\omega}(U)$, we denote by $\mu(\mathfrak{p}; U)$ the cardinal from (BND). Set

$$\mu := \sup \{ \mu(\mathfrak{p}; U) \mid U \subseteq M \text{ with } |U| < \kappa \text{ and } \mathfrak{p} \in S^{<\omega}(U) \}.$$

Since, for every subset $U \subseteq \mathbb{M}$ of size $|U| < \kappa$, there is some automorphism $\pi \in \text{Aut } \mathbb{M}$ with $\pi[U] \subseteq M$, it follows by (INV) that, for all sets $U, C \subseteq \mathbb{M}$ with $|U| < \kappa$ and for every type $\mathfrak{p} \in S^{<\omega}(U)$, there are at most μ types $\mathfrak{q} \in S^{<\omega}(U \cup C)$ with $\mathfrak{p} \leq \mathfrak{q}$.

Fix a set $U \subseteq \mathbb{M}$ and a finite tuple \bar{s} of sorts. For every type $\mathfrak{p} \in S^{\bar{s}}(U)$, we can fix, by (LC), a subset $C(\mathfrak{p}) \subseteq U$ of size $|C(\mathfrak{p})| < \kappa$ such that $\mathfrak{p}|_{C(\mathfrak{p})} \leq \mathfrak{p}$. Let $C \subseteq U$ be a set of size $|C| < \kappa$. Then

- ♦ $|S^{\bar{s}}(C)| \leq 2^{|T| \oplus \kappa}$ and,
- ♦ for every type $\mathfrak{q} \in S^{\bar{s}}(C)$, there are at most μ types $\mathfrak{p} \in S^{\bar{s}}(U)$ with $C(\mathfrak{p}) = C$ and $\mathfrak{p}|_C = \mathfrak{q}$.

Consequently, we have

$$|S^{\bar{s}}(U)| \leq |U|^{<\kappa} \otimes 2^{|T| \oplus \kappa} \otimes \mu.$$

Setting $\lambda_o := \mu \oplus 2^{|T|}$ and $\lambda := \lambda_o^\kappa$, it follows that

$$|S^{\bar{s}}(U)| \leq \lambda^\kappa \otimes \lambda_o^\kappa \otimes \mu = \lambda, \quad \text{for every set } U \text{ of size } |U| \leq \lambda.$$

Hence, T is λ -stable.

(b) Let $\mathfrak{q} \in S^{\bar{s}}(B)$ be an extension of $\mathfrak{p} \in S^{\bar{s}}(A)$. We have to show that $\mathfrak{p} \leq \mathfrak{q}$ if, and only if, \mathfrak{q} is a \downarrow^{f} -free extension of \mathfrak{p} .

(\Rightarrow) Suppose that $\mathfrak{p} \leq \mathfrak{q}$. By (BND), there is a cardinal μ such that, for every set C , \mathfrak{p} has at most μ \leq -free extensions over $A \cup C$. Set $\kappa := \mu^+ \oplus |A|^+$ and let \mathfrak{M} be a κ^+ -saturated and strongly κ^+ -homogeneous model containing B . We use (EXT) to find a type $\mathfrak{r} \geq \mathfrak{q}$ over M . By (TR), it follows that $\mathfrak{p} \leq \mathfrak{r}$. Hence, (INV) implies that \mathfrak{r} has at most μ conjugates over A . By Lemma 4.4, it follows that \mathfrak{r} does not fork over A . In particular, \mathfrak{q} does not fork over A .

(\Leftarrow) Suppose that \mathfrak{q} is a free extension of \mathfrak{p} . Fix a strongly $(|T| \oplus |A|)^+$ -homogeneous model \mathfrak{M} containing B and let \mathfrak{r} be a free extension of \mathfrak{q} over M . By (EXT), there exists a type $\mathfrak{r}' \geq \mathfrak{p}$ over M . By the first part of the proof, $\mathfrak{p} \leq \mathfrak{r}'$ implies that \mathfrak{r}' is a free extension of \mathfrak{p} . Let \bar{a} and \bar{a}' be realisations of, respectively, \mathfrak{r} and \mathfrak{r}' . Then we can use Lemma 3.4 to find an automorphism $\pi \in \text{Aut } \mathbb{M}_A$ such that $\pi(\bar{a}') = \bar{a}$ and $\pi[M] = M$. By (INV),

$$\mathfrak{p} \leq \mathfrak{q} \quad \text{implies} \quad \mathfrak{p} \leq \text{tp}(\pi(\bar{a}')/\pi[M]) = \text{tp}(\bar{a}/M).$$

Hence, it follows by (MON) that $\mathfrak{p} \leq \text{tp}(\bar{a}/B) = \mathfrak{q}$. \square

Translating this theorem into the language of forking relations, we obtain the following characterisation of stable theories.

Theorem 4.6. *Let T be a complete first-order theory. The following statements are equivalent.*

- (1) *T is stable.*
- (2) *$\text{f}/$ is symmetric and $\text{mult}(\text{f}/) < \infty$.*
- (3) *There exists a symmetric forking relation \downarrow with $\text{mult}(\downarrow) < \infty$.*

Proof. (1) \Rightarrow (2) follows by Lemma 4.2 and the implication (2) \Rightarrow (3) is trivial. For (3) \Rightarrow (1), suppose that \downarrow is a symmetric forking relation with $\text{mult}(\downarrow) < \infty$. By Proposition 4.3, the corresponding extension

relation \leq_{\downarrow} satisfies the three axioms from Theorem 4.5 (a). Hence, T is stable. \square

Proposition 4.7. *If \downarrow is a symmetric forking relation with $\text{mult}(\downarrow) < \infty$, then $\downarrow = \downarrow^f$.*

Proof. By Proposition 4.3, the extension relation \leq_{\downarrow} satisfies the six axioms from Theorem 4.5 (b). Consequently, $\leq_{\downarrow} = \leq_{\downarrow^f}$. For a finite tuple \bar{a} and sets $U, B \subseteq \mathbb{M}$, it follows that

$$\begin{aligned} \bar{a} \downarrow_U B & \text{ iff } \text{tp}(\bar{a}/U) \leq_{\downarrow} \text{tp}(\bar{a}/B) \\ & \text{ iff } \text{tp}(\bar{a}/U) \leq_{\downarrow^f} \text{tp}(\bar{a}/B) \quad \text{iff} \quad \bar{a} \downarrow_U^f B. \end{aligned}$$

Hence, finite character implies that $\downarrow = \downarrow^f$. \square

As a further application we derive a characterisation of forking in totally transcendental theories by showing that the relation of being a Morley-free extension (which was defined in Section F2.1) satisfies the conditions of the above theorem.

Corollary 4.8. *Let T be a totally transcendental theory.*

- (a) $\bar{a} \downarrow_U^f B$ iff $\text{rk}_M(\bar{a}/UB) = \text{rk}_M(\bar{a}/U)$, for all finite \bar{a} .
- (b) $\text{mult}_{\downarrow^f}(\mathfrak{p}) < \aleph_0$.

Proof. (a) For types $\mathfrak{p} \in S^s(U)$ and $\mathfrak{q} \in S^s(V)$, we define

$$\mathfrak{p} \leq_M \mathfrak{q} \quad \text{: iff} \quad \mathfrak{q} \text{ is a Morley-free extension of } \mathfrak{p}.$$

It is sufficient to show that \leq_M satisfies the conditions in Theorem 4.5 (b).

(INV) follows immediately from the definition. (BND) and (EXT) were already shown in Lemma F2.1.9 (a) and (b), respectively, while (LC) was proved in Lemma F2.1.6 (c).

For (TR), suppose that $\mathfrak{p} \leq_M \mathfrak{q} \leq_M \mathfrak{r}$. Then

$$\text{rk}_M(\mathfrak{p}) = \text{rk}_M(\mathfrak{q}) = \text{rk}_M(\mathfrak{r}),$$

which implies that $\mathfrak{p} \leq_M \mathfrak{r}$.

(MON) Suppose that $\mathfrak{p} \leq_M \mathfrak{r}$ and $\mathfrak{p} \subseteq \mathfrak{q} \subseteq \mathfrak{r}$. By Lemma F2.1.6 (b), we have

$$\text{rk}_M(\mathfrak{p}) \geq \text{rk}_M(\mathfrak{q}) \geq \text{rk}_M(\mathfrak{r}) = \text{rk}_M(\mathfrak{p}).$$

Hence, $\mathfrak{p} \leq_M \mathfrak{q}$.

(b) We have seen in Lemma F2.1.9 that every type has only finitely many Morley-free extensions. Hence, the claim follows by (a). \square

5. Morley sequences in stable theories

Let us collect several results on Morley sequences in stable theories. Many of the proofs rely on the notion of a stationary type. We start with a proof that we can drop the requirement of indiscernibility from the definition of a Morley sequence if the type in question is stationary.

Lemma 5.1. *Let T be a stable theory, \mathfrak{p} a stationary type over U , and let $(\bar{a}_i)_{i \in I}$ be a sequence of realisations of \mathfrak{p} . If*

$$\bar{a}_i \downarrow_U^f \bar{a}[\langle i \rangle], \quad \text{for all } i \in I,$$

then $(\bar{a}_i)_{i \in I}$ is a \downarrow^f -Morley sequence for \mathfrak{p} over U .

Proof. We have to show that $(\bar{a}_i)_{i \in I}$ is indiscernible over U . By induction on $n < \omega$, we prove that

$$\bar{a}[\bar{i}] \equiv_{U\bar{a}[\langle l \rangle \bar{a}[\langle m \rangle]]} \bar{a}[\bar{k}], \quad \text{for all } \bar{i}, \bar{k} \in [I]^n \text{ with } l < \bar{i}, \bar{k} < m.$$

Hence, let $\bar{i}, \bar{k} \in [I]^n$ and $l < \bar{i}, \bar{k} < m$. By symmetry, we may assume that $i_{n-1} < k_{n-1}$. According to Lemma F2.4.9, we have

$$\bar{a}_{i_{n-1}} \downarrow_U^f \bar{a}[\langle i_{n-1} \rangle \bar{a}[\langle m \rangle]] \quad \text{and} \quad \bar{a}_{k_{n-1}} \downarrow_U^f \bar{a}[\langle i_{n-1} \rangle \bar{a}[\langle m \rangle]],$$

Since \mathfrak{p} is stationary, it follows that

$$\bar{a}_{i_{n-1}} \equiv_{U\bar{a}[\langle i_{n-1} \rangle \bar{a}[\langle m \rangle]]} \bar{a}_{k_{n-1}}.$$

By inductive hypothesis, we further have

$$\bar{a}_{i_{n-2}} \dots \bar{a}_{i_0} \equiv_{U\bar{a}[\langle l \rangle] \bar{a}[\geq k_{n-1}]} \bar{a}_{k_{n-2}} \dots \bar{a}_{k_0}.$$

Consequently,

$$\begin{aligned} \bar{a}_{i_{n-1}} \bar{a}_{i_{n-2}} \dots \bar{a}_{i_0} &\equiv_{U\bar{a}[\langle l \rangle] \bar{a}[\geq m]} \bar{a}_{k_{n-1}} \bar{a}_{i_{n-2}} \dots \bar{a}_{i_0} \\ &\equiv_{U\bar{a}[\langle l \rangle] \bar{a}[\geq m]} \bar{a}_{k_{n-1}} \bar{a}_{k_{n-2}} \dots \bar{a}_{k_0}. \end{aligned} \quad \square$$

Lemma 5.2. *Let T be a stable theory, \mathfrak{p} a stationary type over U and \mathfrak{q} the unique free extensions of \mathfrak{p} over $U \cup C$.*

- (a) *Every \downarrow^f -Morley sequences $(\bar{a}_i)_{i \in I}$ for \mathfrak{q} over $U \cup C$ is also a \downarrow^f -Morley sequence for \mathfrak{p} over U .*
- (b) *Let $(\bar{a}_i)_{i \in I}$ be a \downarrow^f -Morley sequence for \mathfrak{p} over U . If $I_0 \subseteq I$ and $C \downarrow_{U\bar{a}[I_0]}^f \bar{a}[I]$, then $(\bar{a}_i)_{i \in I \setminus I_0}$ is a Morley sequence for \mathfrak{q} over $U \cup C$.*

Proof. (a) As $(\bar{a}_i)_{i \in I}$ is indiscernible over $U \cup C$, it is trivially indiscernible over U . Furthermore,

$$\bar{a}_i \downarrow_U^f C \text{ and } \bar{a}_i \downarrow_{UC}^f \bar{a}[\langle i \rangle] \text{ implies } \bar{a}_i \downarrow_U^f \bar{a}[\langle i \rangle].$$

- (b) Set $A_0 := \bar{a}[I_0]$. We start by showing that

$$C\bar{a}[K] \downarrow_{UA_0}^f \bar{a}[I \setminus K], \text{ for all finite } K \subseteq I \setminus I_0.$$

The proof is by induction on $|K|$. If $K = \emptyset$, the claim holds by assumption. Hence, suppose that $K = K_0 \cup \{k\}$ and that we have already shown that

$$C\bar{a}[K_0] \downarrow_{UA_0}^f \bar{a}[I \setminus K_0].$$

Then

$$C\bar{a}[K_0] \downarrow_{UA_0 \bar{a}_k}^f \bar{a}[I \setminus K].$$

Furthermore, we have seen in Lemma F2.4.9 that

$$\bar{a}_k \downarrow_U^f \bar{a}[I \setminus K],$$

which implies that

$$\bar{a}_k \downarrow_{UA_0}^f \bar{a}[I \setminus K].$$

Consequently, it follows by transitivity that

$$C\bar{a}[K_0] \bar{a}_k \downarrow_{UA_0}^f \bar{a}[I \setminus K].$$

Having proved the claim, it follows by (FIN) that

$$C\bar{a}[\langle i \rangle] \downarrow_{UA_0}^f \bar{a}_i, \text{ for all } i \in I,$$

which implies that

$$\bar{a}_i \downarrow_{UCA_0}^f \bar{a}[\langle i \rangle], \text{ for all } i \in I \setminus I_0.$$

Hence, it follows by Lemma 5.1 that $(\bar{a}_i)_{i \in I \setminus I_0}$ is a \downarrow^f -Morley sequences over $U \cup C \cup A_0$. \square

For stable theories, we can turn every indiscernible sequence into a Morley sequence by increasing the domain of the type.

Proposition 5.3. *Let T be a stable theory, κ an infinite cardinal, and let $(\bar{a}_i)_{i \in I}$ be an infinite indiscernible sequence over U such that $|\bar{a}_i| < \kappa$, for all $i \in I$. There exist a set C of size $|C| < \kappa \oplus \aleph_1$ and a stationary type $\mathfrak{p} \in S^{<\kappa}(U \cup C)$ such that $(\bar{a}_i)_{i \in I}$ is a \downarrow^f -Morley sequence for \mathfrak{p} over $U \cup C$.*

Proof. We have seen in Corollary E5.4.13 and Corollary E5.4.14 that, for every formula $\varphi(\bar{x}; \bar{c})$ the set

$$\llbracket \varphi(\bar{a}_i; \bar{c}) \rrbracket_{i \in I} = \{ i \in I \mid \mathbb{M} \models \varphi(\bar{a}_i; \bar{c}) \}$$

is either finite or cofinite and that, for every set $C \subseteq \mathbb{M}$, the type

$$\text{Av}_1((\bar{a}_i)_i/UC) = \{ \varphi(\bar{x}) \mid \varphi \text{ a formula over } U \cup C \text{ such that} \\ \llbracket \varphi(\bar{a}_i) \rrbracket_{i \in I} \text{ is cofinite} \}$$

is complete. According to Proposition E5.4.12, there exists, for every formula $\varphi(\bar{x}; \bar{y})$ over U , a finite constant $k(\varphi) < \omega$ such that, for all $\bar{c} \subseteq \mathbb{M}$,

$$\varphi(\bar{x}; \bar{c}) \in \text{Av}_1((\bar{a}_i)_i/UC) \quad \text{iff} \quad |I \setminus \llbracket \varphi(\bar{a}_i; \bar{c}) \rrbracket_{i \in I}| \leq k(\varphi).$$

Choose an injective function $\mu : \omega \rightarrow I$ and set $I_o := \text{rng } \mu$. It follows that

$$\begin{aligned} & \varphi(\bar{x}; \bar{c}) \in \text{Av}_1((\bar{a}_i)_i/UC) \\ \text{iff} & \quad \left| \llbracket \varphi(\bar{a}_{\mu(n)}; \bar{c}) \rrbracket_{n < 2k(\varphi)+1} \right| > k(\varphi) \\ \text{iff} & \quad \mathbb{M} \models \bigvee \left\{ \bigwedge_{n \in K} \varphi(\bar{a}_{\mu(n)}; \bar{c}) \mid K \subseteq [2k(\varphi) + 1], |K| = k(\varphi) + 1 \right\}. \end{aligned}$$

Consequently, the type $\text{Av}_1((\bar{a}_i)_i/UC)$ is definable over $\bar{a}[I_o]$, for every $C \subseteq \mathbb{M}$. For $C \subseteq \mathbb{M}$, fix a tuple $\bar{a}(C)$ realising $\text{Av}_1((\bar{a}_i)_i/UC \bar{a}[I_o])$. For every C , we have

$$\bar{a}(C) \equiv_{U\bar{a}[I_o]} \bar{a}(\emptyset) \quad \text{and} \quad \bar{a}(C) \stackrel{\text{df}}{\underset{\bar{a}[I_o]}{\nabla}} UC.$$

Consequently,

$$\bar{a}(\emptyset) \stackrel{*}{\underset{\bar{a}[I_o]}{\nabla}} U$$

By Propositions 3.7 and 2.5 it follows that the type

$$\mathfrak{p} := \text{tp}(\bar{a}(\emptyset)/U\bar{a}[I_o]) = \text{Av}_1((\bar{a}_i)_i/U\bar{a}[I_o])$$

is stationary. Since the tuple \bar{a}_i realises $\text{Av}_1((\bar{a}_i)_i/U\bar{a}[I_o]\bar{a}[<i])$, for $i \in I \setminus I_o$,

$$\bar{a}_i \stackrel{*}{\underset{\bar{a}[I_o]}{\nabla}} U\bar{a}[I_o]\bar{a}[<i] \quad \text{implies} \quad \bar{a}_i \underset{\bar{a}[I_o]}{\nabla} U\bar{a}[I_o]\bar{a}[<i],$$

by Propositions 1.5 and F3.1.12. Therefore, it follows by Lemma 5.1 that $(\bar{a}_i)_{i \in I \setminus I_o}$ is a ∇^f -Morley sequence for \mathfrak{p} over $U \cup \bar{a}[I_o]$. \square

As an application, it follows that every indiscernible sequence can be turned into an indiscernible sequence over a larger set if we remove some of its elements.

Lemma 5.4. *Let T be a stable theory and $(\bar{a}_i)_{i \in I}$ and indiscernible sequence over U with $|\bar{a}_i| < \aleph_o$. For every set $C \subseteq \mathbb{M}$ there exists a set $I_o \subseteq I$ of size $|I_o| \leq |C| \otimes \text{loc}_o(\nabla^f)$ such that $(\bar{a}_i)_{i \in I \setminus I_o}$ is indiscernible over $U \cup C \cup \bar{a}[I_o]$.*

Proof. We have seen in Proposition 5.3 that there exist a set $U' \supseteq U$ and a stationary type \mathfrak{p} over U' such that $(\bar{a}_i)_{i \in I}$ is a Morley sequence for \mathfrak{p} over U' . For every finite $C_o \subseteq C$, we can find a set $J(C_o) \subseteq I$ of size $|J(C_o)| < \text{loc}_o(\nabla^f)$ such that

$$C_o \underset{U'\bar{a}[J(C_o)]}{\nabla^f} \bar{a}[I].$$

Setting $I_o := \bigcup \{ J(C_o) \mid C_o \subseteq C \text{ finite} \}$ it follows that

$$|I_o| \leq |C| \otimes \text{loc}_o(\nabla^f) \quad \text{and} \quad C \underset{U'\bar{a}[I_o]}{\nabla^f} \bar{a}[I].$$

Consequently, we can use Lemma 5.2 to show that $(\bar{a}_i)_{i \in I \setminus I_o}$ is a Morley sequence over $U' C \bar{a}[I_o]$. In particular, it is indiscernible over $U C \bar{a}[I_o]$. \square

In totally transcendental theories, it is particularly simple to find Morley sequences.

Definition 5.5. Let ∇ be an abstract independence relation. A family $(A_i)_{i \in I}$ of sets is ∇ -independent over U if

$$A_k \underset{U}{\nabla} \bigcup_{i \neq k} A_i, \quad \text{for all } k \in I.$$

Lemma 5.6. *Let T be a totally transcendental theory and $\mathfrak{p} \in S^{<\omega}(U)$ a type. Every set $I \subseteq \mathfrak{p}^{\mathbb{M}}$ that is ∇^f -independent over U has a finite partition $I = I_o \cup \dots \cup I_{n-1}$ such that each I_i is totally indiscernible over U .*

Proof. We have seen in Corollary 4.8 (b) that $\text{mult}_{\mathbb{J}^f}(\mathfrak{p}) < \aleph_\alpha$. Thus, \mathfrak{p} has only finitely many free extensions q_0, \dots, q_{n-1} over $\text{acl}^{\text{eq}}(U)$. By Lemma 5.1, each set $I_i := I \cap q_i^{\text{M}}$ forms a Morley sequence over U . In particular, it is totally indiscernible. \square

6. The stability spectrum

The *stability spectrum* of a theory T is the class of all cardinals κ such that T is κ -stable. In this section, we will compute the stability spectrum from two parameters: $\text{fc}(\mathbb{J}^f)$ and $\text{st}(T)$. Recall that $\text{fc}(\mathbb{J}^f)$ is the least cardinal κ such that there is no \mathbb{J}^f -forking chain of length κ for a finite set. The cardinal $\text{st}(T)$ is defined as follows.

Definition 6.1. Let T be a complete theory. $\text{st}(T)$ is the minimal infinite cardinal κ such that T is κ -stable. If there is no such cardinal, we set $\text{st}(T) := \infty$.

The following technical lemma contains the main ingredients to determine the stability spectrum of a theory.

Lemma 6.2. *Let T be a stable theory and κ a cardinal.*

- (a) *If $\kappa < \kappa^{<\text{fc}(\mathbb{J}^f)}$, then T is not κ -stable.*
- (b) $\text{fc}(\mathbb{J}^f) \leq |T|^+$.
- (c) $|S^{<\omega}(U)| \leq \text{st}(T) \leq 2^{|T|}$, for every set U of size $|U| \leq \text{st}(T)$.
- (d) $\text{fc}(\mathbb{J}^f) \oplus \text{mult}(\mathbb{J}^f) \leq \text{st}(T)$.
- (e) *If $\kappa \geq \text{st}(T)$ and $\kappa = \kappa^{<\text{fc}(\mathbb{J}^f)}$, then T is κ -stable.*

Proof. (a) Let μ be the least cardinal with $\kappa^\mu > \kappa$. Then $\kappa < \kappa^{<\text{fc}(\mathbb{J}^f)}$ implies that $\mu < \text{fc}(\mathbb{J}^f)$. Hence, there exist a finite tuple \bar{a} and a \mathbb{J}^f -forking chain $(\bar{b}_\alpha)_{\alpha < \mu}$ for \bar{a} over \emptyset of length μ . We construct a tree $(\bar{c}_\eta)_{\eta \in \kappa^{\leq \mu}}$ as follows. We start with $\bar{c}_{\langle \rangle} := \bar{b}_\emptyset$. For the inductive step, suppose that \bar{c}_η is already defined for all $\eta \in \kappa^{< \mu}$ with $|\eta| < \alpha$ and set

$$C_\alpha := \bigcup \{ \bar{c}_\eta \mid \eta \in \kappa^{< \alpha} \}.$$

If α is a limit ordinal, we choose, for every $\eta \in \kappa^\alpha$, a tuple \bar{c}_η with

$$\bar{c}_\eta \bar{c}[\langle \eta \rangle] \equiv \bar{b}_\alpha \bar{b}[\langle \alpha \rangle] \quad \text{and} \quad \bar{c}_\eta \downarrow_{\bar{c}[\langle \eta \rangle]}^f C_\alpha.$$

For the successor step, suppose that $\alpha = \beta + 1$. For each $\eta \in \kappa^\beta$, we choose

- a tuple \bar{d} such that $\bar{d} \bar{c}[\leq \eta] \equiv \bar{b}_\alpha \bar{b}[\langle \alpha \rangle]$,
- a \mathbb{J}^f -Morley sequence $(\bar{c}'_i)_{i < \kappa}$ for $\text{tp}(\bar{d}/\bar{c}[\leq \eta])$ over $\bar{c}[\leq \eta]$, and
- a sequence $(\bar{c}''_i)_{i < \kappa}$ such that

$$\bar{c}''[\langle \kappa \rangle] \equiv_{\bar{c}[\leq \eta]} \bar{c}'[\langle \kappa \rangle] \quad \text{and} \quad \bar{c}''[\langle \kappa \rangle] \downarrow_{\bar{c}[\leq \eta]}^f C_\alpha.$$

Then we set $\bar{c}_{\eta\alpha} := \bar{c}''_\alpha$, for $\alpha < \kappa$.

Having constructed the tree $(\bar{c}_\eta)_{\eta \in \kappa^{< \mu}}$, we set $U := \bigcup_{\eta \in \kappa^{< \mu}} \bar{c}_\eta$. Then $|U| = \kappa^{< \mu} = \kappa$. For each $\zeta \in \kappa^\mu$, let \bar{a}_ζ be a tuple such that

$$\bar{a}_\zeta \bar{c}[\langle \zeta \rangle] \equiv \bar{a} \bar{b}[\langle \mu \rangle] \quad \text{and} \quad \bar{a}_\zeta \downarrow_{\bar{c}[\langle \zeta \rangle]}^f U.$$

We claim that

$$\bar{a}_\xi \not\equiv_U \bar{a}_\zeta, \quad \text{for } \xi \neq \zeta.$$

This implies that $|S^{<\omega}(U)| \geq \kappa^\mu > \kappa = |U|$. Hence, T is not κ -stable.

It remains to prove the claim. Given $\xi \neq \zeta$, let η be the longest common prefix of ξ and ζ and let $\alpha \neq \beta$ be the indices such that $\eta\alpha < \xi$ and $\eta\beta < \zeta$. We start by showing that

$$\bar{c}[\leq \zeta_\alpha] \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}, \quad \text{for all } \zeta_\alpha < \zeta.$$

The proof is by induction on $|\zeta_\alpha|$. Note that we have

$$\bar{c}_{\eta\beta} \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}$$

by choice of $\bar{c}_{\eta\beta}$ and $\bar{c}_{\eta\alpha}$. By (NOR) this implies that

$$\bar{c}[\leq \eta\beta] \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}.$$

Hence, the claim holds for $\zeta_0 \leq \eta\beta$. For the inductive step, let $\zeta_0 > \eta\beta$. Then

$$\bar{c}_{\zeta_0} \downarrow_{\bar{c}[\prec \zeta_0]}^f C_{|\zeta_0|} \text{ implies } \bar{c}_{\zeta_0} \downarrow_{\bar{c}[\prec \zeta_0]}^f \bar{c}_{\eta\alpha}.$$

By inductive hypothesis, we furthermore have

$$\bar{c}[\prec \zeta_0] \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}.$$

Hence, the claim follows by transitivity.

Having proved the claim, it follows by finite character that

$$\bar{c}[\prec \zeta] \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}.$$

Since $\bar{a}_\zeta \downarrow_{\bar{c}[\prec \zeta]}^f \bar{c}_{\eta\alpha}$ this implies by transitivity that

$$\bar{a}_\zeta \downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}.$$

On the other hand,

$$\bar{a} \not\downarrow_{\bar{b}[\leq \alpha]}^f \bar{b}_{\alpha+1} \text{ implies } \bar{a}_\xi \not\downarrow_{\bar{c}[\leq \eta]}^f \bar{c}_{\eta\alpha}.$$

Consequently, $\bar{a}_\xi \not\equiv_{\bar{c}[\leq \eta\alpha]} \bar{a}_\zeta$.

(b) follows from Theorem F3.2.18.

(c) For the upper bound, it is sufficient to note that, according to Theorem C3.5.17, $(2^{|T|})^{|T|} = 2^{|T|}$ implies that T is $2^{|T|}$ -stable.

For the lower bound, let U be a set of size $|U| \leq \text{st}(T)$. Fixing some set $A \supseteq U$ of size $|A| = \text{st}(T)$, it follows by $\text{st}(T)$ -stability of T that

$$|S^{<\omega}(U)| \leq |S^{<\omega}(A)| = |A| = \text{st}(T).$$

(d) We start by showing that $\text{fc}(\downarrow^f) \leq \text{st}(T)$. Note that

$$\kappa < \text{fc}(\downarrow^f) \text{ implies } \kappa < \kappa^\kappa \leq \kappa^{<\text{fc}(\downarrow^f)}.$$

Therefore, it follows by (a) that T is not κ -stable for $\kappa < \text{fc}(\downarrow^f)$. Thus, $\text{st}(T) \geq \text{fc}(\downarrow^f)$.

It therefore remains to prove that $\text{mult}(\downarrow^f) \leq \text{st}(T)$. Let $\mathfrak{p} \in S^{<\omega}(U)$. By Proposition F2.3.24, there exists a set $U_0 \subseteq U$ of size $|U_0| < \text{loc}_0(\downarrow^f) \leq \text{fc}(\downarrow^f)$ such that $\mathfrak{p}|_{U_0} \leq \mathfrak{p}$. Since every free extension of \mathfrak{p} is also a free extension of $\mathfrak{p}|_{U_0}$, it follows that $\text{mult}(\mathfrak{p}|_{U_0}) \geq \text{mult}(\mathfrak{p})$. As T is $\text{st}(T)$ -stable,

$$|U_0| < \text{fc}(\downarrow^f) \leq \text{st}(T) \text{ implies } |S^{\leq\omega}(U_0)| \leq \text{st}(T).$$

Consequently, there exists a model \mathfrak{M} of size $\text{st}(T)$ that contains U_0 . We claim that $|S^{<\omega}(M)| \geq \text{mult}(\mathfrak{p}|_{U_0})$. As T is $\text{st}(T)$ -stable, it then follows that

$$\text{mult}(\mathfrak{p}) \leq \text{mult}(\mathfrak{p}|_{U_0}) \leq |S^{<\omega}(M)| \leq \text{st}(T).$$

To prove the claim, consider a set $C \supseteq U_0$ and let $(q_\alpha)_{\alpha < \lambda}$ be a sequence of distinct free extensions of $\mathfrak{p}|_{U_0}$ over C . For each $\alpha < \lambda$, choose a free extension $q_\alpha^+ \geq q_\alpha$ over $C \cup M$ and set $r_\alpha := q_\alpha^+|_M$. If we can show that $r_\alpha \neq r_\beta$, for $\alpha \neq \beta$, it will follow that $|S^{<\omega}(M)| \geq \lambda$, as desired.

Hence, suppose that $r_\alpha = r_\beta$. Since q_α^+ and q_β^+ are free extensions of the stationary type $r_\alpha = r_\beta$, it follows that $q_\alpha^+ = q_\beta^+$. In particular, $q_\alpha = q_\beta$, which implies that $\alpha = \beta$.

(e) Let $\kappa \geq \text{st}(T)$ be a cardinal such that $\kappa^{<\text{fc}(\downarrow^f)} = \kappa$ and let U be a set of size $|U| \leq \kappa$. By Proposition F2.3.24, we can find, for every type $\mathfrak{p} \in S^{<\omega}(U)$, a set $U_0 \subseteq U$ of size $|U_0| < \text{loc}_0(\downarrow^f) \leq \text{fc}(\downarrow^f)$ such that $\mathfrak{p}|_{U_0} \leq \mathfrak{p}$. As T is $\text{st}(T)$ -stable,

$$|U_0| < \text{fc}(\downarrow^f) \leq \text{st}(T) \text{ implies } |S^{\leq\omega}(U_0)| \leq \text{st}(T).$$

Consequently, it follows as in Theorem 4.5 that

$$\begin{aligned} |S^{<\omega}(U)| &\leq |U|^{<\text{fc}(\downarrow^f)} \otimes \text{st}(T) \otimes \text{mult}(\downarrow^f) \\ &\leq \kappa^{<\text{fc}(\downarrow^f)} \otimes \text{st}(T) \otimes \text{mult}(\downarrow^f) = \kappa, \end{aligned}$$

where the last equality follows by (c) and our choice of κ . \square

Combining statements (a) and (d) of Lemma 6.2 we obtain the following description of the stability spectrum.

Theorem 6.3. *A stable theory T is κ -stable if, and only if,*

$$\kappa = \text{st}(T) \oplus \kappa^{<\text{fc}(\downarrow^f)}.$$

Proof. (\Leftarrow) follows by Lemma 6.2 (d). For (\Rightarrow), suppose that T is κ -stable. By definition, this implies that $\kappa \geq \text{st}(T)$. Furthermore, it follows by Lemma 6.2 (a) that $\kappa \geq \kappa^{<\text{fc}(\downarrow^f)}$. Since the converse inequality $\kappa \leq \kappa^{<\text{fc}(\downarrow^f)}$ is trivial, the claim follows. \square

Let us consider a subclass of stable theories where the stability spectrum is particularly simple.

Definition 6.4. A complete first-order theory T is called *supersimple* if $\text{loc}(\sqrt{f}) \leq \aleph_0$. If T is supersimple and stable, we call it *superstable*.

Note that it follows by Theorem F2.4.17 that every supersimple theory is simple.

Theorem 6.5. *Let T be a complete first-order theory. The following conditions are equivalent.*

- (1) T is supersimple.
- (2) $\text{fc}(\sqrt{f}) \leq \aleph_0$
- (3) $\text{loc}_0(\sqrt{f}) \leq \aleph_0$

Proof. (2) \Leftrightarrow (3) follows by Proposition F2.3.24 and (1) \Leftrightarrow (3) follows by Lemma F2.3.20. \square

Theorem 6.6. *Let T be a complete first-order theory. The following conditions are equivalent.*

- (1) T is superstable.
- (2) T is κ -stable if, and only if, $\kappa \geq \text{st}(T)$.

- (3) *There is a cardinal λ such that T is κ -stable for all $\kappa \geq \lambda$.*

Proof. (2) \Rightarrow (3) is trivial.

(1) \Rightarrow (2) According to Theorem 6.3, T is κ -stable if, and only if,

$$\kappa \geq \text{st}(T) \quad \text{and} \quad \kappa = \kappa^{<\aleph_0}.$$

As the second condition is vacuously true, the claim follows.

(3) \Rightarrow (1) Fix a cardinal $\kappa \geq \lambda$ with $\text{cf } \kappa = \aleph_0$. As $\kappa \geq \lambda \geq \text{st}(T)$, it follows by Theorem 6.3 that $\kappa = \kappa^{<\text{fc}(\downarrow^f)}$. Since $\kappa^{\aleph_0} = \kappa^{\text{cf } \kappa} > \kappa$, this implies that $\text{fc}(\downarrow^f) \leq \aleph_0$. Hence, the claim follows by Theorem 6.5. \square

Corollary 6.7. *Every \aleph_0 -stable theory is superstable.*

Proof. If T is \aleph_0 -stable, then $\text{fc}(\downarrow^f) \leq \text{st}(T) = \aleph_0$ and it follows by Theorem 6.5 that T is supersimple. \square

We conclude this section by noting that, for countable theories, the characterisation in Theorem 6.3 leaves only four possibilities.

Theorem 6.8. *Every countable complete theory T satisfies exactly one of the following conditions:*

- (1) T is totally transcendental. Then

$$T \text{ is } \kappa\text{-stable} \quad \text{iff} \quad \kappa \geq \aleph_0.$$

- (2) T is superstable, but not totally transcendental. Then

$$T \text{ is } \kappa\text{-stable} \quad \text{iff} \quad \kappa \geq 2^{\aleph_0}.$$

- (3) T is stable, but not superstable. Then

$$T \text{ is } \kappa\text{-stable} \quad \text{iff} \quad \kappa = \kappa^{\aleph_0}.$$

- (4) T is unstable.

Proof. Let T be a stable theory. If T is \aleph_0 -stable, we have seen in Theorem c3.5.18 that T is totally transcendental and κ -stable for all infinite κ .

Hence, suppose that T is not \aleph_0 -stable. Then there exists a countable set $U \subseteq \mathbb{M}$ with $|S^{<\omega}(U)| > \aleph_0$. According to Corollary B5.7.5, this implies that $|S^{<\omega}(U)| = 2^{\aleph_0}$. Consequently, $\text{st}(T) \geq 2^{\aleph_0}$. By Lemma 6.2 (c), it follows that $\text{st}(T) = 2^{\aleph_0}$.

Furthermore, we have $\text{fc}(\downarrow^f) \leq \aleph_1$, according to Lemma 6.2 (b). If $\text{fc}(\downarrow^f) = \aleph_0$, then T is superstable and it follows by Theorem 6.6 that

$$T \text{ is } \kappa\text{-stable} \quad \text{iff} \quad \kappa \geq \text{st}(T) = 2^{\aleph_0}.$$

If $\text{fc}(\downarrow^f) = \aleph_1$, Theorem 6.3 implies that T is κ -stable if, and only if, $\kappa \geq \text{st}(T) = 2^{\aleph_0}$ and $\kappa = \kappa^{\aleph_0}$. Note that $\kappa = \kappa^{\aleph_0}$ implies $2^{\aleph_0} \leq \kappa^{\aleph_0} = \kappa$. Hence, the first condition is superfluous and

$$T \text{ is } \kappa\text{-stable} \quad \text{iff} \quad \kappa = \kappa^{\aleph_0}. \quad \square$$

G2. Models of stable theories

1. Isolation relations

In this chapter, we study the structure of models of a stable theory. Our main tool will be a generalisation of the notion of a construction which we introduced in Section E3.4. This generalisation is based on the notion of a so-called isolation relation.

Definition 1.1. (a) A ternary relation \surd on small subsets of \mathbb{M} is an *isolation relation* if it is an abstract independence relation satisfying the axioms (INV), (BMON), and

(RSH) *Right Shift*.

$$AC \surd_U B \quad \text{and} \quad C \surd_U AB \quad \text{implies} \quad A \surd_U BC.$$

If $\bar{a} \surd_U B$, we say that $\text{tp}(\bar{a}/B)$ is \surd -isolated over U . For $U = B$, we sometimes drop the subscript and abbreviate

$$A \surd_U U \quad \text{by} \quad A \surd U.$$

(b) The *left base-monotonicity cardinal* $\text{lbm}(\surd)$ of an isolation relation \surd is the least cardinal κ such that there are sets A, B, C, U with

$$|C| \leq \kappa, \quad AC \surd_U B, \quad \text{and} \quad A \not\surd_{UC} B.$$

If such a cardinal does not exist, we set $\text{lbm}(\surd) = \infty$.

Thus, the difference between an isolation relation and a preforking relation is that we have dropped (DEF) while we have provided a converse to Lemma F2.2.4 by (RSH).

Lemma 1.2. *Every symmetric preforking relation is an isolation relation with $\text{lbm}(\downarrow) = \infty$.*

Proof. By symmetry and base monotonicity it follows that $\text{lbm}(\downarrow) = \infty$. Hence, we only have to prove (RSH). Let $AC \downarrow_U B$ and $C \downarrow_U A$. Then $B \downarrow_U AC$ and Lemma F2.2.4 implies that $BC \downarrow_U A$. Hence, $A \downarrow_U BC$. \square

Our main example of an isolation relation will be the relation $\overset{\text{at}}{\downarrow}$. To illustrate the concept, we also introduce three variants.

Definition 1.3. Let κ be an infinite cardinal.

$$\begin{aligned} A \overset{\text{at}\kappa}{\downarrow}_U B & : \text{iff} \quad \text{for every finite } \bar{a} \subseteq A, \text{ there exists a set} \\ & \quad \Phi \subseteq \text{tp}(\bar{a}/U) \text{ of size } |\Phi| < \kappa \text{ such that} \\ & \quad \mathbb{M} \models \Phi(\bar{a}') \Rightarrow \bar{a}' \equiv_{UB} \bar{a}. \\ A \downarrow_U^{\text{wo}} B & : \text{iff} \quad A' \equiv_U A \Rightarrow A' \equiv_{UB} A. \\ A \downarrow_U^{\text{a}} B & : \text{iff} \quad A' \equiv_{\text{acl}^{\text{eq}}(U)} A \Rightarrow A' \equiv_{UB} A. \end{aligned}$$

Lemma 1.4. (a) $\overset{\text{at}}{\downarrow} \subseteq \overset{\text{at}\kappa}{\downarrow} \subseteq \downarrow^{\text{wo}} \subseteq \downarrow^{\text{a}}$

- (b) $\overset{\text{at}}{\downarrow}$ is an isolation relation with $\text{lbm}(\overset{\text{at}}{\downarrow}) \geq \aleph_0$.
- (c) $\overset{\text{at}\kappa}{\downarrow}$ satisfies all axioms of an isolation relation except for (RSH). For regular cardinals κ , we have $\text{lbm}(\overset{\text{at}\kappa}{\downarrow}) \geq \kappa$.
- (d) \downarrow^{wo} is a symmetric isolation relation with $\text{lbm}(\downarrow^{\text{wo}}) = \infty$.

Proof. (a) The first two inclusions follow immediately from the definitions. For the last one, suppose that $\bar{a} \downarrow_U^{\text{wo}} \bar{b}$. To prove that $\bar{a} \downarrow_U^{\text{a}} \bar{b}$, let $(\bar{b}_n)_{n < \omega}$ be an indiscernible sequence over U with $\bar{b}_0 = \bar{b}$. According to Lemma F3.1.3, it is sufficient to find a tuple $\bar{a}' \equiv_{U\bar{b}} \bar{a}$ such that

$$\bar{b}_m \equiv_{U\bar{a}'} \bar{b}_n, \quad \text{for all } m, n < \omega.$$

We will show in (d) that \downarrow^{wo} is symmetric. Hence, we also have $\bar{b} \downarrow_U^{\text{wo}} \bar{a}$ and

$$\bar{b}_n \equiv_U \bar{b} \quad \text{implies} \quad \bar{b}_n \equiv_{U\bar{a}} \bar{b}, \quad \text{for all } n < \omega.$$

Consequently, the tuple \bar{a} itself has the desired properties.

(b) We have already seen in Lemma F2.3.3 that $\overset{\text{at}}{\downarrow}$ is an abstract independence relation satisfying (INV) and (BMON). The fact that $\text{lbm}(\overset{\text{at}}{\downarrow}) \geq \aleph_0$ follows by (c) for $\kappa = \aleph_0$. Hence, the only axiom that remains to be verified is (RSH).

Suppose that $AC \overset{\text{at}}{\downarrow}_U B$ and $C \overset{\text{at}}{\downarrow}_U AB$. To check that $A \overset{\text{at}}{\downarrow}_U BC$, consider a finite tuple $\bar{a} \subseteq A$. Since $AC \overset{\text{at}}{\downarrow}_U B$, there is a formula $\varphi(\bar{x})$ over U isolating $\text{tp}(\bar{a}/UB)$. We claim that φ also isolates $\text{tp}(\bar{a}/UBC)$. Let \bar{a}' be a tuple satisfying φ . Then $\bar{a}' \equiv_{UB} \bar{a}$ and we have to show that $\bar{a}' \equiv_{UBC} \bar{a}$. Given a finite tuple $\bar{c} \subseteq C$, we fix some tuple \bar{c}' such that

$$\bar{a}'\bar{c} \equiv_{UB} \bar{a}\bar{c}'.$$

Since $\bar{c} \overset{\text{at}}{\downarrow}_U AB$, $\bar{c}' \equiv_U \bar{c}$ implies that $\bar{c}' \equiv_{UB\bar{a}} \bar{c}$. Hence,

$$\bar{a}'\bar{c} \equiv_{UB} \bar{a}\bar{c}' \equiv_{UB} \bar{a}\bar{c}.$$

We have shown that

$$\bar{a}' \equiv_{UB\bar{c}} \bar{a}, \quad \text{for all finite } \bar{c} \subseteq C.$$

Consequently, $\bar{a}' \equiv_{UBC} \bar{a}$.

(c) (INV) and (FIN) follow immediately from the definition.

(BMON) Suppose that $\bar{a} \overset{\text{at}\kappa}{\downarrow}_U BC$. By (FIN) we may assume that \bar{a} is finite. Hence, there exists a set $\Phi(\bar{x}) \subseteq \text{tp}(\bar{a}/U)$ of size $|\Phi| < \kappa$ such that

$$\mathbb{M} \models \Phi(\bar{a}') \quad \text{implies} \quad \bar{a}' \equiv_{UBC} \bar{a}.$$

Since $\Phi \subseteq \text{tp}(\bar{a}/UC)$, the same set shows that $\bar{a} \overset{\text{at}\kappa}{\downarrow}_{UC} B$.

(MON) Suppose that $\bar{a}\bar{c} \text{ at}\sqrt[\kappa]{U} BD$. We claim that $\bar{a} \text{ at}\sqrt[\kappa]{U} B$. Again we may assume that \bar{a} and \bar{c} are finite. According to the definition, there exists a set $\Phi(\bar{x}, \bar{y}) \subseteq \text{tp}(\bar{a}\bar{c}/U)$ of size $|\Phi| < \kappa$ such that

$$\mathbb{M} \models \Phi(\bar{a}', \bar{c}') \quad \text{implies} \quad \bar{a}'\bar{c}' \equiv_{UBD} \bar{a}\bar{c}.$$

We claim that the set

$$\Psi(\bar{x}) := \{ \exists \bar{y} \wedge \Phi_o \mid \Phi_o \subseteq \Phi \text{ finite} \}$$

is the desired witness for $\bar{a} \text{ at}\sqrt[\kappa]{U} B$. Hence, suppose that $\mathbb{M} \models \Psi(\bar{a}')$. By definition of Ψ it follows that, for every finite subset $\Phi_o \subseteq \Phi$, we can find some tuple \bar{c}' with $\mathbb{M} \models \Phi_o(\bar{a}', \bar{c}')$. By compactness, this implies that there is some tuple \bar{c}' with $\mathbb{M} \models \Phi(\bar{a}', \bar{c}')$. Consequently, $\bar{a}'\bar{c}' \equiv_{UBD} \bar{a}\bar{c}$. In particular, we have $\bar{a}' \equiv_{UB} \bar{a}$.

(NOR) Suppose that $A \text{ at}\sqrt[\kappa]{U} B$. To show that $AU \text{ at}\sqrt[\kappa]{U} BU$, let $\bar{a} \subseteq A$ and $\bar{c} = \langle c_0, \dots, c_{n-1} \rangle \subseteq U$ be finite. There exists a set $\Phi(\bar{x}) \subseteq \text{tp}(\bar{a}/U)$ of size $|\Phi| < \kappa$ such that

$$\mathbb{M} \models \Phi(\bar{a}') \quad \text{implies} \quad \bar{a}' \equiv_{UBC} \bar{a}.$$

Setting

$$\Psi(\bar{x}, \bar{y}) := \Phi(\bar{x}) \cup \{y_0 = c_0, \dots, y_{n-1} = c_{n-1}\}$$

it follows that

$$\mathbb{M} \models \Psi(\bar{a}', \bar{c}') \quad \text{implies} \quad \bar{a}' \equiv_{UBC} \bar{a} \quad \text{and} \quad \bar{c}' = \bar{c}.$$

In particular, we have $\bar{a}'\bar{c}' \equiv_{UB} \bar{a}\bar{c}$.

(LRF) Let $A, B \subseteq \mathbb{M}$. To show that $A \text{ at}\sqrt[\kappa]{A} B$, consider a finite tuple $\bar{a} = \langle a_0, \dots, a_{n-1} \rangle \subseteq A$. We set

$$\Phi(\bar{x}) := \{x_0 = a_0, \dots, x_{n-1} = a_{n-1}\}.$$

Then $\mathbb{M} \models \Phi(\bar{a}')$ implies that $\bar{a}' = \bar{a}$. In particular, we have $\bar{a}' \equiv_{AB} \bar{a}$.

(LTR) Suppose that $A_0 A_1 A_2 \text{ at}\sqrt[\kappa]{A_0 A_1} B$ and $A_0 A_1 \text{ at}\sqrt[\kappa]{A_0} B$. By (NOR) it is sufficient to prove that $A_1 A_2 \text{ at}\sqrt[\kappa]{A_0} B$. Hence, let $\bar{a}_1 \subseteq A_1$ and $\bar{a}_2 \subseteq A_2$ be finite. By assumption, there are sets $\Phi(\bar{x}_1, \bar{x}_2) \subseteq \text{tp}(\bar{a}_1 \bar{a}_2 / A_0 A_1)$ and $\Psi(\bar{x}_1) \subseteq \text{tp}(\bar{a}_1 / A_0)$ of size $|\Phi|, |\Psi| < \kappa$ such that

$$\begin{aligned} \mathbb{M} \models \Phi(\bar{a}'_1, \bar{a}'_2) & \quad \text{implies} \quad \bar{a}'_1 \bar{a}'_2 \equiv_{BA_0 A_1} \bar{a}_1 \bar{a}_2, \\ \text{and } \mathbb{M} \models \Psi(\bar{a}'_1) & \quad \text{implies} \quad \bar{a}'_1 \equiv_{BA_0} \bar{a}_1. \end{aligned}$$

Let $U \subseteq A_1$ be the set of parameters from A_1 that are used in Φ and let Φ' be the set of formulae obtained from Φ by replacing each parameter $c \in U$ by a variable y_c . For every finite $\bar{c} \subseteq U$, there exists a set $\Gamma_{\bar{c}}(\bar{y}) \subseteq \text{tp}(\bar{c} / A_0)$ of size $|\Gamma_{\bar{c}}| < \kappa$ such that

$$\mathbb{M} \models \Psi(\bar{c}') \quad \text{implies} \quad \bar{c}' \equiv_{BA_0} \bar{c}.$$

Suppose that $\bar{x}_1 = \langle x_1^0, \dots, x_1^{n-1} \rangle$ and $\bar{a}_1 = \langle a_1^0, \dots, a_1^{n-1} \rangle$. We set

$$\begin{aligned} \mathcal{E}(\bar{x}_1, \bar{x}_2, (y_c)_{c \in U}) & := \Phi' \cup \Psi \cup \bigcup_{\bar{c} \subseteq U} \Gamma_{\bar{c}} \\ & \cup \{x_1^0 = y_{a_1^0}, \dots, x_1^{n-1} = y_{a_1^{n-1}}\}. \end{aligned}$$

Then

$$\mathbb{M} \models \mathcal{E}(\bar{a}'_1, \bar{a}'_2, \bar{c}') \quad \text{implies} \quad \bar{a}'_1 \equiv_{BA_0} \bar{a}_1 \quad \text{and} \quad \bar{c}' \equiv_{BA_0} \bar{c},$$

where \bar{c} is an enumeration of U . Hence, there exists an automorphism $\pi \in \text{Aut } \mathbb{M}_{BA_0}$ with $\pi[\bar{c}'] = \bar{c}$. By the equations added to \mathcal{E} , it follows that $\pi[\bar{a}'_1] = \bar{a}_1$. Consequently,

$$\mathbb{M} \models \mathcal{E}(\bar{a}'_1, \bar{a}'_2, \bar{c}') \quad \text{implies} \quad \mathbb{M} \models \mathcal{E}(\pi[\bar{a}'_1], \pi[\bar{a}'_2], \pi[\bar{c}']).$$

Hence, $\mathbb{M} \models \Phi(\bar{a}_1, \pi[\bar{a}'_2], \bar{c})$, which means that

$$\pi[\bar{a}'_2] \equiv_{BA_0 A_1} \bar{a}_2.$$

Consequently,

$$\bar{a}'_1 \bar{a}'_2 \equiv_{BA_0} \bar{a}_1 \pi[\bar{a}'_2] \equiv_{BA_0} \bar{a}_1 \bar{a}_2.$$

To compute $\text{lbm}(\text{at}\kappa\sqrt{})$, suppose that κ is regular and $AC \text{at}\kappa\sqrt{U} B$ for $|C| < \kappa$. We have to show that $A \text{at}\kappa\sqrt{UC} B$. Hence, let $\bar{a} \subseteq A$ be finite. For every finite tuple $\bar{c} \subseteq C$, there exists a set $\Phi_{\bar{c}}(\bar{x}, \bar{x}') \subseteq \text{tp}(\bar{a}\bar{c}/U)$ of size $|\Phi_{\bar{c}}| < \kappa$ such that

$$\mathbb{M} \models \Phi_{\bar{c}}(\bar{a}', \bar{c}') \quad \text{implies} \quad \bar{a}'\bar{c}' \equiv_{UB} \bar{a}\bar{c}.$$

We set

$$\Psi(\bar{x}) := \bigcup \{ \Phi_{\bar{c}}(\bar{x}, \bar{c}) \mid \bar{c} \subseteq C \text{ finite} \}.$$

Then $|\Psi| < \kappa$ since κ is regular. Furthermore, $\mathbb{M} \models \Psi(\bar{a}')$ implies

$$\bar{a}'\bar{c} \equiv_{UB} \bar{a}\bar{c}, \quad \text{for all finite } \bar{c} \subseteq C.$$

Hence, $\bar{a}'C \equiv_{UB} \bar{a}C$, which implies that $\bar{a}' \equiv_{UBC} \bar{a}$.

(d) (INV) follows immediately from the definition.

(MON) Suppose that $AC \downarrow_U^{\text{wo}} BD$. To show that $A \downarrow_U^{\text{wo}} B$, consider a set $A' \equiv_U A$. Then there exists a set C' such that $A'C' \equiv_U AC$. By assumption, this implies that $A'C' \equiv_{UBD} AC$. In particular, we have $A' \equiv_{UB} A$.

(BMON) Suppose that $A \downarrow_U^{\text{wo}} BC$. To show that $A \downarrow_{UC}^{\text{wo}} B$, consider a set $A' \equiv_{UC} A$. Then $A' \equiv_U A$, which implies that $A' \equiv_{UBC} A'$.

(NOR) Suppose that $A \downarrow_U^{\text{wo}} B$. To show that $AU \downarrow_U^{\text{wo}} BU$, consider sets $A'U' \equiv_U AU$. Then $U' = U$ and $A' \equiv_U A$. Hence, $A' \equiv_{UB} A$, which implies that $A'U' \equiv_{UB} AU$.

(LRF) Let A and B be sets. To show that $A \downarrow_A^{\text{wo}} B$, let $A' \equiv_A A$. Then $A' = A$, which implies that $A' \equiv_{AB} A$.

(SYM) Suppose that $A \downarrow_U^{\text{wo}} B$. To show that $B \downarrow_U^{\text{wo}} A$, consider a set $B' \equiv_U B$. We fix a set A' such that $B'A \equiv_U BA'$. Then $A' \equiv_U A$, which implies that $A' \equiv_{UB} A$. Hence, $B'A \equiv_U BA' \equiv_U BA$, that is, $B' \equiv_{UA} B$.

(LTR) Since we have already proved symmetry, it is sufficient to show that \downarrow^{wo} is right transitive. Hence, suppose that $A \downarrow_{B_0}^{\text{wo}} B_1$ and $A \downarrow_{B_1}^{\text{wo}} B_2$, for $B_0 \subseteq B_1 \subseteq B_2$. To show that $A \downarrow_{B_0}^{\text{wo}} B_2$, consider a set $A' \equiv_{B_0} A$. Then $A' \equiv_{B_1} A$, which implies that $A' \equiv_{B_2} A$.

(FIN) Suppose that $A_0 \downarrow_U^{\text{wo}} B$, for all finite $A_0 \subseteq A$. To show that $A \downarrow_U^{\text{wo}} B$, let $A' \equiv_U A$. Fix an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ with $\pi[A'] = A$ and consider a finite subset $A'_0 \subseteq A'$. By assumption, $A'_0 \equiv_U \pi[A'_0]$ implies that $A'_0 \equiv_{UB} \pi[A'_0]$. Consequently, we have

$$A'_0 \equiv_{UB} \pi[A'_0], \quad \text{for all finite } A'_0 \subseteq A'.$$

This implies that $A' \equiv_{UB} \pi[A'] = A$.

(RSH) Suppose that $AC \downarrow_U^{\text{wo}} B$ and $C \downarrow_U^{\text{wo}} AB$. By symmetry, it follows that $B \downarrow_U^{\text{wo}} AC$. Hence, $B \downarrow_{UC}^{\text{wo}} A$, which implies that $BC \downarrow_{UC}^{\text{wo}} A$. Together with $C \downarrow_U^{\text{wo}} A$ it follows by left transitivity that $BC \downarrow_U^{\text{wo}} A$. Hence, $A \downarrow_U^{\text{wo}} BC$.

Finally, the fact that $\text{lbm}(\downarrow^{\text{wo}}) = \infty$ follows by symmetry and base monotonicity. \square

Exercise 1.1. Show that \downarrow^{a} satisfies all axioms of a symmetric isolation relation except for (LTR) and (RSH). Furthermore, $\text{lbm}(\downarrow^{\text{a}}) = \infty$.

Remark. One can show that, if the theory in question is stable, the relation \downarrow^{a} is also transitive and, thus, an isolation relation.

Recall that we write $\bar{a} \downarrow_U^{\text{l}} B$ if $\text{tp}(\bar{a}/UB)$ is the unique free extension of $\text{tp}(\bar{a}/U)$ over $U \cup B$. This relation will be used in Section 5 below. It inherits some, but not all properties from \downarrow^{f} . In particular, it is a symmetric isolation relation, but not necessarily a forking relation.

Lemma 1.5. *Let T be a stable theory. The relation \downarrow^{l} is a symmetric isolation relation with $\downarrow^{\text{wo}} \subseteq \downarrow^{\text{l}} \subseteq \downarrow^{\text{f}}$.*

Proof. The second inclusion follows immediately from the definition. For the first one, suppose that $\bar{a} \downarrow_U^{\text{wo}} B$. By Lemma 1.4, it follows that $\bar{a} \downarrow_U^{\text{f}} B$. As T is stable, this is equivalent to $\bar{a} \downarrow_U^{\text{l}} B$. For uniqueness, suppose that $\bar{a}' \equiv_U \bar{a}$ is another tuple with $\bar{a}' \downarrow_U^{\text{l}} B$. Then $\bar{a} \downarrow_U^{\text{wo}} B$ implies that $\bar{a}' \equiv_{UB} \bar{a}$. Consequently, $\bar{a} \downarrow_U^{\text{l}} B$.

It remains to prove that \downarrow^{l} is a symmetric isolation relation.

(INV) follows immediately from the definition.

(MON) Suppose that $\bar{a}_0 \bar{a}_1 \downarrow_U^! B$ and let $B_0 \subseteq B$. If $\text{tp}(\bar{a}_0/UB)$ is the unique free extension of $\text{tp}(\bar{a}_0/U)$ over $U \cup B$, then $\text{tp}(\bar{a}_0/UB_0)$ is its unique free extension over $U \cup B_0$. Hence, it is sufficient show that $\bar{a}_0 \bar{a}_1 \downarrow_U^! B$ implies $\bar{a}_0 \downarrow_U^! B$. By monotonicity of $\downarrow^!$, we only need to prove uniqueness.

Consider a tuple $\bar{a}'_0 \equiv_U \bar{a}_0$ with $\bar{a}'_0 \downarrow_U^! B$. We have to show that $\bar{a}'_0 \equiv_{UB} \bar{a}_0$. Choose \bar{a}'_1 such that $\bar{a}'_0 \bar{a}'_1 \equiv_U \bar{a}_0 \bar{a}_1$, and let \bar{a}''_1 be a tuple with

$$\bar{a}''_1 \equiv_{U\bar{a}'_0} \bar{a}'_1 \quad \text{and} \quad \bar{a}''_1 \downarrow_{U\bar{a}'_0}^! B.$$

Since $\bar{a}'_0 \downarrow_U^! B$, transitivity implies that

$$\bar{a}'_0 \bar{a}''_1 \downarrow_U^! B.$$

As $\bar{a}'_0 \bar{a}''_1 \equiv_U \bar{a}'_0 \bar{a}'_1 \equiv_U \bar{a}_0 \bar{a}_1$, it follows that

$$\bar{a}_0 \bar{a}_1 \downarrow_U^! B \quad \text{implies} \quad \bar{a}'_0 \bar{a}''_1 \equiv_{UB} \bar{a}_0 \bar{a}_1.$$

In particular, $\bar{a}'_0 \equiv_{UB} \bar{a}_0$.

(NOR) Suppose that $\bar{a} \downarrow_U^! B$. Then $\bar{a} \downarrow_U^! B$ implies $U\bar{a} \downarrow_U^! BU$. Hence, we only need to prove uniqueness. Let \bar{c} be an enumeration of U and suppose that there are tuples $\bar{a}'\bar{c}' \equiv_U \bar{a}\bar{c}$ such that $\bar{a}'\bar{c}' \downarrow_U^! BU$. Then $\bar{c}' = \bar{c}$, and $\bar{a}' \equiv_U \bar{a}$ implies that $\bar{a}' \equiv_{UB} \bar{a}$. Hence, $\bar{a}'\bar{c}' = \bar{a}'\bar{c} \equiv_{UB} \bar{a}\bar{c}$.

(BMON) Suppose that $\bar{a} \downarrow_U^! BC$. We claim that $\bar{a} \downarrow_{UC}^! B$. Since $\downarrow^!$ is base monotone, we only need to prove uniqueness. Hence, consider a tuple $\bar{a}' \equiv_{UC} \bar{a}$ with $\bar{a}' \downarrow_{UC}^! B$. Then

$$\bar{a}' \equiv_U \bar{a} \quad \text{and} \quad \bar{a} \downarrow_U^! BC \quad \text{implies} \quad \bar{a}' \equiv_{UBC} \bar{a}.$$

(FIN) If $A \downarrow_U^! B$, then (MON) implies that $A_0 \downarrow_U^! B$, for every finite $A_0 \subseteq A$. Conversely, suppose that $A \not\downarrow_U^! B$. If $A \not\downarrow_U^! B$, there is a finite subset $A_0 \subseteq A$ with $A_0 \not\downarrow_U^! B$ and we are done. Hence, suppose that $A \downarrow_U^! B$. Then there is a set $A' \equiv_U A$ such that $A' \downarrow_U^! B$ and $A' \not\equiv_{UB} A$. Let $\pi \in \text{Aut } \mathbb{M}_U$ be an automorphism with $\pi[A] = A'$. Since $A' \not\equiv_{UB} A$,

we can find a finite tuple $\bar{a} \subseteq A$ with $\pi(\bar{a}) \not\equiv_{UB} \bar{a}$. As $\pi(\bar{a}) \equiv_U \bar{a}$ and $\pi(\bar{a}) \downarrow_U^! B$, it follows that $\bar{a} \not\downarrow_U^! B$.

(SYM) Let $\bar{a} \downarrow_U^! \bar{b}$. By symmetry of $\downarrow^!$, we have $\bar{b} \downarrow_U^! \bar{a}$. For uniqueness, consider a tuple $\bar{b}' \equiv_U \bar{b}$ with $\bar{b}' \downarrow_U^! \bar{a}$. There exists an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ with $\pi(\bar{b}') = \bar{b}$. Since $\downarrow^!$ is invariant under automorphisms,

$$\bar{b}' \downarrow_U^! \bar{a} \quad \text{implies} \quad \bar{b} \downarrow_U^! \pi(\bar{a}).$$

By symmetry, $\pi(\bar{a}) \downarrow_U^! \bar{b}$. Since $\pi(\bar{a}) \equiv_U \bar{a}$ and $\bar{a} \downarrow_U^! \bar{b}$, it follows that

$$\pi(\bar{a}) \equiv_{U\bar{b}} \bar{a}.$$

For every formula $\varphi(\bar{x}, \bar{y})$ over U , we therefore have

$$\begin{aligned} \mathbb{M} \models \varphi(\bar{a}, \bar{b}) & \quad \text{iff} \quad \mathbb{M} \models \varphi(\pi(\bar{a}), \bar{b}) \\ & \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}, \pi^{-1}(\bar{b})) \quad \text{iff} \quad \mathbb{M} \models \varphi(\bar{a}, \bar{b}'). \end{aligned}$$

Consequently, $\bar{b}' \equiv_{U\cup\bar{a}} \bar{b}$.

(LTR) As we already have proved symmetry, it is sufficient to show that, for sets $B_0 \subseteq B_1 \subseteq B_2$,

$$\bar{a} \downarrow_{B_0}^! B_1 \quad \text{and} \quad \bar{a} \downarrow_{B_1}^! B_2 \quad \text{implies} \quad \bar{a} \downarrow_{B_0}^! B_2.$$

By transitivity of $\downarrow^!$, we only need to prove uniqueness. Hence, consider a tuple $\bar{a}' \equiv_{B_0} \bar{a}$ with $\bar{a}' \downarrow_{B_0}^! B_2$. Then

$$\bar{a} \downarrow_{B_0}^! B_1 \quad \text{implies} \quad \bar{a}' \equiv_{B_1} \bar{a}.$$

Hence,

$$\bar{a} \downarrow_{B_1}^! B_2 \quad \text{implies} \quad \bar{a}' \equiv_{B_2} \bar{a}.$$

(RSH) Suppose that $AC \downarrow_U^! B$ and $C \downarrow_U^! AB$. Then

$$B \downarrow_U^! AC \quad \text{and} \quad C \downarrow_U^! A,$$

and it follows by Lemma F2.2.4 that $BC \downarrow_U^! A$. Thus, $A \downarrow_U^! BC$. \square

2. Constructions

We can use isolation relations to stratify a structure such that every part is isolated over the previous ones. This leads to a generalised notion of a construction.

Throughout the chapter, we will use the following notation. Given a sequence $(A_\alpha)_{\alpha < \gamma}$ of sets, an ordinal $\alpha \leq \gamma$, and a set $I \subseteq \gamma$, we will write

$$A[I] := \bigcup_{i \in I} A_i, \quad A[<\alpha] := \bigcup_{i < \alpha} A_i, \quad \text{and} \quad A[\leq\alpha] := \bigcup_{i \leq \alpha} A_i.$$

Definition 2.1. Let $\sqrt{}$ be a ternary relation on small subsets of \mathbb{M} and $A, U \subseteq \mathbb{M}$.

(a) A $\sqrt{}$ -stratification of A over U is a sequence $\zeta = (B_\alpha)_{\alpha < \gamma}$ of disjoint sets $B_\alpha \subseteq A$ such that $A = B[<\gamma]$ and

$$B_\alpha \sqrt{UB[<\alpha]}, \quad \text{for all } \alpha < \gamma.$$

(b) A $\sqrt{}$ -stratification ζ is a $\sqrt{}$ -construction if each set B_α is a singleton. In this case, we identify ζ with the corresponding sequence $(b_\alpha)_{\alpha < \gamma}$ of elements $b_\alpha \in B_\alpha$. We say that a set A is $\sqrt{}$ -constructible over U if there exists a $\sqrt{}$ -construction of A over U .

(c) Let $\zeta = (B_\alpha)_{\alpha < \gamma}$ be a $\sqrt{}$ -stratification. The *locality cardinal* $\text{loc}(\zeta)$ of ζ is the least cardinal κ such that, for every $\alpha < \gamma$, there exists a set $C_\alpha \subseteq U \cup B[<\alpha]$ of size $|C_\alpha| < \kappa$ such that

$$B_\alpha \sqrt{C_\alpha} UB[<\alpha].$$

Remark. In this terminology, the kind of constructions introduced in Section E3.4 are $\text{at}\sqrt{}$ -constructions.

We will use $\sqrt{}$ -stratifications to study the structure of $\sqrt{}$ -constructible models. To do so, we will frequently be interested in whether a given subset of a $\sqrt{}$ -constructible set is itself $\sqrt{}$ -constructible. A simple sufficient condition is given by the notion of a *closed set*.

Definition 2.2. Let $\zeta = (B_\alpha)_{\alpha < \gamma}$ be a $\sqrt{}$ -stratification of A . A subset $I \subseteq \gamma$ of indices is ζ -closed over U if

$$B_\alpha \sqrt{UB[I \cap \alpha]} UB[<\alpha], \quad \text{for all } \alpha \in I.$$

Similarly, we call a set $C \subseteq A$ ζ -closed if it is of the form $C = B[I]$, for some ζ -closed set $I \subseteq \gamma$.

Lemma 2.3. Let $\sqrt{}$ be an isolation relation and $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\sqrt{}$ -stratification of A over U . If $I \subseteq \gamma$ is ζ -closed over U , then $(B_\alpha)_{\alpha \in I}$ is a $\sqrt{}$ -stratification of $B[I]$ over U .

Proof. Consider an index $\alpha \in I$. As I is ζ -closed, we have

$$B_\alpha \sqrt{UB[I \cap \alpha]} UB[<\alpha].$$

By (MON), this implies that

$$B_\alpha \sqrt{UB[I \cap \alpha]} UB[I \cap \alpha]. \quad \square$$

In particular, ζ -closed subsets of a $\sqrt{}$ -constructible set are themselves $\sqrt{}$ -constructible. Before proving further properties of ζ -closed sets, let us present a lemma with several ways to construct such sets.

Lemma 2.4. Let $\sqrt{}$ be a relation satisfying (BMON), $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\sqrt{}$ -stratification of A over U , and let $\kappa \geq \text{loc}(\zeta)$ be a regular cardinal.

- (a) A union of ζ -closed sets is ζ -closed.
- (b) If $I \subseteq \gamma$ is ζ -closed over U , then so is $I \cap \beta$ for every $\beta \leq \gamma$.
- (c) Every index $\alpha < \gamma$ is contained in a ζ -closed set $I \subseteq \downarrow \alpha$ over U of size $|I| < \kappa$.
- (d) For every set $C \subseteq A$ of size $|C| < \kappa$, there is some ζ -closed set $I \subseteq \gamma$ over U of size $|I| < \kappa$ with $C \subseteq B[I]$.

Proof. (a) Let \mathcal{I} be a set of ζ -closed sets. To show that $K := \bigcup \mathcal{I}$ is ζ -closed, consider an index $\alpha \in K$. Then $\alpha \in I$, for some ζ -closed set $I \in \mathcal{I}$, and

$$B_\alpha \sqrt{UB[I \cap \alpha]} UB[<\alpha] \text{ implies } B_\alpha \sqrt{UB[K \cap \alpha]} UB[<\alpha]$$

by (bMON).

(b) Let I be ζ -closed and fix $\beta \leq \gamma$. For $\alpha \in I \cap \beta$, we have $I \cap \alpha = (I \cap \beta) \cap \alpha$. Hence,

$$B_\alpha \sqrt{UB[I \cap \alpha]} UB[<\alpha] \text{ implies } B_\alpha \sqrt{UB[(I \cap \beta) \cap \alpha]} UB[<\alpha].$$

(c) We prove the claim by induction on α . There exists a set $C_\alpha \subseteq U \cup B[<\alpha]$ of size $|C_\alpha| < \text{loc}(\zeta) \leq \kappa$ such that

$$B_\alpha \sqrt{C_\alpha} UB[<\alpha].$$

Set $J := \{\beta < \alpha \mid C_\alpha \cap B_\beta \neq \emptyset\}$. By inductive hypothesis, every index $\beta \in J$ is contained in some ζ -closed set $I_\beta \subseteq \downarrow \beta$ of size $|I_\beta| < \kappa$. By (a), the union $I := \bigcup_{\beta \in J} I_\beta$ is also ζ -closed. Since

$$C_\alpha \subseteq U \cup B[J] \subseteq U \cup B[I] = U \cup B[I \cap \alpha],$$

it follows by (bMON) that

$$B_\alpha \sqrt{C_\alpha} UB[<\alpha] \text{ implies } B_\alpha \sqrt{UB[I \cap \alpha]} UB[<\alpha].$$

Since $I \subseteq \alpha$, this implies that $I \cup \{\alpha\}$ is also ζ -closed. Furthermore, $|I \cup \{\alpha\}| < \kappa$, as κ is regular.

(d) Given $C \subseteq A$, set

$$J := \{\alpha < \gamma \mid C \cap B_\alpha \neq \emptyset\}.$$

By (c), every $\alpha \in J$ is contained in some ζ -closed set $K_\alpha \subseteq \gamma$ of size $|K_\alpha| < \kappa$. By (a), the union $I := \bigcup_{\alpha < \gamma} K_\alpha$ is ζ -closed. As κ is regular and $|J| \leq |C| < \kappa$, we have $|I| < \kappa$. Since $C \subseteq B[J] \subseteq B[I]$, the claim follows. \square

The next proposition collects several properties of ζ -closed sets. In particular, we show that a $\sqrt{\quad}$ -stratification over U is also a $\sqrt{\quad}$ -stratification over $U \cup C$, for every ζ -closed set C .

Proposition 2.5. *Let $\sqrt{\quad}$ be an isolation relation and $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\sqrt{\quad}$ -stratification of A over U .*

(a) *If $I \subseteq \gamma$ is ζ -closed over U , then*

$$B[I] \sqrt{UB[I \cap \alpha]} B[<\alpha], \text{ for all } \alpha \leq \gamma.$$

(b) *ζ is a $\sqrt{\quad}$ -stratification of A over $U \cup B[I]$, for every ζ -closed set $I \subseteq \gamma$.*

(c) *If $K \subseteq \gamma$ is a ζ -closed set over U of size $|B[K]| < \text{lmb}(\sqrt{\quad})$, then every set $I \subseteq \gamma$ that is ζ -closed over U is also ζ -closed over $U \cup B[K]$.*

Proof. (a) Fix $\alpha \leq \gamma$. We prove the statement by induction on the minimal ordinal β such that $I \subseteq \beta$. If $I \subseteq \alpha$, then (LRF) and (NOR) imply that

$$B[I] \sqrt{UB[I]} B[<\alpha].$$

As $I = I \cap \alpha$, the claim follows.

For the successor step, suppose that $I = \{\beta\} \cup I_0$ where $I_0 \subseteq \beta$ and $\beta \geq \alpha$. Since I is ζ -closed, we have

$$B_\beta \sqrt{UB[I \cap \beta]} B[<\beta]$$

which, by (MON), implies that

$$B_\beta \sqrt{UB[I_0]B[I \cap \alpha]} B[<\alpha].$$

Furthermore, the set $I_0 = I \cap \beta$ is ζ -closed according to Lemma 2.4 (b). Consequently, the inductive hypothesis yields

$$B[I_0] \sqrt{UB[I_0 \cap \alpha]} B[<\alpha].$$

Since $I_0 \cap \alpha = I \cap \alpha$, it follows by (NOR) and (LTR) that

$$B_\beta B[I_0] \sqrt{UB[I \cap \alpha]} B[<\alpha].$$

Finally, suppose that I has no maximal element. We have seen in Lemma 2.4 (b) that $I \cap \beta$ is ζ -closed, for all $\beta < \gamma$. By inductive hypothesis, it therefore follows that

$$B[I \cap \beta] \sqrt{UB[I \cap \alpha]} B[<\alpha], \quad \text{for all } \beta \in I.$$

Consequently, (FIN) implies that

$$B[I] \sqrt{UB[I \cap \alpha]} B[<\alpha].$$

(b) We have to show that

$$B_\alpha \sqrt{UB[I]B[<\alpha]}, \quad \text{for all } \alpha < \gamma.$$

Hence, let $\alpha < \gamma$. If $\alpha \in I$, the claim follows by (NOR). Thus, suppose that $\alpha \notin I$. By (a), we have

$$B[I] \sqrt{UB[I \cap (\alpha+1)]} B[\leq \alpha].$$

Since $\alpha \notin I$, this implies by (BMON) that

$$B[I] \sqrt{UB[<\alpha]} B[<\alpha] B_\alpha.$$

Hence, (BMON) and (NOR) yield

$$UB[I] B_\alpha \sqrt{UB[<\alpha] B_\alpha} B[<\alpha].$$

As $B_\alpha \sqrt{UB[<\alpha]} B[<\alpha]$, it follows by (NOR) and (LTR) that

$$B[I] B_\alpha \sqrt{UB[<\alpha]} B[<\alpha].$$

Hence, $B[I] \sqrt{UB[<\alpha]} B[<\alpha] B_\alpha$ implies by (RSH) that

$$B_\alpha \sqrt{UB[<\alpha]} B[I] B[<\alpha].$$

Consequently, it follows by (BMON) and (NOR) that

$$B_\alpha \sqrt{UB[I]B[<\alpha]} UB[I] B[<\alpha].$$

(c) Let $I, K \subseteq \gamma$ be ζ -closed over U . Then $I \cup K$ is also ζ -closed. Hence, it follows by (a) that

$$B[I \cup K] \sqrt{UB[(I \cup K) \cap \alpha]} B[<\alpha].$$

For $\alpha \in I$, this implies that

$$B_\alpha B[K] \sqrt{UB[K \cap \alpha] B[I \cap \alpha]} B[<\alpha].$$

Since $|B[K]| < \text{lbn}(\sqrt{\cdot})$, it follows that

$$B_\alpha \sqrt{UB[K] B[I \cap \alpha]} B[<\alpha],$$

as desired. \square

Corollary 2.6. *Let $\sqrt{\cdot}$ be an isolation relation and $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\sqrt{\cdot}$ -stratification of A over U . Then $A \sqrt{UB[<\alpha]}$, for all $\alpha \leq \gamma$.*

Proof. Since the set $I := \gamma$ is ζ -closed, this follows immediately from Proposition 2.5 (a). \square

We conclude this section by presenting conditions for the existence of a stratification or a construction.

Lemma 2.7. *Let $\sqrt{\cdot}$ be an isolation relation, $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\sqrt{\cdot}$ -stratification of A over U , and let $(C_\alpha)_{\alpha < \delta}$ be a sequence of subsets of A such that each set C_α is ζ -closed over $U \cup C[<\alpha]$.*

(a) *The union $C[<\delta]$ is ζ -closed over U .*

(b) $(C_\alpha \setminus C[<\alpha])_{\alpha<\delta}$ is a $\sqrt{\text{-}}$ -stratification of $C[<\delta]$ over U .

Proof. (a) For each $\beta < \delta$, fix a set $I_\beta \subseteq \gamma$ such that $C_\beta = B[I_\beta]$. We prove that $I[<\delta]$ is ζ -closed over U by induction on δ . Consider an index $\alpha \in I[<\delta]$. Then $\alpha \in I_\beta$ for some ordinal $\beta < \delta$. By inductive hypothesis, $I[<\beta]$ is ζ -closed over U . Hence, it follows by Proposition 2.5 (a) that

$$C[<\beta] \sqrt{UB[I[<\beta] \cap \alpha]} B[<\alpha].$$

By (BMON), this implies that

$$C[<\beta] \sqrt{UB[I[<\beta] \cap \alpha] B[I_\beta \cap \alpha]} B[<\alpha].$$

Furthermore, as C_β is ζ -closed over $U \cup C[<\beta]$, we have

$$B_\alpha \sqrt{UC[<\beta] B[I_\beta \cap \alpha]} B[<\alpha].$$

Therefore, it follows by (LTR) that

$$B_\alpha C[<\beta] \sqrt{UB[I[<\beta] \cap \alpha] B[I_\beta \cap \alpha]} B[<\alpha].$$

By (BMON), this implies that

$$B_\alpha \sqrt{UB[I[<\delta] \cap \alpha]} B[<\alpha].$$

(b) Let $\alpha < \delta$. By (a) and Proposition 2.5 (b), ζ is a $\sqrt{\text{-}}$ -stratification of A over $U \cup C[<\alpha]$. As C_α is ζ -closed over $U \cup C[<\alpha]$, it follows therefore by Proposition 2.5 (a) that $C_\alpha \sqrt{UC[<\alpha]}$. \square

Lemma 2.8. *Let $\sqrt{\text{-}}$ be an isolation relation and $(B_\alpha)_{\alpha<\gamma}$ a sequence of sets.*

- (a) *If every B_α is $\sqrt{\text{-}}$ -constructible over $U \cup B[<\alpha]$, then $B[<\gamma]$ is $\sqrt{\text{-}}$ -constructible over U .*
- (b) *Let ζ be a $\sqrt{\text{-}}$ -construction of some set $A \supseteq B[<\gamma]$ over U . If each B_α is ζ -closed over U , then $B[<\gamma]$ is $\sqrt{\text{-}}$ -constructible over U .*

Proof. (a) For each $\alpha < \gamma$, fix a $\sqrt{\text{-}}$ -construction $(b_i^\alpha)_{i<\eta(\alpha)}$ of B_α over $U \cup B[<\alpha]$. Set $\delta(\alpha) := \sum_{i<\alpha} \eta(i)$ and let $(a_\beta)_{\beta<\delta(\gamma)}$ be the concatenation of all sequences $(b_i^\alpha)_i$, for $\alpha < \gamma$, that is,

$$a_{\delta(\alpha)+i} := b_i^\alpha, \quad \text{for } \alpha < \gamma \text{ and } i < \eta(\alpha).$$

To prove that $(a_\beta)_{\beta<\delta(\gamma)}$ is a $\sqrt{\text{-}}$ -construction of $B[<\gamma]$ over U , consider an index $\beta < \delta(\gamma)$. Then $\beta = \delta(\alpha) + i$, for some $\alpha < \gamma$ and $i < \eta(\alpha)$, and

$$b_i^\alpha \sqrt{UB[<\alpha] b^\alpha[<i]} \text{ implies } a_\beta \sqrt{Ua[<\beta]}.$$

(b) According to Lemma 2.7 (a), B_α is $\sqrt{\text{-}}$ -constructible over U . Furthermore, Lemma 2.4 (a) implies that each set of the form $B[<\alpha]$ is ζ -closed over U . Hence, it follows by Proposition 2.5 (b) that B_α is $\sqrt{\text{-}}$ -constructible over $U \cup B[<\alpha]$. Consequently, the claim follows by (a). \square

Lemma 2.9. *Let $\sqrt{\text{-}}$ be an isolation relation. A set A of size $|A| \leq \text{lbm}(\sqrt{\text{-}})$ is $\sqrt{\text{-}}$ -constructible over a set U if, and only if, $A \sqrt{U}$.*

Proof. (\Rightarrow) follows by Corollary 2.6. For (\Leftarrow) , let $A \sqrt{U}$ and let $\zeta = (a_\alpha)_{\alpha<\kappa}$ be an enumeration of A of length $\kappa := |A|$. We claim that ζ is a $\sqrt{\text{-}}$ -construction of A over U . For each $\alpha < \kappa$,

$$A \sqrt{U} \text{ implies } a_\alpha a[<\alpha] \sqrt{U}.$$

Since $|a[<\alpha]| < \kappa \leq \text{lbm}(\sqrt{\text{-}})$, it follows that $a_\alpha \sqrt{Ua[<\alpha]} U$. \square

Corollary 2.10. *Let $\sqrt{\text{-}}$ be an isolation relation and let $(B_\alpha)_{\alpha<\gamma}$ be a $\sqrt{\text{-}}$ -stratification of A over U where*

$$|B_\alpha| \leq \text{lbm}(\sqrt{\text{-}}), \quad \text{for all } \alpha < \gamma.$$

Then A is $\sqrt{\text{-}}$ -constructible over U .

Proof. Since $B_\alpha \sqrt{UB[<\alpha]}$, it follows by Lemma 2.9 that each B_α is $\sqrt{\text{-}}$ -constructible over $U \cup B[<\alpha]$. Consequently, the claim follows by Lemma 2.8 (a). \square

Clearly, if a set A has a $\sqrt{\quad}$ -construction of length γ , then $|A| \leq \gamma < |A|^+$. The next lemma can be used to obtain constructions of length exactly $|A|$.

Lemma 2.11. *Let ζ be a $\sqrt{\quad}$ -construction of A over U . If $\text{loc}(\zeta)^{\text{reg}} \leq |A|$, then A has a $\sqrt{\quad}$ -construction over U of length $|A|$.*

Proof. Let $(a_\alpha)_{\alpha < \kappa}$ be an enumeration of A of length $\kappa := |A|$. By induction on $\alpha < \kappa$, we can use Lemma 2.4 to choose subsets $B_\alpha \subseteq A$ of size $|B_\alpha| < \text{loc}(\zeta)^{\text{reg}}$ such that $a_\alpha \in B_\alpha$ and B_α is ζ -closed over $U \cup B[<\alpha]$. By Lemma 2.3, each set $B_\alpha \setminus B[<\alpha]$ has a $\sqrt{\quad}$ -construction $\xi_\alpha = (b_i^\alpha)_{i < \gamma_\alpha}$ of length

$$\gamma_\alpha < |B_\alpha|^+ \leq \text{loc}(\zeta)^{\text{reg}} \leq \kappa.$$

We have seen in the proof of Lemma 2.8 (a) that the concatenation of these $\sqrt{\quad}$ -constructions is a $\sqrt{\quad}$ -construction of $B[<\kappa] = A$ over U of length $\sum_{\alpha < \kappa} \gamma_\alpha = \kappa$. \square

3. Prime models

Using $\sqrt{\quad}$ -constructions we can generalise the results of Section E3.4 to arbitrary isolation relations. One important property of the relation $\text{at}\sqrt{\quad}$ that is not captured by the notion of an isolation relation is the fact that every model realises all isolated types. When generalising certain results about $\text{at}\sqrt{\quad}$ -constructions we have to require this property separately. This leads to the notion of $\sqrt{\quad}$ -saturation.

Definition 3.1. Let $\sqrt{\quad}$ be an isolation relation, κ a cardinal, and $A, U \subseteq \mathbb{M}$ sets.

(a) A is $\sqrt{\quad}$ - κ -saturated if, for all sets $C \subseteq A$ of size $|C| < \kappa$ and every finite set $B \subseteq \mathbb{M}$ with $B \sqrt{\quad} C$, there is some set $B' \subseteq A$ with $B' \equiv_C B$.

(b) A is $\sqrt{\quad}$ - κ -prime over U if it is $\sqrt{\quad}$ - κ -saturated and, for every $\sqrt{\quad}$ - κ -saturated set $B \supseteq U$, there exists an automorphism $\pi \in \text{Aut } \mathbb{M}_U$ with $\pi[A] \subseteq B$.

Our aim is to prove that $\sqrt{\quad}$ -prime models are unique, up to isomorphism. We start by setting up the required back-and-forth machinery.

Lemma 3.2. *Let A be $\sqrt{\quad}$ - κ -saturated and $U \subseteq A$ a set of size $|U| < \kappa$. For every $\sqrt{\quad}$ -constructible set B over U of size $|B| \leq \kappa$, there exists some set $B' \subseteq A$ with $B' \equiv_U B$.*

Proof. Let $\zeta = (b_\alpha)_{\alpha < \gamma}$ be a $\sqrt{\quad}$ -construction of B over U of length $\gamma \leq \kappa$. We inductively construct a sequence $(b'_\alpha)_{\alpha < \gamma}$ in A such that

$$b'[\alpha] \equiv_U b[\alpha], \quad \text{for all } \alpha \leq \gamma.$$

Suppose that we have already defined b'_α , for all $\alpha < \beta$. Fix an element c such that

$$b_\beta b[<\beta] \equiv_U c b'[\alpha].$$

Then

$$b_\beta \sqrt{\quad} U b[<\beta] \quad \text{implies} \quad c \sqrt{\quad} U b'[\alpha].$$

Since A is $\sqrt{\quad}$ - κ -saturated and $|U| \oplus |\beta| < \kappa$, we can therefore find an element $b'_\beta \in A$ with $b'_\beta \equiv_{U b'[\alpha]} c$. It follows that

$$b'_\beta b'[\alpha] \equiv_U c b'[\alpha] \equiv_U b_\beta b[<\beta]. \quad \square$$

Lemma 3.3. *Let $\sqrt{\quad}$ be an isolation relation, $\kappa \geq \text{lcm}(\sqrt{\quad})$ an uncountable cardinal, $M, N \subseteq \mathbb{M}$ $\sqrt{\quad}$ - κ -saturated, $U \subseteq \mathbb{M}$ a set of size $|U| < \kappa$, and let ξ and ζ be $\sqrt{\quad}$ -constructions of, respectively, M and N over U such that $\text{loc}(\xi)^{\text{reg}}, \text{loc}(\zeta)^{\text{reg}} \leq \text{lcm}(\sqrt{\quad})$. Then*

$$H : M \overset{\kappa}{\cong}_{\text{iso}} N,$$

where H is the set of all elementary maps $f : A \rightarrow B$ such that

- ♦ $A \subseteq M$ is a ξ -closed set of size $|A| < \kappa$,
- ♦ $B \subseteq N$ is a ζ -closed set of size $|B| < \kappa$, and

- ◆ $f \upharpoonright U = \text{id}_U$.

Proof. Set $\lambda := \text{loc}(\xi)^{\text{reg}} \oplus \text{loc}(\zeta)^{\text{reg}}$. By symmetry it is sufficient to check the forth property. Consider a map $f : A \rightarrow B$ in H and an element $c \in M$. We will construct an increasing chain of elementary maps $g_n : C_n \rightarrow D_n$, for $n < \omega$, such that

- ◆ $f \subseteq g_0$ and $c \in \text{dom}(g_0)$,
- ◆ $|C_n \setminus C_{n-1}|, |D_n \setminus D_{n-1}| < \lambda$ (where $C_{-1} := A$ and $D_{-1} := B$),
- ◆ C_n is $\sqrt{\cdot}$ -constructible over $U \cup C_{n-1}$,
- ◆ D_n is $\sqrt{\cdot}$ -constructible over $U \cup D_{n-1}$,
- ◆ C_n is ξ -closed over U , for even $n < \omega$, and
- ◆ D_n is ζ -closed over U , for odd $n < \omega$.

Then we can set $g := \bigcup_{n < \omega} g_n$. By Lemma 2.4 (a),

$$\text{dom}(g) = \bigcup_{n < \omega} C_n = \bigcup_{n < \omega} C_{2n}$$

is ξ -closed and $\text{rng}(g) = \bigcup_{n < \omega} D_{2n+1}$ is ζ -closed. Furthermore,

$$|\text{dom}(g)| < |A|^+ \oplus \lambda \oplus \aleph_1 \leq \kappa.$$

Hence, $g \in H$.

It remains to construct $(g_n)_n$. By Lemma 2.4 (d), we can find a ξ -closed set $C'_0 \subseteq M$ of size $|C'_0| < \lambda$ with $c \in C'_0$. Choose a set $D'_0 \subseteq \mathbb{M}$ with

$$AC'_0 \equiv_U BD'_0.$$

Note that it follows by Proposition 2.5 (b) that C'_0 is $\sqrt{\cdot}$ -constructible over $U \cup A$. Consequently, D'_0 is $\sqrt{\cdot}$ -constructible over $U \cup B$. Since N is $\sqrt{\cdot}$ - κ -saturated, we can therefore use Lemma 3.2 to find a set $D''_0 \subseteq N$ with

$$D''_0 \equiv_{UB} D'_0.$$

Consequently, $AC'_0 \equiv_U BD''_0$. Let $g_0 : A \cup C'_0 \rightarrow B \cup D''_0$ be the corresponding extension of f .

For the successor step, suppose that we have already defined $g_n : C_n \rightarrow D_n$. First, consider the case where n is even. As λ is regular, we can use Lemma 2.4 (d) to find a ζ -closed set $D'_{n+1} \subseteq N$ of size $|D'_{n+1}| < \lambda$ with $D_n \setminus D_{n-1} \subseteq D'_{n+1}$. Choose a set $C'_{n+1} \subseteq \mathbb{M}$ with

$$C_n C'_{n+1} \equiv_U D_n D'_{n+1}.$$

By Proposition 2.5 (b), D'_{n+1} is $\sqrt{\cdot}$ -constructible over $U \cup D_{n-1}$. According to Lemma 2.9 this implies that

$$D'_{n+1} \sqrt{\cdot}_{UD_{n-1}} UD_{n-1}.$$

Since $|D_n \setminus D_{n-1}| < \lambda \leq \text{lcm}(\sqrt{\cdot})$, it follows that

$$D'_{n+1} \sqrt{\cdot}_{UD_n} UD_n.$$

Applying Lemma 2.9 again, we see that the set D'_{n+1} is $\sqrt{\cdot}$ -constructible over $U \cup D_n$. By invariance, it follows that C'_{n+1} is $\sqrt{\cdot}$ -constructible over $U \cup C_n$. Hence, we can use Lemma 3.2 to find a set $C''_{n+1} \subseteq M$ with

$$C''_{n+1} \equiv_{UC_n} C'_{n+1}.$$

Consequently, $C_n C''_{n+1} \equiv_U D_n D'_{n+1}$. Let $g_{n+1} : C_n \cup C''_{n+1} \rightarrow D_n \cup D'_{n+1}$ be the corresponding extension of g_n .

If n is odd, we proceed similarly by choosing a ξ -closed set $C_{n+1} \subseteq M$ containing $C_n \setminus C_{n-1}$. \square

With the back-and-forth machinery in place we can construct isomorphisms between $\sqrt{\cdot}$ - κ -saturated sets.

Proposition 3.4. *Let $\sqrt{\cdot}$ be an isolation relation and $A, B \subseteq \mathbb{M}$ $\sqrt{\cdot}$ - κ -saturated sets that are $\sqrt{\cdot}$ -constructible over some set $U \subseteq A$ of size $|U| < \kappa$.*

- If $|A| \leq \kappa$, there exists an automorphism $\pi : \text{Aut } \mathbb{M}_U$ such that $\pi[A] \subseteq B$.

(b) If $|A|, |B| \leq \kappa$, there exists an automorphism $\pi : \text{Aut } \mathbb{M}_U$ such that $\pi[A] = B$.

Proof. By Lemma 3.3, we have $H : A \equiv_{\text{iso}}^{\kappa} B$. Consequently, we can use Lemma C4.4.10 (a) or (b) to find an elementary embedding $h : A \rightarrow B$ such that $h \upharpoonright U = \text{id}_U$ and, in case $|B| \leq \kappa$, such that h is surjective. As \mathbb{M} is strongly κ^+ -homogeneous, we can extend h to the desired automorphism π . \square

Corollary 3.5. *Let U be a set of size $|U| < \kappa$. Up to isomorphism, there is at most one $\sqrt{-\kappa}$ -saturated set A of size $|A| \leq \kappa$ that is $\sqrt{-}$ -constructible over U .*

Proof. If A and B are $\sqrt{-\kappa}$ -saturated sets of size at most κ that are $\sqrt{-}$ -constructible over U , we can use Proposition 3.4 (b) to find an automorphism mapping A to B . \square

Corollary 3.6. *Let A be a $\sqrt{-\kappa}$ -saturated set of size $|A| \leq \kappa$ that is $\sqrt{-}$ -constructible over a set $U \subseteq A$ of size $|U| < |A|$. Then A is $\sqrt{-\kappa}$ -prime over U .*

Proof. If $B \supseteq U$ is $\sqrt{-\kappa}$ -saturated and $\sqrt{-}$ -constructible over U , we can use Proposition 3.4 (a) to find the desired automorphism mapping A to a subset of B . \square

Having proved that $\sqrt{-\kappa}$ -saturated sets are unique, it remains to show that such sets exist. For the relation $\sqrt[\text{at}]{}$ we will prove below that every model is $\sqrt[\text{at}]{-\kappa}$ -saturated. For isolation relations $\sqrt{-}$ that satisfy the extension axiom, we have the following lemma.

Lemma 3.7. *Let $\sqrt{-}$ be an isolation relation satisfying (EXT) such that $\text{lbm}(\sqrt{-}) \geq \aleph_0$, and let κ be an infinite cardinal. For every set U , there exists some $\sqrt{-\kappa}$ -saturated set A that is $\sqrt{-}$ -constructible over U .*

Proof. We construct an increasing sequence $(A_\alpha)_{\alpha < \kappa^+}$ of sets $A_\alpha \subseteq \mathbb{M}$ such that each A_α is $\sqrt{-}$ -constructible over $U \cup A[\langle \alpha \rangle]$ and, for every

$\alpha < \kappa^+$, every set $C \subseteq A[\langle \alpha \rangle]$ of size $|C| < \kappa$, and every finite set $B \subseteq \mathbb{M}$, there is some set $B' \subseteq A[\leq \alpha]$ with $B' \equiv_C B$.

We start with $A_0 := U$. For the inductive step, suppose that we have already defined A_α , for all $\alpha < \beta$. To find A_β , we fix an enumeration $\langle C_\alpha, \mathfrak{p}_\alpha \rangle_{\alpha < \gamma}$ of all pairs $\langle C, \mathfrak{p} \rangle$ where $C \subseteq A[\langle \beta \rangle]$ has size $|C| < \kappa$ and $\mathfrak{p} \in S^{<\omega}(C)$ is a type such that $\bar{a} \sqrt{-} C$, for every realisation \bar{a} of \mathfrak{p} . By induction on $\alpha < \gamma$, we choose finite tuples $\bar{b}_\alpha \in M^{<\omega}$ as follows. Let \bar{b}'_α be a realisation of \mathfrak{p}_α . Then $\bar{b}'_\alpha \sqrt{-} C$ implies, by (EXT), that there is some tuple $\bar{b}_\alpha \equiv_C \bar{b}'_\alpha$ with

$$\bar{b}_\alpha \sqrt{-}_C A[\langle \beta \rangle] \bar{b}[\langle \alpha \rangle].$$

We set $A_\beta := A[\langle \beta \rangle] \cup \bar{b}[\langle \gamma \rangle]$. Note that $(\bar{b}_\alpha \setminus \bar{b}[\langle \alpha \rangle])_{\alpha < \gamma}$ is a $\sqrt{-}$ -stratification of A_β over $A[\langle \beta \rangle]$. Since $\text{lbm}(\sqrt{-}) \geq \aleph_0$, it follows by Corollary 2.10 that A_β is $\sqrt{-}$ -constructible over $A[\langle \beta \rangle]$.

Having constructed $(A_\alpha)_{\alpha < \kappa^+}$, we claim that the union $A := A[\langle \kappa^+ \rangle]$ has the desired properties. Since every set A_α is $\sqrt{-}$ -constructible over $U \cup A[\langle \alpha \rangle]$, it follows by Lemma 2.8 (a) that A is $\sqrt{-}$ -constructible over U . To show that it is also $\sqrt{-\kappa}$ -saturated, let $C \subseteq A$ be a set of size $|C| < \kappa$ and let $\bar{b} \subseteq \mathbb{M}^{<\omega}$ be a tuple with $\bar{b} \sqrt{-} C$. As κ^+ is regular, there is some index $\alpha < \kappa^+$ such that $C \subseteq A[\langle \alpha \rangle]$. Since the pair $\langle C, \text{tp}(\bar{b}/C) \rangle$ appears in the sequence used in the construction of A_α , it follows that there is some $\bar{b}' \subseteq A_\alpha$ with $\bar{b}' \equiv_C \bar{b}$. \square

4. $\sqrt[\text{at}]{}$ -constructible models

In this section we take a closer look at $\sqrt[\text{at}]{}$ -constructible sets. We have already seen in Section E3.4 that a model which is $\sqrt[\text{at}]{}$ -constructible over some set U is atomic over U , prime over U , and unique up to isomorphisms over U . These facts also follow from the general results we have derived in the present chapter once we have shown that models are always $\sqrt[\text{at}]{}$ -saturated.

Proposition 4.1. *Every model $M \subseteq \mathbb{M}$ is $\overset{\text{at}}{\sqrt{}}$ -saturated.*

Proof. Suppose that $\bar{a} \overset{\text{at}}{\sqrt{}} U$ where $\bar{a} \in M^{<\omega}$ and $U \subseteq M$. Let $\varphi(\bar{x})$ be a formula over U isolating $\text{tp}(\bar{a}/U)$. Then

$$\mathbb{M} \models \varphi(\bar{a}) \quad \text{implies} \quad \mathbb{M} \models \exists \bar{x} \varphi(\bar{x}) \quad \text{implies} \quad \mathfrak{M} \models \exists \bar{x} \varphi(\bar{x}).$$

Hence, there is some tuple $\bar{a}' \in M^{<\omega}$ with $\mathfrak{M} \models \varphi(\bar{a}')$. By choice of φ this implies that $\text{tp}(\bar{a}'/U) = \text{tp}(\bar{a}/U)$. \square

Example. Consider the theory T of the structure $\mathfrak{C} := \langle 2^\omega, (P_n)_{n < \omega} \rangle$ where

$$P_n := \{ \alpha \in 2^\omega \mid \alpha(n) = 1 \}.$$

For this theory, we have

$$A \overset{\text{d}}{\sqrt{}} U B \quad \text{iff} \quad A \cap B \subseteq U,$$

and $A \overset{\text{at}}{\sqrt{}} U B \quad \text{iff} \quad A \subseteq U.$

Furthermore,

$$|S^{<\omega}(U)| \leq |U| \oplus |S^{<\omega}(\emptyset)| = |U| \oplus 2^{\aleph_0},$$

since

$$\text{tp}(\bar{a}/\bar{a} \cap U) \models \text{tp}(\bar{a}/U), \quad \text{for all } \bar{a}, U \subseteq \mathbb{M}.$$

Consequently, the theory T is superstable with $\text{st}(T) = 2^{\aleph_0}$, and a set A is $\overset{\text{at}}{\sqrt{}}$ -constructible over U if, and only if, $A \subseteq U$. In particular, no model of T is $\overset{\text{at}}{\sqrt{}}$ -constructible over \emptyset .

For stable theories one can show that subsets of $\overset{\text{at}}{\sqrt{}}$ -constructible sets are again $\overset{\text{at}}{\sqrt{}}$ -constructible. We start with three technical lemmas.

Lemma 4.2. *Let \downarrow be a symmetric preforking relation. If $A \downarrow_U B$ then, for every set $D \subseteq \mathbb{M}$, there exists a set $C \subseteq A$ of size $|C| < \text{loc}(\downarrow) \oplus |D|^+$ such that*

$$A \downarrow_{UC} BD.$$

Proof. By right locality, we can choose a set $C \subseteq A$ of size

$$|C| < \text{loc}(\downarrow) \oplus |D|^+ \quad \text{such that} \quad D \downarrow_{UBC} UBA.$$

It follows that $D \downarrow_{UBC} A$. Furthermore, $A \downarrow_U B$ implies $B \downarrow_{UC} A$. By transitivity it follows that $BD \downarrow_{UC} A$. \square

Lemma 4.3. *Let \downarrow be a symmetric preforking relation, $\sqrt{\quad}$ an isolation relation, and let $\kappa \geq \text{loc}(\downarrow)$ be a regular cardinal. Let ζ be a $\sqrt{\quad}$ -construction of some set A over U with $\text{loc}(\zeta) \leq \kappa$ and let $C \subseteq A$ be a subset. For every ζ -closed set $B \subseteq A$ with*

$$B \downarrow_{U(C \cap B)} C$$

and every set $D \subseteq A$ of size $|D| < \kappa$, there exists a ζ -closed set $B_+ \supseteq B \cup D$ such that

$$|B_+ \setminus B| < \kappa \quad \text{and} \quad B_+ \downarrow_{U(C \cap B_+)} C.$$

Proof. Let $(d_\alpha)_{\alpha < \gamma}$ be an enumeration of D . Starting with $B_0 := B$, we construct an increasing chain $(B_\alpha)_{\alpha < \gamma}$ of ζ -closed sets such that

$$d_\alpha \in B_{\alpha+1}, \quad |B_\alpha \setminus B| < \kappa, \quad \text{and} \quad B_\alpha \downarrow_{U(C \cap B_\alpha)} C.$$

Then we can set $B_+ := \bigcup_{\alpha < \gamma} B_\alpha$.

For the successor step, suppose that we have already defined B_α . By Lemma 2.4 (c), there exists a ζ -closed set $Z \subseteq A$ of size $|Z| < \kappa$ containing d_α . We can choose a set $W \subseteq U \cup B_\alpha \cup C$ of size

$$|W| < \text{loc}(\downarrow) \oplus |Z|^+ \leq \kappa \quad \text{such that} \quad Z \downarrow_W UB_\alpha C.$$

It follows that $Z \downarrow_{UB_\alpha(W \cap C)} C$. Furthermore,

$$B_\alpha \downarrow_{U(C \cap B_\alpha)} C \text{ implies } B_\alpha \downarrow_{U(C \cap B_\alpha)(C \cap Z)(C \cap W)} C.$$

With transitivity it follows that

$$B_\alpha Z(W \cap C) \downarrow_{U(C \cap B_\alpha)(C \cap Z)(C \cap W)} C.$$

Setting $B_{\alpha+1} := B_\alpha \cup Z \cup (W \cap C)$, we have

$$|B_{\alpha+1} \setminus B| \leq |B_\alpha \setminus B| \oplus |Z| \oplus |W \cap C| < \kappa,$$

as desired.

For the limit step, let δ be a limit ordinal and suppose that B_α is already defined for all $\alpha < \delta$. Then we set $B_\delta := \bigcup_{\alpha < \delta} B_\alpha$. \square

Lemma 4.4. *Let T be a stable theory.*

$$A \sqrt[\text{at}]{} UB \text{ and } A \downarrow_U^f B \text{ implies } A \sqrt[\text{at}]{} U.$$

Proof. Let $\bar{a} \subseteq A$ be finite. We have to show that $\text{tp}(\bar{a}/U)$ is isolated. Since $\bar{a} \sqrt[\text{at}]{} UB$, the type $\mathfrak{p} := \text{tp}(\bar{a}/UB)$ is isolated, that is, the set $\{\mathfrak{p}\}$ is open in $\mathfrak{S}^s(UB)$. By assumption, \mathfrak{p} does not fork over U . Hence, $\mathfrak{p} \in F^s(UB/U)$ in the notation of the Open Mapping Theorem. According to that theorem, the restriction map $\mathfrak{F}^s(UB/U) \rightarrow \mathfrak{S}^s(U)$ is open. Consequently, the image $\{\mathfrak{p}|_U\}$ is open in $\mathfrak{S}^s(U)$ and $\mathfrak{p}|_U$ is isolated. \square

Using these lemmas, we can show that subsets of constructible sets are again constructible.

Theorem 4.5. *Let T be a stable theory with $\text{fc}(\downarrow^f) \leq \aleph_1$ and let A be a $\sqrt[\text{at}]{}-constructible$ set over U . Every subset $C \subseteq A$ is also $\sqrt[\text{at}]{}-constructible$ over U .*

Proof. Let ζ be a $\sqrt[\text{at}]{}-construction$ of A over U . Since

$$\text{loc}(\zeta) \leq \text{loc}_o(\sqrt[\text{at}]{}) \leq \aleph_0$$

and $\text{loc}(\downarrow^f) \leq \text{loc}_o(\downarrow^f)^{\text{reg}} = \text{fc}(\downarrow^f)^{\text{reg}} \leq \aleph_1,$

we can use Lemma 4.3 to inductively construct an increasing sequence $(B_\alpha)_{\alpha < \gamma}$ of sets $B_\alpha \subseteq A$ such that

- ♦ $B_0 = \emptyset$ and $B[<\gamma] = A$,
- ♦ each B_α is ζ -closed over $U \cup B[<\alpha]$,
- ♦ $|B_\alpha \setminus B[<\alpha]| \leq \aleph_0$,
- ♦ $B[<\alpha] \downarrow_{U(C \cap B[<\alpha])}^f C$.

Set $C_\alpha := C \cap (B_\alpha \setminus B[<\alpha])$, for $\alpha < \gamma$. Then $|C_\alpha| \leq \aleph_0 = \text{lbn}(\sqrt[\text{at}]{})$ and, by Corollary 2.10, it is sufficient to prove that $(C_\alpha)_{\alpha < \gamma}$ is a $\sqrt[\text{at}]{}-stratification$ of C over U .

Hence, let $\alpha < \gamma$. Since B_α is ζ -closed over $U \cup B[<\alpha]$, it follows by Lemma 2.8 (a) that $(B_\alpha)_{\alpha < \gamma}$ is a $\sqrt[\text{at}]{}-stratification$ of A over U . Hence,

$$B_\alpha \sqrt[\text{at}]{} UB[<\alpha], \text{ which implies that } C_\alpha \sqrt[\text{at}]{} UB[<\alpha].$$

Since $C_\alpha \downarrow_{UC[<\alpha]}^f B[<\alpha]$, Lemma 4.4 implies that $C_\alpha \sqrt[\text{at}]{} UC[<\alpha]$, as desired. \square

Corollary 4.6. *Let T be a stable theory with $\text{fc}(\downarrow^f) \leq \aleph_1$ and let \mathfrak{M} be a $\sqrt[\text{at}]{}-constructible$ model over U . Then \mathfrak{M} is the unique prime model of T over U .*

Proof. By Corollary 3.6 \mathfrak{M} is prime over U . For uniqueness, let \mathfrak{N} be another prime model over U . Then there exists an elementary embedding $h : \mathfrak{N} \rightarrow \mathfrak{M}$. By Theorem 4.5, \mathfrak{N} is $\sqrt[\text{at}]{}-constructible$ over U . Hence, it follows by Corollary 3.5 that $\mathfrak{N} \cong \mathfrak{M}$. \square

Finally, let us take a look at $\sqrt[\text{at}]{}-constructible$ sets in totally transcendental theories. We will prove below that a model of such a theory is prime if, and only if, it is $\sqrt[\text{at}]{}-constructible$. We also give a characterisation in terms of the length of indiscernible sequences. One direction is contained in the following proposition.

Proposition 4.7. *Let T be totally transcendental and let A be $\sqrt[\text{at}]{}-constructible$ over U . Every indiscernible sequence over U that is contained in A is countable.*

Proof. Let $\zeta = (a_\alpha)_{\alpha < \gamma}$ be a $\sqrt[\text{at}]{}-construction$ of A over U and suppose that A contains an uncountable indiscernible sequence $(c_\alpha)_{\alpha < \omega_1}$ over U . Note that $\text{loc}(\zeta) \leq \text{loc}_o(\sqrt[\text{at}]{} \zeta) = \aleph_o$. Hence, we can use Lemma 2.4 (c) to fix, for every $\alpha < \omega_1$, a finite ζ -closed set $B_\alpha \subseteq A$ over U with $c_\alpha \in B_\alpha$. By Lemma E5.3.11 there exists, for every $\alpha < \omega_1$, an ordinal $\delta_\alpha < \omega_1$ such that $(c_\beta)_{\delta_\alpha \leq \beta < \omega_1}$ is indiscernible over $U \cup B[\langle \alpha \rangle]$. For $\alpha < \omega_1$, set

$$\gamma_o^\alpha := \alpha, \quad \gamma_{n+1}^\alpha := \delta_{\gamma_n^\alpha}, \quad \text{and} \quad \gamma_*^\alpha := \sup_{n < \omega} \gamma_n^\alpha.$$

Note that $\gamma_*^\alpha < \omega_1$, for all $\alpha < \omega_1$. Since $(c_\beta)_{\beta \geq \gamma_*^\alpha}$ is indiscernible over $U \cup B[\langle \gamma_n^\alpha \rangle]$, for all $n < \omega$, it follows that it is indiscernible over

$$\bigcup_{n < \omega} (U \cup B[\langle \gamma_n^\alpha \rangle]) = U \cup B[\langle \gamma_*^\alpha \rangle].$$

Set

$$D_o := \{ \gamma_*^\alpha \mid \alpha < \omega_1 \} \quad \text{and} \quad D := \{ \sup I \mid I \subseteq D_o, \sup I < \omega_1 \}.$$

The set D is closed by definition and it is unbounded since $\gamma_*^\alpha \geq \alpha$, for all α . By Lemma A4.6.8 (a) it follows in particular that D is stationary. Furthermore, for every $\delta \in D$, the suffix $(c_\alpha)_{\delta \leq \alpha < \omega_1}$ is indiscernible over $U \cup B[\langle \delta \rangle]$. By Lemma 2.4 and Proposition 2.5, $B[\langle \delta \rangle]$ is ζ -closed and ζ is a $\sqrt[\text{at}]{}-construction$ over $U \cup B[\langle \delta \rangle]$. By Corollary 2.6, it follows that

$$A \sqrt[\text{at}]{} UB[\langle \delta \rangle], \quad \text{which implies that} \quad c_\delta \sqrt[\text{at}]{} UB[\langle \delta \rangle].$$

Since $\text{loc}(\sqrt[\text{at}]{} \zeta) = \aleph_o$, we can fix, for every $\delta \in D$, a finite set $W_\delta \subseteq U \cup B[\langle \delta \rangle]$ such that

$$c_\delta \sqrt[\text{at}]{}_{W_\delta} UB[\langle \delta \rangle].$$

By the Theorem of Fodor, there exist an index $\delta_o \in D$ and a stationary set $E \subseteq D$ such that $W_\varepsilon \subseteq B[\langle \delta_o \rangle]$, for all $\varepsilon \in E$. Fix two ordinals $\varepsilon < \eta$ in E . By indiscernibility, we have $c_\varepsilon \equiv_{U \cup B[\langle \delta_o \rangle]} c_\eta$. But $W_\eta \subseteq U \cup B[\langle \delta_o \rangle]$ implies that

$$\text{tp}(c_\eta / UB[\langle \delta_o \rangle]) \equiv \text{tp}(c_\eta / UB[\langle \delta_o \rangle] c_\varepsilon).$$

Hence, $c_\eta \neq c_\varepsilon$ implies that $c_\varepsilon \neq c_\varepsilon$. A contradiction. \square

For the converse, we need several lemmas.

Definition 4.8. Let \mathfrak{M} be a model. A set $A \subseteq M$ is *invariant* over $U \subseteq M$ if, for all finite tuples $\bar{a}, \bar{b} \in M^{<\omega}$,

$$\bar{a} \equiv_U \bar{b} \quad \text{implies} \quad \bar{a} \subseteq A \Leftrightarrow \bar{b} \subseteq A.$$

Lemma 4.9. *Let T be a totally transcendental theory, \mathfrak{M} a model of T , and let $U \subseteq M$ be a set such that $M \sqrt[\text{at}]{} U$. Then $M \sqrt[\text{at}]{} UC$ for every set $C \subseteq M$ that is invariant over U .*

Proof. Let $\bar{a} \in M^{<\omega}$. Then the type $\text{tp}(\bar{a}/U)$ is isolated by some formula $\varphi(\bar{x})$ over U . We have seen in Lemma E3.4.12 that the isolated types in $\mathfrak{S}^{<\omega}(UC)$ are dense. Consequently, we can find some isolated type

$$\mathfrak{p} \in \langle \varphi \rangle_{\mathfrak{S}^{<\omega}(UC)}.$$

Let $\psi(\bar{x}, \bar{d})$ be a formula over U isolating \mathfrak{p} with $\bar{d} \subseteq C$ and fix a tuple \bar{b} realising \mathfrak{p} . Then $\mathfrak{M} \models \varphi(\bar{b})$ implies that $\bar{a} \equiv_U \bar{b}$. Consequently, we can find some tuple \bar{c} with $\bar{a}\bar{c} \equiv_U \bar{b}\bar{d}$. Then $\bar{d} \subseteq C$ implies $\bar{c} \subseteq C$ by invariance of C over U . Furthermore,

$$\mathfrak{M} \models \psi(\bar{b}, \bar{d}) \quad \text{implies} \quad \mathfrak{M} \models \psi(\bar{a}, \bar{c}).$$

We claim that $\psi(\bar{x}; \bar{c})$ isolates $\text{tp}(\bar{a}/UC)$. Let $\vartheta(\bar{x}; \bar{c}') \in \text{tp}(\bar{a}/UC)$. Fix some tuple \bar{d}' such that

$$\bar{a}\bar{c}\bar{c}' \equiv_U \bar{b}\bar{d}\bar{d}'.$$

By invariance of C over U , it follows that $\bar{d}' \subseteq U \cup C$. Since $\psi(\bar{x}; \bar{d})$ isolates $\text{tp}(\bar{b}/UC)$ and $\mathfrak{M} \models \vartheta(\bar{a}; \bar{c}')$ implies $\mathfrak{M} \models \vartheta(\bar{b}; \bar{d}')$, we have

$$T(U \cup \bar{d}\bar{d}') \models \psi(\bar{x}; \bar{d}) \rightarrow \vartheta(\bar{x}; \bar{d}').$$

Consequently,

$$T(U \cup \bar{c}\bar{c}') \models \psi(\bar{x}; \bar{c}) \rightarrow \vartheta(\bar{x}; \bar{c}'),$$

as desired. \square

Lemma 4.10. *Let T be a totally transcendental theory, $\mathfrak{M}, \mathfrak{N}$ models of T , $\mathfrak{p} \in S^1(U)$ a type over $U \subseteq M$, and $U \subseteq C \subseteq M$ a set that is invariant over U . If $M \sqrt{\text{at}}/U$ and M does not contain an uncountable indiscernible sequence over U , then every elementary map $f : C \rightarrow N$ can be extended to an elementary map $C \cup \mathfrak{p}^{\mathfrak{M}} \rightarrow N$.*

Proof. We prove the claim by induction on $\alpha := \text{rk}_M(\mathfrak{p})$. If $\alpha = \mathfrak{o}$, then $\mathfrak{p}^{\mathfrak{M}} \subseteq \text{acl}(U)$ and the claim holds trivially.

For the inductive step, suppose that we have proved the statement already for all types of Morley rank less than α . Let $I \subseteq \mathfrak{p}^{\mathfrak{M}}$ be a maximal set such that

$$a \not\downarrow_U^f I \setminus \{a\}, \quad \text{for all } a \in I.$$

According to Lemma G1.5.6, we can partition I into finitely many totally indiscernible sequences. By assumption, each of them is countable. Hence \square so is I . Let $(a_n)_{n < \omega}$ be an enumeration of I and set

$$C_n := C \cup \{c \in M \mid \text{rk}_M(c/Ua[<n]) < \alpha\}, \quad \text{for } n < \omega.$$

Then $(C_n)_{n < \omega}$ forms an increasing chain starting with $C_0 = C$. Furthermore, every element $b \in \mathfrak{p}^{\mathfrak{M}} \setminus \bigcup_{n < \omega} C_n$ satisfies

$$\text{rk}_M(b/UI) = \text{rk}_M(b/U).$$

By Corollary G1.4.8 (a), this implies that $b \not\downarrow_U^f I$. By maximality of I , it therefore follows that

$$\mathfrak{p}^{\mathfrak{M}} \subseteq \bigcup_{n < \omega} C_n.$$

Note that $M \sqrt{\text{at}}/U$ implies $M \sqrt{\text{at}}/Ua[<n]$ since $\text{lcm}(\sqrt{\text{at}}) \geq \aleph_0$. As C_n is invariant over $U \cup a[<n]$, it follows by Lemma 4.9 that $M \sqrt{\text{at}}/C_n$. We prove by induction on n , that f can be extended to C_n . For $n = \mathfrak{o}$, we have $C_0 = C$ and there is nothing to do. For the inductive step, suppose that we have already extended f to C_n . We first extend f to $C_n \cup \{a_n\}$. Since $M \sqrt{\text{at}}/C_n$, there exists some formula $\varphi(x; \bar{c})$ with parameters $\bar{c} \subseteq C_n$ isolating $\text{tp}(a_n/C_n)$. Then

$$\mathfrak{M} \models \exists x \varphi(x; \bar{c}) \quad \text{implies} \quad \mathfrak{N} \models \exists x \varphi(x; f(\bar{c})).$$

Hence, there exists some element $b \in N$ such that we can extend f by setting $f(a_n) := b$. Having done so we obtain the desired extension of f to C_{n+1} by applying the inductive hypothesis on α to C_n (for U) and $C_n \cup \{a_n\}$ (for C). \square

We obtain the following characterisation of prime models of totally transcendental theories.

Theorem 4.11. *Let T be a totally transcendental theory and U a set of parameters.*

- (a) T has a unique prime model over U .
- (b) For a model \mathfrak{M} of T , the following statements are equivalent:
 - (1) \mathfrak{M} is prime over U
 - (2) \mathfrak{M} is $\sqrt{\text{at}}$ -constructible over U .

- (3) \mathfrak{M} is atomic over U and M does not contain an uncountable indiscernible sequence over U .

Proof. (a) The existence and uniqueness of a prime model were already proved in Theorem E3.4.14.

(b) (2) \Rightarrow (1) was shown in Proposition E3.4.3.

(1) \Rightarrow (3) Suppose that \mathfrak{M} is prime over U . By Proposition E3.4.13, there exists a model \mathfrak{A} that is \aleph_1 -constructible over U . As \mathfrak{M} is prime over U , we can find an elementary embedding $h : \mathfrak{M} \rightarrow \mathfrak{A}$ fixing U . By Corollary 2.6, we have $A \aleph_1 \text{ } U$. Since $h[M] \subseteq A$, it follows by invariance that $M \aleph_1 \text{ } U$. Thus, \mathfrak{M} is atomic over U .

Furthermore, every indiscernible sequence in M is mapped by h to an indiscernible sequence in A . We have seen in Proposition 4.7 that A only contains countable indiscernible sequences over U . Hence, so does M .

(3) \Rightarrow (2) Let \mathfrak{N} be an arbitrary model of $T(U)$. To find the desired elementary embedding $\mathfrak{M}_U \rightarrow \mathfrak{N}$, we choose a maximal elementary map $f : C \rightarrow N$ with a domain $C \subseteq M$ that is invariant over U . We claim that $C = M$. For a contradiction, suppose otherwise. Then there is some element $a \in M \setminus C$. By Lemma 4.9, we have $M \aleph_1 \text{ } C$. Hence, setting $\mathfrak{p} := \text{tp}(a/C)$ we can use Lemma 4.10 to extend f to an elementary map $C \cup \mathfrak{p}^{\mathfrak{M}} \rightarrow N$. This contradicts the maximality of f . \square

5. Strongly independent stratifications

In contrast to \aleph_1 , the isolation relation \downarrow^1 admits arbitrarily large constructible sets. Consequently, the length of \downarrow^1 -constructions is unbounded. But we will show that there are \downarrow^1 -stratifications of bounded length such that every enumeration refining them is a \downarrow^1 -construction.

Unique free extensions

As a preliminary step, we start with computing $\text{fc}(\downarrow^1)$. We employ the following characterisation of \downarrow^1 which is a variant of Lemma G1.3.6.

Definition 5.1. For sets $A, B, U \subseteq \mathbb{M}$, we define

$$A \downarrow_U^{\text{do}} B \quad \text{iff} \quad \text{every relation that is definable over both } A \cup U \text{ and } B \cup U \text{ is already definable over } U.$$

If $A \downarrow_U^{\text{do}} B$, we call A and B *definably orthogonal* over U .

Proposition 5.2. Let T be a stable theory. Then

$$A \downarrow_U^f C \quad \text{and} \quad C \downarrow_U^{\text{do}} \text{acl}^{\text{eq}}(U) \quad \text{implies} \quad A \downarrow_U^1 C.$$

Proof. For a contradiction, suppose that $C \downarrow_U^{\text{do}} \text{acl}^{\text{eq}}(U)$ and there are tuples \bar{a} and \bar{b} such that

$$\bar{a} \downarrow_U^f C, \quad \bar{b} \downarrow_U^f C, \quad \text{and} \quad \bar{a} \equiv_U \bar{b},$$

but $\mathfrak{p} := \text{tp}(\bar{a}/UC)$ and $\mathfrak{q} := \text{tp}(\bar{b}/UC)$ are different. By Lemma ??, we can find some $\chi \in \text{FE}(U)$ such that

$$\mathfrak{p}(\bar{x}) \cup \mathfrak{q}(\bar{y}) \models \neg \chi(\bar{x}, \bar{y}).$$

Let E be the set of those equivalence classes $[\bar{c}]_\chi$ containing some tuple $\bar{c}' \in [\bar{c}]_\chi$ realising \mathfrak{p} . Then E is finite and we can choose an enumeration $[\bar{c}_0]_\chi, \dots, [\bar{c}_{m-1}]_\chi$ of E where each representative \bar{c}_i realises \mathfrak{p} . We set

$$\varphi(x) := \bigvee_{i < m} \chi(\bar{x}, \bar{c}_i).$$

Then $\mathbb{M} \models \varphi(\bar{a})$, while

$$\mathfrak{p}(\bar{x}) \cup \mathfrak{q}(\bar{y}) \models \neg \chi(\bar{x}, \bar{y}) \quad \text{implies} \quad \mathbb{M} \models \neg \chi(\bar{c}_i, \bar{b}),$$

for all $i < m$. Hence, $\mathbb{M} \models \varphi(\bar{a}) \wedge \neg\varphi(\bar{b})$ and it is sufficient to prove that φ is equivalent to a formula over U .

Since $\varphi^{\mathbb{M}} \equiv \bigvee_{i < m} \iota_{\chi} \bar{x} = [\bar{c}_i]_{\chi}$ is definable over $\text{acl}^{\text{eq}}(U)$ (recall that the structure \mathbb{M}^{eq} is equipped with projection functions $\iota_{\chi} : \mathbb{M}^{\text{eq}} \rightarrow \mathbb{M}_{\chi}$), it is sufficient to show that it is also definable over C . By Theorem E2.1.11 we only have to check that $\pi[\varphi^{\mathbb{M}}] = \varphi^{\mathbb{M}}$, for all $\pi \in \text{Aut } \mathbb{M}_C$. Hence, consider an automorphism $\pi \in \text{Aut } \mathbb{M}_C$. Then π can be extended to an automorphism of \mathbb{M}_C^{eq} and, therefore, it induces a permutation on the equivalence classes of χ . Since $\pi[\mathfrak{p}^{\mathbb{M}}] = \mathfrak{p}^{\mathbb{M}}$ it follows that π induces a permutation of E . Hence, we have

$$\begin{aligned} \pi[\varphi^{\mathbb{M}}] &= \bigcup_{i < m} \pi[\chi(\bar{x}, \bar{c}_i)]^{\mathbb{M}} = \bigcup_{i < m} \chi(\bar{x}, \pi(\bar{c}_i))^{\mathbb{M}} \\ &= \bigcup_{i < m} \chi(\bar{x}, \bar{c}_i)^{\mathbb{M}} = \varphi^{\mathbb{M}}, \end{aligned}$$

as desired. \square

Proposition 5.3. *Let T be a stable theory.*

- (a) $\text{fc}(\downarrow^f) \leq \text{fc}(\downarrow^!) \leq |T|^+$.
- (b) $\text{fc}(\downarrow^!) \leq \text{fc}(\downarrow^f)^{\text{reg}} \oplus \text{mult}(\downarrow^f)^+$.
- (c) *If T is \aleph_0 -stable, then $\text{fc}(\downarrow^!) \leq \aleph_0$.*

Proof. (a) For the lower bound, note that $\downarrow^! \subseteq \downarrow^f$ implies that every \downarrow^f -forking chain is also a $\downarrow^!$ -forking chain. For the upper bound, we prove that $\text{loc}_o(\downarrow^!) \leq |T|^+$. Since $|T|^+$ is regular, it then follows by Proposition F2.3.24 that $\text{fc}(\downarrow^!) \leq \text{loc}_o(\downarrow^!)^{\text{reg}} \leq |T|^+$.

Hence, consider a finite set A and an arbitrary set B . We construct an increasing sequence $(U_n)_{n < \omega}$ of subsets $U_n \subseteq B$ of size $|U_n| \leq |T|$ such that the union $U := \bigcup_{n < \omega} U_n$ is a set of size $|U| \leq |T|$ with $A \downarrow_U^! B$.

We start with some set $U_0 \subseteq B$ of size $|U_0| < \text{fc}(\downarrow^!) \leq |T|^+$ such that $A \downarrow_{U_0}^f B$. For the inductive step, suppose that U_n is already defined. Note that there are at most

$$|T^{\text{eq}}| \oplus |\text{acl}^{\text{eq}}(U_n)| \leq |T^{\text{eq}}| \oplus |U_n| = |T|$$

formulae over $\text{acl}^{\text{eq}}(U_n)$ and, consequently, at most that many relations that are definable over both B and $\text{acl}^{\text{eq}}(U_n)$. Consequently, we can choose a set $C_n \subseteq B$ of size $|C_n| \leq |T|$ such that every relation definable over both B and $\text{acl}^{\text{eq}}(U_n)$ is definable over C_n . Setting $U_{n+1} := U_n \cup C_n$, it follows that

$$B \perp_{U_{n+1}}^{\text{do}} \text{acl}^{\text{eq}}(U_n) \quad \text{and} \quad |U_{n+1}| = |U_n| \oplus |C_n| \leq |T|.$$

To see that $U := \bigcup_{n < \omega} U_n$ has the desired properties, first note that every relation definable over $\text{acl}^{\text{eq}}(U) = \bigcup_{n < \omega} \text{acl}^{\text{eq}}(U_n)$ is definable over $\text{acl}^{\text{eq}}(U_n)$, for some n , and hence over $U_{n+1} \subseteq U$. Consequently,

$$B \perp_U^{\text{do}} \text{acl}^{\text{eq}}(U)$$

and it follows by Proposition 5.2 that $A \downarrow_U^f B$ implies $A \downarrow_U^! B$.

(b), (c) We prove both bounds simultaneously. Let κ be the least regular cardinal such that

$$\text{mult}_{\downarrow^f}(\mathfrak{p}) < \kappa, \quad \text{for all types } \mathfrak{p}.$$

Then $\text{mult}(\downarrow^f) \leq \kappa \leq \text{mult}(\downarrow^f)^+$. Furthermore, for an \aleph_0 -stable theory, we have $\text{fc}(\downarrow^f)^{\text{reg}} \leq \text{st}(T)^{\text{reg}} = \aleph_0$ and we have seen in Corollary G1.4.8 that $\kappa \leq \aleph_0$. Therefore, both (b) and (c) follow if we can prove that

$$\text{fc}(\downarrow^!) \leq \text{fc}(\downarrow^f)^{\text{reg}} \oplus \kappa.$$

For a contradiction, suppose that we can find some $\downarrow^!$ -forking chain $(B_{\alpha})_{\alpha < \gamma}$ for some finite tuple \bar{a} over \emptyset whose length is $\gamma := \text{fc}(\downarrow^f)^{\text{reg}} \oplus \kappa$. Let $I \subseteq \gamma$ be the set of all indices $\alpha < \gamma$ such that $\bar{a} \not\downarrow_{B[\langle \alpha \rangle]}^f B_{\alpha}$. Then $(B_{\alpha})_{\alpha \in I}$ is a $\downarrow^!$ -forking chain for \bar{a} over \emptyset since

$$A \not\downarrow_{B[\langle \alpha \rangle]}^f B_{\alpha} \quad \text{implies} \quad A \not\downarrow_{B[I] \cap B[\langle \alpha \rangle]}^f B_{\alpha}, \quad \text{for all } \alpha \in I.$$

Hence, $|I| < \text{fc}(\downarrow^!) \leq \gamma$. As γ is regular, it follows that $I \subseteq \beta_0$, for some index $\beta_0 < \gamma$. Note that $\beta_0 + \gamma = \gamma$. Hence, replacing $(B_{\alpha})_{\alpha < \gamma}$ by the subsequence $(B_{\alpha})_{\beta_0 \leq \alpha < \gamma}$ and setting $U := B[\langle \beta_0 \rangle]$, we may assume that

$$\bar{a} \not\downarrow_{UB[\langle \alpha \rangle]}^! B_{\alpha} \quad \text{and} \quad \bar{a} \downarrow_{UB[\langle \alpha \rangle]}^f B_{\alpha}, \quad \text{for all } \alpha < \gamma.$$

We construct a sequence $(\bar{c}_\alpha)_{\alpha < \gamma}$ of tuples as follows. As

$$\bar{a} \not\downarrow_{UB[<\alpha]}^1 B_\alpha \quad \text{and} \quad \bar{a} \downarrow_{UB[<\alpha]}^f B_\alpha,$$

there exists a tuple \bar{c}'_α such that

$$\bar{c}'_\alpha \downarrow_{UB[<\alpha]}^f B_\alpha, \quad \bar{c}'_\alpha \equiv_{UB[<\alpha]} \bar{a}, \quad \text{and} \quad \bar{c}'_\alpha \not\equiv_{UB[<\alpha]B_\alpha} \bar{a}.$$

We choose some tuple $\bar{c}_\alpha \equiv_{UB[<\alpha]B_\alpha} \bar{c}'_\alpha$ such that

$$\bar{c}_\alpha \downarrow_{UB[<\alpha]}^f B[<\gamma].$$

By a straightforward induction on α , one can show that $\bar{a} \downarrow_U^f B[<\alpha]$. Consequently, we have $\bar{c}_\alpha \downarrow_U^f B[<\alpha]$ and it follows by transitivity that $\bar{c}_\alpha \downarrow_U^f B[<\gamma]$. Since

$$\bar{c}_\alpha \not\equiv_{UB[<\alpha+1]} \bar{a} \equiv_{UB[<\alpha+1]} c_{\alpha'}, \quad \text{for all } \alpha < \alpha' < \gamma,$$

the types $\text{tp}(\bar{c}_\alpha/UB[<\gamma])$ are distinct \downarrow^f -free extensions of $\text{tp}(\bar{a}/U)$. By choice of κ , it follows that $\gamma < \kappa$. A contradiction. \square

Strongly independent stratifications

Instead of working with single \downarrow^1 -constructions, we will work with families of them that are encoded by a stratification for the following relation.

Definition 5.4. Let \mathfrak{M} be the model of a stable theory. We define

$$A \overset{\text{si}}{\downarrow}_U B \quad \text{:iff} \quad a \downarrow_U^1 B \cup (A \setminus \{a\}), \quad \text{for all } a \in A.$$

A set A is *strongly independent* over U if $A \overset{\text{si}}{\downarrow}_U U$.

Note that $A \overset{\text{si}}{\downarrow}_U B$ implies that every enumeration of A is a \downarrow^1 -construction over U . It follows that every $\overset{\text{si}}{\downarrow}$ -stratification can be refined to a whole family of \downarrow^1 -constructions.

Lemma 5.5. *If there exists a $\overset{\text{si}}{\downarrow}$ -stratification ζ of A over U , then A is \downarrow^1 -constructible over U .*

Proof. Suppose that $\zeta = (B_\beta)_{\beta < \gamma}$. By Lemma 2.8 (a), it is sufficient to prove that each set B_β is \downarrow^1 -constructible over $U \cup B[<\beta]$. Hence, let $(b_\alpha)_{\alpha < \delta}$ be an arbitrary enumeration of B_β . Then

$$B_\beta \overset{\text{si}}{\downarrow} UB[<\beta] \quad \text{implies} \quad b_\alpha \downarrow_{UB[<\beta]}^1 b[<\alpha], \quad \text{for all } \alpha < \delta. \square$$

Lemma 5.6. *In a stable theory, the relation $\overset{\text{si}}{\downarrow}$ satisfies all axioms of an isolation relation except for (LTR).*

Proof. (INV) follows immediately from the definition.

(MON) Suppose that $AC \overset{\text{si}}{\downarrow}_U BD$. Then

$$a \downarrow_U^1 BD \cup ((A \cup C) \setminus \{a\}), \quad \text{for all } a \in A \cup C,$$

which implies that

$$a \downarrow_U^1 B \cup (A \setminus \{a\}), \quad \text{for all } a \in A.$$

Hence, $A \overset{\text{si}}{\downarrow}_U B$.

(BMON) Suppose that $A \overset{\text{si}}{\downarrow}_U BC$. Then

$$a \downarrow_U^1 BC \cup (A \setminus \{a\}), \quad \text{for all } a \in A,$$

which implies that

$$a \downarrow_{UC}^1 B \cup (A \setminus \{a\}), \quad \text{for all } a \in A.$$

Hence, $A \overset{\text{si}}{\downarrow}_{UC} B$.

(NOR) Suppose that $A \overset{\text{si}}{\downarrow}_U B$. Then

$$a \downarrow_U^1 B \cup (A \setminus \{a\}), \quad \text{for all } a \in A,$$

which implies that

$$a \downarrow_U^! UB \cup (A \setminus \{a\}), \quad \text{for all } a \in A.$$

Since, by (NOR),

$$c \downarrow_U^! UB \cup (A \setminus \{c\}), \quad \text{for all } c \in U,$$

it follows that $AU \overset{\text{si}}{\sqrt{U}} BU$.

(LRF) Let A and B be sets. Since

$$a \downarrow_A^! B \cup (A \setminus \{a\}), \quad \text{for all } a \in A,$$

it follows that $A \overset{\text{si}}{\sqrt{A}} B$.

(FIN) Suppose that $A_{\circ} \overset{\text{si}}{\sqrt{U}} B$, for all finite $A_{\circ} \subseteq A$. To show that $A \overset{\text{si}}{\sqrt{U}} B$, consider an element $a \in A$. For every finite $A_{\circ} \subseteq A \setminus \{a\}$,

$$A_{\circ} a \overset{\text{si}}{\sqrt{U}} B \quad \text{implies} \quad a \downarrow_U^! BA_{\circ}.$$

Since $\downarrow^!$ is a symmetric relation with finite character, it follows that

$$a \downarrow_U^! B(A \setminus \{a\}).$$

(RSH) Suppose that $AC \overset{\text{si}}{\sqrt{U}} B$ and $C \overset{\text{si}}{\sqrt{U}} AB$. In order to show that $A \overset{\text{si}}{\sqrt{U}} BC$, we consider an element $a \in A$. Then

$$AC \overset{\text{si}}{\sqrt{U}} B \quad \text{implies} \quad a \downarrow_U^! B \cup ((A \cup C) \setminus \{a\}).$$

If $a \notin C$, then $B \cup ((A \cup C) \setminus \{a\}) = B \cup C \cup (A \setminus \{a\})$ and we are done. Hence, suppose that $a \in C$. Then

$$C \overset{\text{si}}{\sqrt{U}} AB \quad \text{implies} \quad a \downarrow_U^! AB \cup (C \setminus \{a\}).$$

As $AB \cup (C \setminus \{a\}) = BC \cup (A \setminus \{a\})$, the claim follows. \square

The next lemma is our main tool to construct $\overset{\text{si}}{\sqrt{U}}$ -stratifications.

Lemma 5.7. *Let T be a stable theory and $c \in \mathbb{M}$. Then*

$$A \overset{\text{si}}{\sqrt{U}} B \quad \text{and} \quad c \downarrow_U^! AB \quad \text{implies} \quad Ac \overset{\text{si}}{\sqrt{U}} B.$$

Proof. Let $a \in A$ and set $A_{\circ} := A \setminus \{a\}$. We have to show that

$$a \downarrow_U^! BA_{\circ}c.$$

By symmetry, $c \downarrow_{UBA_{\circ}}^! a$ implies $a \downarrow_{UBA_{\circ}}^! c$. Since $a \downarrow_U^! BA_{\circ}$, it follows by transitivity that $a \downarrow_U^! BA_{\circ}c$, as desired. \square

We are finally able to prove that $\overset{\text{si}}{\sqrt{U}}$ -stratifications always exist and that their length can be bounded.

Theorem 5.8. *Let T be a stable theory. Every set $A \subseteq \mathbb{M}$ has a $\overset{\text{si}}{\sqrt{U}}$ -stratification $\zeta = (B_{\alpha})_{\alpha < \gamma}$ over \emptyset of length $\gamma \leq \text{fc}(\downarrow^!)$.*

Proof. Set $\kappa := \text{fc}(\downarrow^!)$. By induction on α , we choose a sequence $(B_{\alpha})_{\alpha < \kappa}$ of disjoint subsets $B_{\alpha} \subseteq A$ as follows. Suppose that B_{α} has already been defined for all $\alpha < \beta$. Since the union of an increasing chain of strongly independent sets is again strongly independent, we can use the Lemma of Zorn to find a maximal subset $B_{\beta} \subseteq A \setminus B[<\beta]$ such that

$$B_{\beta} \overset{\text{si}}{\sqrt{U}} B[<\beta].$$

The sequence $(B_{\alpha})_{\alpha < \kappa}$ defined in this way is a $\overset{\text{si}}{\sqrt{U}}$ -stratification of $B[<\kappa]$ over \emptyset .

It remains to prove that $B[<\kappa] = A$. For a contradiction, suppose that there is some element $a \in A \setminus B[<\kappa]$. By definition of $\kappa = \text{fc}(\downarrow^!)$ there exists some index $\alpha < \kappa$ such that $a \downarrow_{B[<\alpha]}^! B_{\alpha}$. Since $B_{\alpha} \overset{\text{si}}{\sqrt{B[<\alpha]}} B[<\alpha]$, it follows by Lemma 5.7 that

$$B_{\alpha} a \overset{\text{si}}{\sqrt{B[<\alpha]}} B[<\alpha].$$

This contradicts the maximality of B_{α} . \square

By the special nature of the relation $\sqrt[\text{si}]{} / \sqrt[\text{si}]{} /$ -stratifications can always be refined.

Definition 5.9. Let $(B_\alpha)_{\alpha < \gamma}$ and $(C_\alpha)_{\alpha < \delta}$ be partitions of a set A . We call $(B_\alpha)_{\alpha < \gamma}$ a *refinement* of $(C_\alpha)_{\alpha < \delta}$ if there exists an increasing function $f : \gamma \rightarrow \delta$ such that

$$B_\alpha \subseteq C_{f(\alpha)}, \quad \text{for all } \alpha < \gamma.$$

Lemma 5.10. Let $\zeta = (C_\alpha)_{\alpha < \delta}$ be a $\sqrt[\text{si}]{} /$ -stratification of A over U . Every refinement $(B_\alpha)_{\alpha < \gamma}$ of ζ is also a $\sqrt[\text{si}]{} /$ -stratification of A over U .

Proof. Let $f : \gamma \rightarrow \delta$ be the function such that $B_\alpha \subseteq C_{f(\alpha)}$. To show that $B_\alpha \sqrt[\text{si}]{} / B[<\alpha]$, we consider an element $a \in B_\alpha$. Since

$$C[<f(\alpha)] \subseteq B[<\alpha] \quad \text{and} \quad B[\leq\alpha] \subseteq C[\leq f(\alpha)],$$

it follows by monotonicity of $\downarrow^!$ that

$$a \downarrow_{UC[<f(\alpha)]}^! C[\leq f(\alpha)] \setminus \{a\}$$

implies $a \downarrow_{UB[<\alpha]}^! B[\leq\alpha] \setminus \{a\}$. □

When considering $\sqrt[\text{si}]{} /$ -stratifications as families of $\downarrow^!$ -constructions, we need to modify the notion of a closed set as follows.

Definition 5.11. Let $\zeta = (B_\alpha)_{\alpha < \gamma}$ be a $\sqrt[\text{si}]{} /$ -stratification of A over U .

(a) For every $\alpha < \gamma$ and every element $a \in B_\alpha$, let $W(a) \subseteq U \cup B[<\alpha]$. We call the family $(W(a))_{a \in A}$ a *system of bases* for ζ if

$$a \downarrow_{W(a)}^! U \cup (B[\leq\alpha] \setminus \{a\}), \quad \text{for all } \alpha < \gamma \text{ and all } a \in B_\alpha.$$

(b) Let $(W(a))_{a \in A}$ be a system of bases for ζ . A set $C \subseteq A$ is *W-closed* if $W(a) \subseteq U \cup C$, for all $a \in C$.

Lemma 5.12. Let T be a stable theory and let $\zeta = (B_\alpha)_{\alpha < \gamma}$ be a $\sqrt[\text{si}]{} /$ -stratification of A over U . There exists a system of bases $(W(a))_{a \in A}$ for ζ such that $|W(a)| < \text{loc}_0(\downarrow^!)$, for all $a \in A$.

Proof. For every $a \in B_\alpha$, we choose a set $W(a) \subseteq U \cup B[<\alpha]$ of size $|W(a)| < \text{loc}_0(\downarrow^!)$ such that

$$a \downarrow_{W(a)}^! UB[<\alpha].$$

Since $a \downarrow_{UB[<\alpha]}^! U \cup (B[\leq\alpha] \setminus \{a\})$, it follows by transitivity that

$$a \downarrow_{W(a)}^! U \cup (B[\leq\alpha] \setminus \{a\}). \quad \square$$

Lemma 5.13. Let $(B_\alpha)_{\alpha < \gamma}$ be a $\sqrt[\text{si}]{} /$ -stratification of A over U with system of bases $(W(a))_{a \in A}$. If $C \subseteq A$ is *W-closed*, then $(B_\alpha \cap C)_{\alpha < \gamma}$ is a $\sqrt[\text{si}]{} /$ -stratification of C over U .

Proof. Set $C_\alpha := B_\alpha \cap C$, for $\alpha < \gamma$. Consider an element $a \in C_\alpha$. We have to show that $a \downarrow_{UC[<\alpha]}^! C[\leq\alpha] \setminus \{a\}$. Note that $W(a) \subseteq U \cup C[<\alpha]$ and

$$a \downarrow_{W(a)}^! U \cup (B[\leq\alpha] \setminus \{a\}),$$

by choice of $W(a)$. By monotonicity, it follows that

$$a \downarrow_{UC[<\alpha]}^! U \cup (C[\leq\alpha] \setminus \{a\}),$$

as desired. □

6. Representations

We have introduced indiscernible systems at the end of Section E5.3. Intuitively, if $a : I \rightarrow M$ in an indiscernible system over \mathfrak{J} , the structure \mathfrak{M} is at least as complicated as \mathfrak{J} . If $a : I \rightarrow M$ is surjective, the converse is also true to some extent. In this section, we will characterise theories T by classes \mathcal{C} of structures such that, every model \mathfrak{M} of T has a bijective indiscernible system $a : I \rightarrow M$ with $\mathfrak{J} \in \mathcal{C}$.

Definition 6.1. Let \mathcal{C} be a class of structures.

(a) A *representation* of a structure \mathfrak{M} in \mathcal{C} is a bijective indiscernible system $r : \mathfrak{S} \rightarrow \mathfrak{M}$, for some $\mathfrak{S} \in \mathcal{C}$. If the system $r : \mathfrak{S} \rightarrow \mathfrak{M}$ is only QF-indiscernible, i.e., if

$$\text{atp}(\bar{i}) = \text{atp}(\bar{k}) \quad \text{implies} \quad \text{atp}(r[\bar{i}]) = \text{atp}(r[\bar{k}]),$$

we call r a *quantifier-free representation*.

(b) We say that a theory T has \mathcal{C} -*representations* if every model \mathfrak{M} of T has a representation in \mathcal{C} . Similarly, we say that a class \mathcal{K} has *quantifier-free \mathcal{C} -representations* if every structure $\mathfrak{M} \in \mathcal{K}$ has a quantifier-free representation in \mathcal{C} .

First, let us note that representations are closed under composition.

Lemma 6.2. *If T has \mathcal{K} -representations and \mathcal{K} has quantifier-free \mathcal{C} -representations, then T has \mathcal{C} -representations.*

Lemma 6.3. *If $\mathcal{C} \subseteq \mathcal{K}$, then \mathcal{C} has quantifier-free \mathcal{K} -representations.*

In this section we will characterise stable theories in terms of representations in the following classes.

Definition 6.4. Let κ and λ be cardinals.

(a) $Y_{\kappa\lambda}$ is the signature consisting of unary predicates P_α , for $\alpha < \kappa$, and unary function symbols f_α , for $\alpha < \lambda$.

(b) We denote the class of all $Y_{\kappa\lambda}$ -structures by $\text{Un}(\kappa, \lambda)$ and the subclass consisting of all structures that are *locally finite* by $\text{Lf}(\kappa, \lambda)$. Finally, $\text{Wf}(\kappa, \lambda) \subseteq \text{Un}(\kappa, \lambda)$ is the subclass of all structures such that the inverse R^{-1} of the relation $R := \bigcup_{\alpha < \lambda} f_\alpha$ is *well-founded*.

First, let us give some simple relationships between these classes.

Lemma 6.5. *Let κ and λ be infinite cardinals.*

- (a) $\text{Wf}(\kappa, \lambda)$ has *quantifier-free $\text{Un}(\kappa, \lambda)$ -representations*.
- (b) If $n < \aleph_\circ$, $\text{Wf}(\kappa, n)$ has *quantifier-free $\text{Lf}(\kappa, n)$ -representations*.

(c) If $\kappa_\circ \leq \kappa$ and $\lambda_\circ \leq \lambda$, $\text{Un}(\kappa_\circ, \lambda_\circ)$ has *quantifier-free $\text{Un}(\kappa, \lambda)$ -representations*.

(d) $\text{Un}(\kappa, \lambda)$ has *quantifier-free $\text{Un}(\circ, \lambda \oplus \kappa)$ -representations*.

(e) $\text{Wf}(\kappa, \lambda)$ has *quantifier-free $\text{Wf}(\circ, \lambda \oplus \kappa)$ -representations*.

Proof. (a) follows by Lemma 6.3.

(b) If n is finite, every structure in $\text{Wf}(\kappa, n)$ is locally finite. Therefore $\text{Wf}(\kappa, n) \subseteq \text{Lf}(\kappa, n)$ and the claim follows again by Lemma 6.3.

(c) For $\mathfrak{M} \in \text{Un}(\kappa_\circ, \lambda_\circ)$, we construct a quantifier-free representation $r : I \rightarrow M$ where $\mathfrak{S} \in \text{Un}(\kappa, \lambda)$ is the expansion of \mathfrak{M} by the following functions and relations:

$$\begin{aligned} R_\alpha &:= \emptyset, & \text{for } \kappa_\circ \leq \alpha < \kappa, \\ f_\alpha &:= \text{id}, & \text{for } \lambda_\circ \leq \alpha < \lambda. \end{aligned}$$

It follows that the identity function $\text{id} : M \rightarrow M$ is a quantifier-free indiscernible system over \mathfrak{S} .

(d) Consider a structure $\mathfrak{M} \in \text{Un}(\kappa, \lambda)$. If $|M| \leq 1$, let $\mathfrak{S} \in \text{Un}(\circ, \lambda \oplus \kappa)$ be the unique structure of size $|I| = |M|$ and let $r : I \rightarrow M$ be the corresponding bijection. Then $r : I \rightarrow M$ is a quantifier-free indiscernible system.

If $|M| > 1$, we proceed as follows. Choosing distinct elements $\circ, 1 \in M$ we construct the structure \mathfrak{S} with universe $I := M$ and functions

$$\begin{aligned} f_\alpha^{\mathfrak{S}}(a) &:= f_\alpha^{\mathfrak{M}}(a), & \text{for } \alpha < \lambda, \\ g_\beta^{\mathfrak{S}}(a) &:= \begin{cases} \circ & \text{if } a \notin P_\beta, \\ 1 & \text{if } a \in P_\beta, \end{cases} & \text{for } \beta < \kappa, \\ h^{\mathfrak{S}}(a) &:= \circ. \end{aligned}$$

Note that \mathfrak{S} has $\kappa \oplus \lambda \oplus 1$ functions. Hence, after renaming the functions we obtain a structure in $\text{Un}(\circ, \lambda \oplus \kappa)$. The identity $\text{id} : I \rightarrow M$ is a quantifier-free indiscernible system.

(e) Let $\mathfrak{M} \in \text{Wf}(\kappa, \lambda)$. If $|M| \leq 1$, we proceed as in (d). If $|M| \geq 2$, we modify the construction in (d) as follows. Since $\mathfrak{M} \in \text{Wf}(\kappa, \lambda)$, we can choose $0, 1 \in M$ such that $f_\alpha^{\mathfrak{M}}(0) = 0$ and $f_\alpha^{\mathfrak{M}}(1) \in \{0, 1\}$, for all α . Let \mathfrak{S} be the structure with universe M and functions

$$\begin{aligned} f_\alpha^{\mathfrak{S}}(a) &:= f_\alpha^{\mathfrak{M}}(a), & \text{for } \alpha < \lambda, \\ g_\beta^{\mathfrak{S}}(a) &:= \begin{cases} 0 & \text{if } a \notin P_\beta \text{ or } a = 0, \\ 1 & \text{if } a \in P_\beta \text{ and } a \neq 0, \end{cases} & \text{for } \beta < \kappa, \\ h^{\mathfrak{S}}(a) &:= 0. \end{aligned}$$

After a suitable renaming of the functions, we obtain a structure $\mathfrak{S} \in \text{Wf}(0, \lambda \oplus \kappa)$. Again $\text{id} : I \rightarrow M$ is the desired quantifier-free indiscernible system. \square

Representable theories are stable

We start by showing that theories represented in one of the above classes are stable. The key argument is contained in the following two Ramsey results.

Lemma 6.6. *Let $\mathfrak{M} \in \text{Un}(\kappa, \lambda)$, for infinite cardinals κ and λ , and let $n < \omega$. For every sequence $(\bar{a}^\alpha)_{\alpha < \mu}$ where $\bar{a}^\alpha \in M^n$ and $\mu := (2^{\kappa \oplus \lambda})^+$, there exists an infinite subset $I \subseteq \mu$ such that the subsequence $(\bar{a}^\alpha)_{\alpha \in I}$ is totally QF-indiscernible.*

Proof. Every finitely generated substructure of \mathfrak{M} has size at most λ . By Lemma B1.1.5, there are, up to isomorphism, at most $2^{\kappa \oplus \lambda}$ such substructures. Since $\mu > 2^{\kappa \oplus \lambda}$, there exists a subset $I_0 \subseteq \mu$ of size $|I_0| = \mu$ such that

$$\langle \langle \bar{a}^\alpha \rangle \rangle_{\mathfrak{M}}, \bar{a}^\alpha \cong \langle \langle \bar{a}^\beta \rangle \rangle_{\mathfrak{M}}, \bar{a}^\beta, \quad \text{for all } \alpha, \beta \in I_0.$$

For each $\alpha \in I_0$, we fix an enumeration $\bar{b}^\alpha = (b_i^\alpha)_{i < \gamma}$, without repetitions, of $\langle \langle \bar{a}^\alpha \rangle \rangle_{\mathfrak{M}}$ such that, for all $\alpha, \beta \in I_0$, the map $b_i^\alpha \mapsto b_i^\beta$, $i < \gamma$,

induces an isomorphism

$$\langle \langle \bar{a}^\alpha \rangle \rangle_{\mathfrak{M}}, \bar{a}^\alpha \rightarrow \langle \langle \bar{a}^\beta \rangle \rangle_{\mathfrak{M}}, \bar{a}^\beta.$$

For every $\nu < \mu$, we have $\nu^{<\lambda^+} = \nu^\lambda \leq (2^{\kappa \oplus \lambda})^\lambda = 2^{\kappa \oplus \lambda} < \mu$. Hence, we can use Lemma A4.6.11 to find a subset $U \subseteq M$ and a subset $I_1 \subseteq I_0$ of size $|I_1| = \mu$ such that

$$\bar{b}^\alpha \cap \bar{b}^\beta = U, \quad \text{for all } \alpha \neq \beta \text{ in } I_1.$$

Since $\mu > 2^\lambda$, there exists a subset $I_2 \subseteq I_1$ of size $|I_2| = \mu$ and a set $K \subseteq \lambda$ such that, for all $\alpha \in I_2$,

$$K = \{ i < \gamma \mid b_i^\alpha \in U \}.$$

We claim that the sequence $(\bar{a}^i)_{i \in I_2}$ is totally QF-indiscernible. Let $\varphi(\bar{x})$ be an atomic formula. We have to show that

$$\mathfrak{M} \models \varphi(\bar{a}[\bar{\alpha}]) \leftrightarrow \varphi(\bar{a}[\bar{\beta}]), \quad \text{for all } \bar{\alpha}, \bar{\beta} \subseteq I_2.$$

First, we consider the case where $\varphi = (s = t)$ is an equation. Then there are indices $\xi_0, \dots, \xi_{m-1}, \eta_0, \dots, \eta_{n-1} < \lambda$ and variables x, y such that

$$s(x) = f_{\xi_{m-1}} \cdots f_{\xi_0} x \quad \text{and} \quad t(y) = f_{\eta_{n-1}} \cdots f_{\eta_0} y.$$

For each component $i < n$ of \bar{a}^α , there are indices i' and i'' such that $b_{i'}^\alpha = s(a_i^\alpha)$ and $b_{i''}^\alpha = t(a_i^\alpha)$. Since we have chosen \bar{b}^α without repetitions, it follows that

$$\begin{aligned} \mathfrak{M} \models s(a_i^\alpha) = t(a_k^\beta) & \quad \text{iff} \quad b_{i'}^\alpha = b_{k''}^\beta \\ & \quad \text{iff} \quad i' = k'' \text{ and } (\alpha = \beta \text{ or } i' \in K). \end{aligned}$$

The latter condition is invariant under permutations of I_2 .

It remains to consider the case where $\varphi = P_\zeta t$ for $t = f_{\xi_{m-1}} \cdots f_{\xi_0} x$. Again we can find, for every component i of \bar{a}^α an index i' such that $b_{i'}^\alpha = t(a_i^\alpha)$. Therefore,

$$\mathfrak{M} \models P_\zeta(t(a_i^\alpha)) \quad \text{iff} \quad b_{i'}^\alpha \in P_\zeta.$$

The latter condition does not depend on α , since the substructures induced by the tuples \bar{b}^α are isomorphic. \square

Lemma 6.7. *Let $\mathfrak{M} \in \text{Lf}(\kappa, \lambda)$, let $\mu > \kappa, \lambda$ be an uncountable cardinal, and $n < \omega$. For every sequence $(\bar{a}^\alpha)_{\alpha < \mu}$ of n -tuples $\bar{a}^\alpha \in M^n$ of length μ and every set $U \subseteq M$ of size $|U| < \mu$, there exists a subset $I \subseteq \mu$ of size $|I| = \mu$ such that*

$$\text{atp}(\bar{a}^\alpha/U) = \text{atp}(\bar{a}^\beta/U), \quad \text{for all } \alpha, \beta \in I.$$

Proof. Since each substructure $\langle\langle \bar{a}^\alpha \rangle\rangle_{\mathfrak{M}}$ is finite and there are at most $\kappa \oplus \lambda \oplus \aleph_0 < \mu$ finite substructures, there is a subset $I_0 \subseteq \mu$ of size $|I_0| = \mu$ such that

$$\langle\langle \bar{a}^\alpha \rangle\rangle_{\mathfrak{M}}, \bar{a}^\alpha \cong \langle\langle \bar{a}^\beta \rangle\rangle_{\mathfrak{M}}, \bar{a}^\beta, \quad \text{for all } \alpha, \beta \in I_0.$$

Let $V := \langle\langle U \rangle\rangle_{\mathfrak{M}}$. Since there are at most $|V|^{<\omega} = |V| \oplus \aleph_0 < \mu$ finite subsets of V , we can find a subset $I_1 \subseteq I_0$ of size $|I_1| = \mu$ and a finite set $W \subseteq V$ such that

$$\langle\langle \bar{a}^\alpha \rangle\rangle_{\mathfrak{M}} \cap V = W, \quad \text{for all } \alpha \in I_1.$$

Let \bar{c} be an enumeration of W . There exists a subset $I_2 \subseteq I_1$ of size $|I_2| = \mu$ such that

$$\langle\langle \bar{a}^\alpha \rangle\rangle_{\mathfrak{M}}, \bar{a}^\alpha, \bar{c} \cong \langle\langle \bar{a}^\beta \rangle\rangle_{\mathfrak{M}}, \bar{a}^\beta, \bar{c}, \quad \text{for all } \alpha, \beta \in I_2.$$

It follows that

$$\text{atp}(\bar{a}^\alpha/V) = \text{atp}(\bar{a}^\beta/V), \quad \text{for all } \alpha, \beta \in I_2. \quad \square$$

Using these two lemmas we can show that theories with representations in one of the above classes are stable.

Proposition 6.8. *Let T be a complete first-order theory.*

- (a) *If T has $\text{Un}(\kappa, \lambda)$ -representations, for some cardinals κ, λ , it is stable.*
- (b) *If T has $\text{Lf}(\kappa, \kappa)$ -representations, it is λ -stable, for all $\lambda \geq \kappa \oplus \aleph_0$.*
- (c) *If T has $\text{Lf}(\kappa, \lambda)$ -representations, it is superstable with $\text{st}(T) \leq \kappa \oplus \lambda \oplus \aleph_0$.*

Proof. (a) With out loss of generality, we may assume that κ and λ are infinite. For a contradiction, assume that T is unstable. Then we can use Theorem E5.3.13 to find a model \mathfrak{M} containing an infinite indiscernible sequence $(\bar{a}^i)_{i \in I}$ that is not totally indiscernible. By Lemma E5.3.9, we can find such a sequence where $I = \mu$, for $\mu := (2^{\kappa \oplus \lambda})^+$. Let $r : \mathfrak{S} \rightarrow \mathfrak{M}$ be a representation of \mathfrak{M} in $\text{Un}(\kappa, \lambda)$ and set $\bar{b}^i := r^{-1}(\bar{a}^i)$. By Lemma 6.6, there exists an infinite subset $I_0 \subseteq I$ such that the subsequence $(\bar{b}^i)_{i \in I_0}$ is totally QF-indiscernible. It follows that $(\bar{a}^i)_{i \in I_0}$ is totally indiscernible. As $\text{Av}(\bar{a}^i)_{i \in I_0} = \text{Av}(\bar{a}^i)_{i \in I}$, the sequence $(\bar{a}^i)_{i \in I}$ is also totally indiscernible. A contradiction.

(b) For a contradiction, suppose that $|S^{\bar{s}}(U)| > \lambda$, for some finite tuple \bar{s} of sorts and some set U of parameters of size $|U| = \lambda \geq \kappa \oplus \aleph_0$. Fix a sequence $(\bar{a}^\alpha)_{\alpha < \lambda^+}$ such that

$$\text{tp}(\bar{a}^\alpha/U) \neq \text{tp}(\bar{a}^\beta/U), \quad \text{for all } \alpha \neq \beta.$$

Let $r : \mathfrak{S} \rightarrow \mathfrak{M}$ be a representation of \mathfrak{M} in $\text{Lf}(\kappa, \kappa)$. We set

$$V := \langle\langle r^{-1}[U] \rangle\rangle_{\mathfrak{S}} \quad \text{and} \quad \bar{b}^\alpha := r^{-1}(\bar{a}^\alpha), \quad \text{for } \alpha < \lambda^+.$$

By Lemma 6.7, there exists a subset $I \subseteq \lambda^+$ of size $|I| = \lambda^+$ such that

$$\text{atp}(\bar{b}^\alpha/V) = \text{atp}(\bar{b}^\beta/V), \quad \text{for all } \alpha, \beta \in I.$$

Since r is a representation, it follows that

$$\text{tp}(\bar{a}^\alpha/U) = \text{tp}(\bar{a}^\beta/U), \quad \text{for all } \alpha, \beta \in I,$$

in contradiction to our choice of $(\bar{a}^\alpha)_\alpha$.

(c) According to Theorem G1.6.6, a theory T is superstable if, and only if, there exists some cardinal λ such that T is κ -stable, for all $\kappa \geq \lambda$. Consequently, the claim follows from (b). \square

Stable theories have representations

For the converse statements, we employ $\text{si}/$ -stratifications. The following two technical lemmas contain the key argument.

Lemma 6.9. *Let T be stable, $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\text{si}/$ -stratification of some set M over \emptyset such that $|B_0| = |B_1| = 1$, and let $(W(a))_{a \in M}$ be a system of bases for ζ .*

Suppose that $p : C \rightarrow D$ is a bijective function with $C, D \subseteq M$ satisfying the following conditions:

- ◆ $p[B_\alpha] \subseteq B_\alpha$, for all $\alpha < \gamma$.
- ◆ C is W -closed.
- ◆ $p[W(a)] = W(p(a))$, for all $a \in C$.
- ◆ $p[\text{tp}(a/W(a))] = \text{tp}(p(a)/W(p(a)))$, for all $a \in C$.

Then p is an elementary map.

Proof. Set $C_\alpha := C \cap B_\alpha$ and $D_\alpha := D \cap B_\alpha$. By assumption, $p[C_\alpha] = D_\alpha$. We will show by induction on $\alpha < \gamma$ that

$$p[\text{tp}(\bar{a}/C[<\alpha])] = \text{tp}(p(\bar{a})/D[<\alpha]), \quad \text{for all finite } \bar{a} \subseteq C_\alpha.$$

For $\alpha < 2$, the claim holds trivially since $|B_\alpha| = 1$. For $\alpha \geq 2$, we prove the statement by induction on $|\bar{a}|$. Hence, suppose that $\bar{a} = b\bar{c}$ and that we have already shown that

$$p[\text{tp}(\bar{c}/C[<\alpha])] = \text{tp}(p(\bar{c})/D[<\alpha]).$$

By assumption, we have

$$W(p(b)) = p[W(b)],$$

$$\text{and } \text{tp}(p(b) / W(p(b))) = p(\text{tp}(b/W(b))).$$

Since $b \downarrow_{W(b)}^! B[\leq \alpha] \setminus \{b\}$, the type $\mathfrak{p} := \text{tp}(b/C[<\alpha]\bar{c})$ is the unique free extension of $\text{tp}(b/W(b))$. As we have already shown that the map $p \upharpoonright C[<\alpha]\bar{c}$ is elementary, it follows that the image $p(\mathfrak{p})$ does not fork over $p[W(b)] = W(p(b))$. Since

$$\text{tp}(p(b)/W(p(b))) = \text{tp}(p(b)/p[W(b)]) = p(\text{tp}(b/W(b)))$$

and $p(b) \downarrow_{W(p(b))}^! D[<\alpha]p(\bar{c})$, it follows that

$$p(\mathfrak{p}) = \text{tp}(p(b) / D[<\alpha]p(\bar{c})).$$

Consequently,

$$p(\text{tp}(b\bar{c}/C[<\alpha])) = \text{tp}(p(b\bar{c})/D[<\alpha]). \quad \square$$

Lemma 6.10. *Let \mathfrak{M} be a structure, $\zeta = (B_\alpha)_{\alpha < \gamma}$ a $\text{si}/$ -stratification of M over \emptyset such that $|B_0| = |B_1| = 1$, and let $(W(a))_{a \in M}$ be a system of bases for ζ . If*

$$\mathfrak{S} := \langle M, (f_\alpha)_{\alpha < \kappa}, (P_\alpha)_{\alpha < \lambda}, (Q_\alpha)_{\alpha < \mu} \rangle$$

is a structure such that

- ◆ $\{f_\alpha(a) \mid \alpha < \kappa\} = \{a\} \cup W(a)$, for all $a \in M$,
- ◆ $a \in P_\beta \Leftrightarrow b \in P_\beta$, for all $\beta < \lambda$, implies that $a \in B_\alpha \Leftrightarrow b \in B_\alpha$, for all $\alpha < \gamma$,
- ◆ $a \in Q_\beta \Leftrightarrow b \in Q_\beta$, for all $\beta < \mu$, implies that

$$a(f_\alpha(a))_{\alpha < \kappa} \equiv b(f_\alpha(b))_{\alpha < \kappa},$$

then the identity map $\text{id} : M \rightarrow M$ is a representation of \mathfrak{M} in \mathfrak{S} .

Proof. Let $\bar{a}, \bar{b} \subseteq M$ be finite tuples such that

$$\langle\langle \bar{a} \rangle\rangle_{\mathfrak{S}, \bar{a}} \cong \langle\langle \bar{b} \rangle\rangle_{\mathfrak{S}, \bar{b}}.$$

We have to show that $\bar{a} \equiv \bar{b}$ in \mathfrak{M} . Let $p : \langle\langle \bar{a} \rangle\rangle_{\mathfrak{S}} \rightarrow \langle\langle \bar{b} \rangle\rangle_{\mathfrak{S}}$ be an isomorphism with $p(\bar{a}) = \bar{b}$. It is sufficient to prove that p satisfies the conditions of Lemma 6.9.

By assumption on the predicates P_β we have

$$p(c) \in B_\alpha \quad \text{iff} \quad c \in B_\alpha.$$

Hence, $p[B_\alpha] \subseteq B_\alpha$. Furthermore, if $c \in \text{dom}(p) = \langle\langle \bar{a} \rangle\rangle_{\mathfrak{S}}$, then

$$\{c\} \cup W(c) = \{f_\alpha(c) \mid \alpha < \kappa\} \subseteq \langle\langle \bar{a} \rangle\rangle_{\mathfrak{S}} = \text{dom}(p),$$

$$\begin{aligned} \text{and} \quad p[W(c)] &= \{p(f_\alpha(c)) \mid \alpha < \kappa, f_\alpha(c) \neq c\} \\ &= \{f_\alpha(p(c)) \mid \alpha < \kappa, f_\alpha(p(c)) \neq p(c)\} \\ &= W(p(c)). \end{aligned}$$

Hence, $\text{dom}(p)$ is W -closed and $p[W(c)] = W(p(c))$.

Finally, for $c \in \text{dom}(p)$, it follows by assumption on the predicates Q_α that

$$c\bar{d} \equiv p(c\bar{d}),$$

where \bar{d} is an enumeration of $W(c)$. Hence,

$$p[\text{tp}(c/W(c))] = \text{tp}(p(c)/W(p(c))). \quad \square$$

Using these two lemmas we can construct representations for stable theories.

Proposition 6.11. *Let T be a stable theory and set $\kappa := \min\{\text{st}(T), |T|\}$ and $\lambda := \min\{\text{fc}(\downarrow), |T|\}$.*

- (a) T has $\text{Wf}(\kappa, \lambda)$ -representations.
- (b) If $\text{fc}(\downarrow) \leq \aleph_0$, T has $\text{Lf}(\kappa, \aleph_0)$ -representations.

Proof. Let \mathfrak{M} be a model of T . By Theorem 5.8, there exists a $\sqrt[\lambda]{}$ -stratification $\zeta = (B_n)_{n < \gamma}$ of M over \emptyset of length $\gamma \leq \text{fc}(\downarrow)$. Since

$$\text{fc}(\downarrow) \leq \text{fc}(\downarrow) \oplus \text{mult}(\downarrow)^+ \leq \text{st}(T)^+ \quad \text{and} \quad \text{fc}(\downarrow) \leq |T|^+,$$

it follows that $\gamma \leq \kappa^+$. Taking a suitable refinement of ζ we may assume by Lemma 5.10 that $|B_0| = |B_1| = 1$. By Lemma 5.12, there exists a system of bases $(W(a))_{a \in M}$ with

$$|W(a)| < \text{loc}_0(\downarrow) \leq \text{fc}(\downarrow) \leq |T|^+.$$

Consequently, $|W(a)| \leq \lambda$.

We define $\mathfrak{S} := \langle M, (f_\alpha)_{\alpha < \lambda}, (P_\alpha)_{\alpha < \kappa}, (Q_\alpha)_{\alpha < \kappa} \rangle$ as follows. We choose the functions $(f_\alpha)_{\alpha < \lambda}$ such that, for every $a \in M$, $(f_\alpha(a))_{\alpha < \lambda}$ is an enumeration of $\{a\} \cup W(a)$. Fixing an injective function $h : 2^\kappa \rightarrow \wp(\kappa)$, we define

$$P_\alpha := \bigcup \{B_\beta \mid \alpha \in h(\beta)\}.$$

(Note that $\gamma \leq \kappa^+ \leq 2^\kappa$.) As there are at most $\text{st}(T)$ many types over the empty set and there are only $|T|$ formulae, we can fix an enumeration $(\varphi_\alpha(x))_{\alpha < \kappa}$ of all formulae (up to logical equivalence) of the form

$$\varphi(x; f_{\beta_0}x, \dots, f_{\beta_{n-1}}x),$$

where φ is a formula over the signature of T , $n < \omega$, and $\beta_0, \dots, \beta_{n-1} < \kappa$. We set

$$Q_\alpha := \{a \in M \mid \langle \mathfrak{M}, (f_\alpha)_\alpha \rangle \models \varphi_\alpha(a)\}.$$

It is straightforward to check that the structure \mathfrak{S} satisfies the conditions of Lemma 6.10. Hence, the identity function $\text{id} : M \rightarrow M$ is a representation of \mathfrak{M} in \mathfrak{S} .

Finally, note that $\mathfrak{S} \in \text{Wf}(\kappa, \lambda)$ since,

$$a \in B_\alpha \quad \text{implies} \quad f_\beta(a) \in \{a\} \cup B[<\alpha].$$

If $\text{fc}(\downarrow) \leq \aleph_0$, the sets $\{f_\alpha(a) \mid \alpha < \kappa\} = W(a) \cup \{a\}$ are finite, for all $a \in M$. By the Lemma of Kőnig, it follows that $\mathfrak{S} \in \text{Lf}(\kappa, \aleph_0)$. \square

The following two theorems summarise the results of this section.

Theorem 6.12 (Cohen, Shelah). *Let T be a complete first-order theory. The following conditions are equivalent:*

- (1) T is stable.
- (2) T has $\text{Un}(\kappa, \lambda)$ -representations, for some cardinals κ and λ .
- (3) T has $\text{Wf}(\mathfrak{o}, |T|)$ -representations.
- (4) T has $\text{Wf}(|T|, |T|)$ -representations.

Proof. (2) \Rightarrow (1) has been shown in Proposition 6.8 (a), the implications (4) \Rightarrow (3) \Rightarrow (2) follow from Lemmas 6.5 and 6.2, and (1) \Rightarrow (4) follows by Proposition 6.11. \square

Theorem 6.13 (Cohen, Shelah). *Let T be a complete first-order theory. The following conditions are equivalent:*

- (1) T is \aleph_0 -stable.
- (2) T has $\text{Lf}(\aleph_0, \aleph_0)$ -representations.

Proof. (2) \Rightarrow (1) follows by Proposition 6.8 (b) and (1) \Rightarrow (2) follows by Proposition 6.11. \square

Recommended Literature

Set theory

M. D. Potter, *Sets. An Introduction*, Oxford University Press 1990.

A. Lévy, *Basic Set Theory*, Springer 1979, Dover 2002.

K. Kunen, *Set Theory. An Introduction to Independence Proofs*, North-Holland 1983.

T. J. Jech, *Set Theory*, 3rd ed., Springer 2003.

Algebra and Category Theory

G. M. Bergman, *An Invitation to General Algebra and Universal Constructions*, 2nd ed., Springer 2015.

P. M. Cohn, *Universal Algebra*, 2nd ed., Springer 1981.

P. M. Cohn, *Basic Algebra*, Springer 2003.

S. Lang, *Algebra*, 3rd ed., Springer 2002.

F. Borceux, *Handbook of Categorical Algebra*, Cambridge University Press 1994.

S. MacLane, *Categories for the Working Mathematician*, 2nd ed., Springer 1998.

J. Adámek, J. Rosický, and M. Vitale, *Algebraic Theories*, Cambridge University Press 2011.

J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge University Press 1994.

Topology and lattice theory

R. Engelking, *General Topology*, 2nd ed., Heldermann 1989.
 C.-A. Faure, A. Frölicher, *Modern Projective Geometry*, Kluwer 2000.
 P. T. Johnstone, *Stone Spaces*, Cambridge University Press 1982.
 G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. Mislove, and D. S. Scott, *Continuous Lattices and Domains*, Cambridge University Press 2003.

Model theory

K. Tent and M. Ziegler, *A Course in Model Theory*, Cambridge University Press 2012.
 W. Hodges, *Model Theory*, Cambridge University Press 1993.
 B. Poizat, *A Course in Model Theory*, Springer 2000.
 C. C. Chang and H. J. Keisler, *Model Theory*, 3rd ed., North-Holland 1990.

General model theory

J. Barwise and S. Feferman, eds., *Model-Theoretic Logics*, Springer 1985.
 J. T. Baldwin, *Categoricity*, AMS 2010.
 R. Diaconescu, *Institution-Independent Model Theory*, Birkhäuser 2008.
 H.-D. Ebbinghaus and J. Flum, *Finite Model Theory*, Springer 1995.

Stability theory

S. Buechler, *Essential Stability Theory*, Springer 1996.
 E. Casanovas, *Simple Theories and Hyperimaginaries*, Cambridge University Press 2011.
 A. Pillay, *Geometric Stability Theory*, Oxford Science Publications 1996.
 F. O. Wagner, *Simple Theories*, Kluwer Academic Publishers 2000.
 S. Shelah, *Classification Theory*, 2nd ed., North-Holland 1990.

Symbol Index

Chapter A1

\mathbb{S}	universe of sets, 5
$a \in b$	membership, 5
$a \subseteq b$	subset, 5
HF	hereditary finite sets, 7
$\cap A$	intersection, 11
$A \cap B$	intersection, 11
$A \setminus B$	difference, 11
$\text{acc}(A)$	accumulation, 12
$\text{fnd}(A)$	founded part, 13
$\cup A$	union, 21
$A \cup B$	union, 21
$\wp(A)$	power set, 21
cut A	cut of A , 22

Chapter A2

$\langle a_0, \dots, a_{n-1} \rangle$	tuple, 27
$A \times B$	cartesian product, 27
$\text{dom } f$	domain of f , 28
$\text{rng } f$	range of f , 29
$f(a)$	image of a under f , 29
$f : A \rightarrow B$	function, 29
B^A	set of all functions $f : A \rightarrow B$, 29

id_A	identity function, 30
$S \circ R$	composition of relations, 30
$g \circ f$	composition of functions, 30
R^{-1}	inverse of R , 30
$R^{-1}(a)$	inverse image, 30
$R _C$	restriction, 30
$R \upharpoonright C$	left restriction, 31
$R[C]$	image of C , 31
$(a_i)_{i \in I}$	sequence, 37
$\prod_i A_i$	product, 37
pr_i	projection, 37
\bar{a}	sequence, 38
$\cup_i A_i$	disjoint union, 38
$A \sqcup B$	disjoint union, 38
in_i	insertion map, 39
\mathfrak{Q}^{op}	opposite order, 40
$\downarrow X$	initial segment, 41
$\uparrow X$	final segment, 41
$\downarrow X$	initial segment, 41
$\uparrow X$	final segment, 41
$[a, b]$	closed interval, 41
(a, b)	open interval, 41
$\max X$	greatest element, 42
$\min X$	minimal element, 42
$\sup X$	supremum, 42

$\inf X$	infimum, 42
$\mathfrak{A} \cong \mathfrak{B}$	isomorphism, 44
$\text{fix } f$	fixed points, 48
$\text{lfp } f$	least fixed point, 48
$\text{gfp } f$	greatest fixed point, 48
$[a]_{\sim}$	equivalence class, 54
A/\sim	set of \sim -classes, 54
$\text{TC}(R)$	transitive closure, 55

Chapter A3

a^+	successor, 59
$\text{ord}(\mathfrak{A})$	order type, 64
On	class of ordinals, 64
On_o	von Neumann ordinals, 69
$\rho(a)$	rank, 73
$A^{<\infty}$	functions $\downarrow \alpha \rightarrow A$, 74
$\mathfrak{A} + \mathfrak{B}$	sum, 85
$\mathfrak{A} \cdot \mathfrak{B}$	product, 86
$\mathfrak{A}^{(\mathfrak{B})}$	exponentiation of well-orders, 86
$\alpha + \beta$	ordinal addition, 89
$\alpha \cdot \beta$	ordinal multiplication, 89
$\alpha^{(\beta)}$	ordinal exponentiation, 89

Chapter A4

$ A $	cardinality, 113
∞	cardinality of proper classes, 113
Cn	class of cardinals, 113
\aleph_α	aleph alpha, 115
$\kappa \oplus \lambda$	cardinal addition, 116
$\kappa \otimes \lambda$	cardinal multiplication, 116

κ^λ	cardinal exponentiation, 116
$\sum_i \kappa_i$	cardinal sum, 122
$\prod_i \kappa_i$	cardinal product, 122
$\text{cf } \alpha$	cofinality, 124
\neg_α	beth alpha, 127
$(<\kappa)^\lambda$	$\sup_\mu \mu^\lambda$, 128
$\kappa^{<\lambda}$	$\sup_\mu \kappa^\mu$, 128

Chapter B1

$R^{\mathfrak{A}}$	relation of \mathfrak{A} , 149
$f^{\mathfrak{A}}$	function of \mathfrak{A} , 149
A^s	$A_{s_0} \times \dots \times A_{s_n}$, 151
$\mathfrak{A} \subseteq \mathfrak{B}$	substructure, 152
$\text{Sub}(\mathfrak{A})$	substructures of \mathfrak{A} , 152
$\mathfrak{S}ub(\mathfrak{A})$	substructure lattice, 152
$\mathfrak{A} _X$	induced substructure, 152
$\langle\langle X \rangle\rangle_{\mathfrak{A}}$	generated substructure, 154
$\mathfrak{A} _\Sigma$	reduct, 155
$\mathfrak{A} _T$	restriction to sorts in T , 155
$\mathfrak{A} \cong \mathfrak{B}$	isomorphism, 156
$\ker f$	kernel of f , 158
$h(\mathfrak{A})$	image of h , 162
\mathcal{C}^{obj}	class of objects, 162
$\mathcal{C}(a, b)$	morphisms $a \rightarrow b$, 162
$g \circ f$	composition of morphisms, 162
id_a	identity, 163
\mathcal{C}^{mor}	class of morphisms, 163
$\mathfrak{S}et$	category of sets, 163
$\mathfrak{H}om(\Sigma)$	category of homomorphisms, 163
$\mathfrak{H}om_s(\Sigma)$	category of strict homomorphisms, 163

$\mathfrak{Emb}(\Sigma)$	category of embeddings, 163
$\mathfrak{S}et_*$	category of pointed sets, 163
$\mathfrak{S}et^2$	category of pairs, 163
\mathcal{C}^{op}	opposite category, 166
F^{op}	opposite functor, 168
$(F \downarrow G)$	comma category, 170
$F \cong G$	natural isomorphism, 172
$\text{Cong}(\mathfrak{A})$	set of congruence relations, 176
$\mathfrak{C}ong(\mathfrak{A})$	congruence lattice, 176
\mathfrak{A}/\sim	quotient, 180

Chapter B2

$ x $	length of a sequence, 189
$x \cdot y$	concatenation, 189
\leq	prefix order, 189
\leq_{lex}	lexicographic order, 189
$ v $	level of a vertex, 192
$\text{frk}(v)$	foundation rank, 194
$a \sqcap b$	infimum, 197
$a \sqcup b$	supremum, 197
a^*	complement, 200
\mathfrak{L}^{op}	opposite lattice, 206
$\text{cl}_i(X)$	ideal generated by X , 206
$\text{cl}_f(X)$	filter generated by X , 206
\mathfrak{B}_2	two-element boolean algebra, 210
$\text{ht}(a)$	height of a , 218
$\text{rk}_p(a)$	partition rank, 222
$\text{deg}_p(a)$	partition degree, 226

Chapter B3

$T[\Sigma, X]$	finite Σ -terms, 231
t_v	subterm at v , 232
$\text{free}(t)$	free variables, 235
$t^{\mathfrak{A}}[\beta]$	value of t , 235
$\mathfrak{T}[\Sigma, X]$	term algebra, 236
$t[x/s]$	substitution, 238
$\mathfrak{S}ig\mathfrak{Var}$	category of signatures and variables, 239
$\mathfrak{S}ig$	category of signatures, 240
\mathfrak{Var}	category of variables, 240
$\mathfrak{T}erm$	category of terms, 240
$\mathfrak{A} _\mu$	μ -reduct of \mathfrak{A} , 241
$\text{Str}[\Sigma]$	class of Σ -structures, 241
$\text{Str}[\Sigma, X]$	class of all Σ -structures with variable assignments, 241
$\mathfrak{S}tr\mathfrak{Var}$	category of structures and assignments, 241
$\mathfrak{S}tr$	category of structures, 241
$\prod_i \mathfrak{A}^i$	direct product, 243
$\llbracket \varphi \rrbracket$	set of indices, 245
$\bar{a} \sim_u \bar{b}$	filter equivalence, 245
$u _J$	restriction of u to J , 246
$\prod_i \mathfrak{A}^i / u$	reduced product, 246
\mathfrak{A}^u	ultrapower, 247
$\varinjlim D$	directed colimit, 255
$\varinjlim D$	colimit of D , 257
$\varprojlim D$	directed limit, 260
$f * \mu$	componentwise composition for cocones, 262
$G[\mu]$	image of a cocone under a functor, 265
\mathfrak{Z}_n	partial order of an alternating path, 276

\mathfrak{Z}_n^\perp	partial order of an alternating path, 276
$f \rightsquigarrow g$	alternating-path equivalence, 277
$[f]_F^\rightsquigarrow$	alternating-path equivalence class, 277
$s * t$	componentwise composition of links, 280
π_t	projection along a link, 281
in_D	inclusion link, 281
$D[t]$	image of a link under a functor, 284
$\text{Ind}_{\mathcal{P}}(\mathcal{C})$	inductive \mathcal{P} -completion, 285
$\text{Ind}_{\text{all}}(\mathcal{C})$	inductive completion, 285

Chapter B4

$\text{Ind}_\kappa^{\lambda}(\mathcal{C})$	inductive (κ, λ) -completion, 295
$\text{Ind}(\mathcal{C})$	inductive completion, 296
\mathcal{C}	loop category, 317
$\ \mathfrak{a}\ $	cardinality in an accessible category, 333
$\mathfrak{Sub}_{\mathcal{K}}(\mathfrak{a})$	category of \mathcal{K} -subobjects, 341
$\mathfrak{Sub}_\kappa(\mathfrak{a})$	category of κ -presentable subobjects, 341

Chapter B5

$\text{cl}(A)$	closure of A , 347
$\text{int}(A)$	interior of A , 347

∂A	boundary of A , 347
$\text{rk}_{\text{CB}}(x/A)$	Cantor-Bendixson rank, 369
$\text{spec}(\mathfrak{L})$	spectrum of \mathfrak{L} , 374
$\langle x \rangle$	basic closed set, 374
$\text{clop}(\mathfrak{S})$	algebra of clopen subsets, 378

Chapter B6

$\mathfrak{Aut} \mathfrak{M}$	automorphism group, 390
G/U	set of cosets, 390
$\mathfrak{S}/\mathfrak{N}$	factor group, 392
$\mathfrak{Sym} \Omega$	symmetric group, 393
$g a$	action of g on a , 394
$G \bar{a}$	orbit of \bar{a} , 394
$\mathfrak{S}_{(X)}$	pointwise stabiliser, 395
$\mathfrak{S}_{\{X\}}$	setwise stabiliser, 395
$\langle \bar{a} \mapsto \bar{b} \rangle$	basic open set of the group topology, 399
$\deg p$	degree, 403
$\mathfrak{I}(\mathfrak{R})$	lattice of ideals, 404
$\mathfrak{R}/\mathfrak{a}$	quotient of a ring, 406
$\text{Ker } h$	kernel, 406
$\text{spec}(\mathfrak{R})$	spectrum, 406
$\bigoplus_i \mathfrak{M}_i$	direct sum, 409
$\mathfrak{M}^{(I)}$	direct power, 409
$\dim \mathfrak{Q}$	dimension, 413
$\text{FF}(\mathfrak{R})$	field of fractions, 415
$\mathfrak{R}(\bar{a})$	subfield generated by \bar{a} , 418
$p[x]$	polynomial function, 419
$\text{Aut}(\mathfrak{L}/\mathfrak{R})$	automorphisms over K , 427
$ a $	absolute value, 430

Chapter C1

$\text{ZL}[\mathfrak{R}, X]$	Zariski logic, 447
\models	satisfaction relation, 448
$\text{BL}(\mathfrak{B})$	boolean logic, 448
$\text{FO}_{\kappa\aleph_0}[\Sigma, X]$	infinitary first-order logic, 449
$\neg\varphi$	negation, 449
$\wedge \Phi$	conjunction, 449
$\vee \Phi$	disjunction, 449
$\exists x\varphi$	existential quantifier, 449
$\forall x\varphi$	universal quantifier, 449
$\text{FO}[\Sigma, X]$	first-order logic, 449
$\mathfrak{A} \models \varphi[\beta]$	satisfaction, 450
true	true, 451
false	false, 451
$\varphi \vee \psi$	disjunction, 451
$\varphi \wedge \psi$	conjunction, 451
$\varphi \rightarrow \psi$	implication, 451
$\varphi \leftrightarrow \psi$	equivalence, 451
$\text{free}(\varphi)$	free variables, 454
$\text{qr}(\varphi)$	quantifier rank, 457
$\text{Mod}_L(\Phi)$	class of models, 458
$\Phi \models \varphi$	entailment, 464
\equiv	logical equivalence, 464
$\Phi^=$	closure under entailment, 464
$\text{Th}_L(\mathfrak{I})$	L -theory, 465
\equiv_L	L -equivalence, 466
$\text{DNF}(\varphi)$	disjunctive normal form, 471
$\text{CNF}(\varphi)$	conjunctive normal form, 471
$\text{NNF}(\varphi)$	negation normal form, 473
\mathfrak{Logic}	category of logics, 482
$\exists^{\lambda} x\varphi$	cardinality quantifier, 485

$\text{FO}_{\kappa\aleph_0}(\text{wo})$	FO with well-ordering quantifier, 486
W	well-ordering quantifier, 486
$Q_{\mathcal{K}}$	Lindström quantifier, 486
$\text{SO}_{\kappa\aleph_0}[\Sigma, \Xi]$	second-order logic, 487
$\text{MSO}_{\kappa\aleph_0}[\Sigma, \Xi]$	monadic second-order logic, 487
\mathfrak{PO}	category of partial orders, 492
\mathfrak{Lb}	Lindenbaum functor, 492
$\neg\varphi$	negation, 494
$\varphi \vee \psi$	disjunction, 494
$\varphi \wedge \psi$	conjunction, 494
$L _{\Phi}$	restriction to Φ , 495
L/Φ	localisation to Φ , 495
\models_{Φ}	consequence modulo Φ , 495
\equiv_{Φ}	equivalence modulo Φ , 495

Chapter C2

$\mathfrak{Emb}_L(\Sigma)$	category of L -embeddings, 497
$\text{QF}_{\kappa\aleph_0}[\Sigma, X]$	quantifier-free formulae, 498
$\exists\Delta$	existential closure of Δ , 498
$\forall\Delta$	universal closure of Δ , 498
$\exists_{\kappa\aleph_0}$	existential formulae, 498
$\forall_{\kappa\aleph_0}$	universal formulae, 498
$\exists_{\kappa\aleph_0}^+$	positive existential formulae, 498
\leq_{Δ}	Δ -extension, 502
\leq	elementary extension, 502

Φ_{Δ}^{\pm} Δ -consequences of Φ , 525
 \leq_{Δ} preservation of
 Δ -formulae, 525

Chapter c3

$S(L)$ set of types, 531
 $\langle \Phi \rangle$ types containing Φ , 531
 $\text{tp}_L(\bar{a}/\mathfrak{M})$ L -type of \bar{a} , 532
 $S_L^s(T)$ type space for a theory, 532
 $S_L^s(U)$ type space over U , 532
 $\mathfrak{C}(L)$ type space, 537
 $f(p)$ conjugate of p , 547
 $\mathfrak{C}_{\Delta}(L)$ $\mathfrak{C}(L|_{\Delta})$ with topology
induced from $\mathfrak{C}(L)$, 561
 $\langle \Phi \rangle_{\Delta}$ closed set in $\mathfrak{C}_{\Delta}(L)$, 561
 $p|_{\Delta}$ restriction to Δ , 564
 $\text{tp}_{\Delta}(\bar{a}/U)$ Δ -type of \bar{a} , 564

Chapter c4

\equiv_{α} α -equivalence, 581
 \equiv_{∞} ∞ -equivalence, 581
 $\text{pIso}_{\kappa}(\mathfrak{A}, \mathfrak{B})$ partial isomorphisms,
582
 $\bar{a} \mapsto \bar{b}$ map $a_i \mapsto b_i$, 582
 \emptyset the empty function, 582
 $I_{\alpha}(\mathfrak{A}, \mathfrak{B})$ back-and-forth system, 583
 $I_{\infty}(\mathfrak{A}, \mathfrak{B})$ limit of the system, 585
 \cong_{α} α -isomorphic, 585
 \cong_{∞} ∞ -isomorphic, 585
 $m =_k n$ equality up to k , 587
 $\phi_{\mathfrak{A}, \bar{a}}^{\alpha}$ Hintikka formula, 590

$\text{EF}_{\alpha}(\mathfrak{A}, \bar{a}, \mathfrak{B}, \bar{b})$
Ehrenfeucht-Fraïssé
game, 593
 $\text{EF}_{\infty}^{\kappa}(\mathfrak{A}, \bar{a}, \mathfrak{B}, \bar{b})$
Ehrenfeucht-Fraïssé
game, 593
 $I_{\text{FO}}^{\kappa}(\mathfrak{A}, \mathfrak{B})$ partial FO-maps of size κ ,
602
 $\sqsubseteq_{\text{iso}}^{\kappa}$ $\infty\kappa$ -simulation, 603
 $\cong_{\text{iso}}^{\kappa}$ $\infty\kappa$ -isomorphic, 603
 $\mathfrak{A} \sqsubseteq_{\circ}^{\kappa} \mathfrak{B}$ $I_{\circ}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \sqsubseteq_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathfrak{A} \equiv_{\circ}^{\kappa} \mathfrak{B}$ $I_{\circ}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \equiv_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathfrak{A} \sqsubseteq_{\text{FO}}^{\kappa} \mathfrak{B}$ $I_{\text{FO}}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \sqsubseteq_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathfrak{A} \equiv_{\text{FO}}^{\kappa} \mathfrak{B}$ $I_{\text{FO}}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \equiv_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathfrak{A} \sqsubseteq_{\infty}^{\kappa} \mathfrak{B}$ $I_{\infty}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \sqsubseteq_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathfrak{A} \equiv_{\infty}^{\kappa} \mathfrak{B}$ $I_{\infty}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \equiv_{\text{iso}}^{\kappa} \mathfrak{B}$, 603
 $\mathcal{G}(\mathfrak{A})$ Gaifman graph, 609

Chapter c5

$L \leq L'$ L' is as expressive as L , 617
(A) algebraic, 618
(B) boolean closed, 618
(B₊) positive boolean closed,
618
(C) compactness, 618
(CC) countable compactness,
618
(FOP) finite occurrence property,
618
(KP) Karp property, 618
(LSP) Löwenheim-Skolem
property, 618
(REL) closed under
relativisations, 618

(SUB) closed under substitutions,
618
(TUP) Tarski union property, 618
 $\text{hn}_{\kappa}(L)$ Hanf number, 622
 $\text{ln}_{\kappa}(L)$ Löwenheim number, 622
 $\text{wn}_{\kappa}(L)$ well-ordering number, 622
 $\text{occ}(L)$ occurrence number, 622
 $\text{pr}_T(\mathcal{K})$ T -projection, 640
 $\text{PC}_{\kappa}(L, \Sigma)$ projective L -classes, 641
 $L_0 \leq_{\text{PC}}^{\kappa} L_1$ projective reduction, 641
 $\text{RPC}_{\kappa}(L, \Sigma)$ relativised projective
 L -classes, 645
 $L_0 \leq_{\text{RPC}}^{\kappa} L_1$ relativised projective
reduction, 645
 $\Delta(L)$ interpolation closure, 653
 $\text{ifp } f$ inductive fixed point, 662
 $\text{lim inf } f$ least partial fixed point,
662
 $\text{lim sup } f$ greatest partial fixed point,
663
 f_{φ} function defined by φ , 668
 $\text{FO}_{\kappa\aleph_0}(\text{LFP})$ least fixed-point logic,
668
 $\text{FO}_{\kappa\aleph_0}(\text{IFP})$ inflationary fixed-point
logic, 669
 $\text{FO}_{\kappa\aleph_0}(\text{PFP})$ partial fixed-point
logic, 669
 \triangleleft_{φ} stage comparison, 679

Chapter d1

$\text{tor}(\mathfrak{B})$ torsion subgroup, 709
 a/n divisor, 710
DAG theory of divisible
torsion-free abelian

groups, 710
ODAG theory of ordered divisible
abelian groups, 710
 $\text{div}(\mathfrak{B})$ divisible closure, 711
 F field axioms, 714
ACF theory of algebraically
closed fields, 714
RCF theory of real closed fields,
715

Chapter d2

$\langle <\mu \rangle^{\lambda}$ $\bigcup_{\kappa < \mu} \kappa^{\lambda}$, 727
 $\text{HO}_{\infty}[\Sigma, X]$ infinitary Horn
formulae, 740
 $\text{SH}_{\infty}[\Sigma, X]$ infinitary strict Horn
formulae, 740
 $\text{H}\forall_{\infty}[\Sigma, X]$ infinitary universal
Horn formulae, 740
 $\text{SH}\forall_{\infty}[\Sigma, X]$ infinitary universal
strict Horn formulae,
740
 $\text{HO}[\Sigma, X]$ first-order Horn formulae,
740
 $\text{SH}[\Sigma, X]$ first-order strict Horn
formulae, 740
 $\text{H}\forall[\Sigma, X]$ first-order universal Horn
formulae, 740
 $\text{SH}\forall[\Sigma, X]$ first-order universal
strict Horn formulae,
740
 $\langle C; \Phi \rangle$ presentation, 745
 $\text{Prod}(\mathcal{K})$ products, 749
 $\text{Sub}(\mathcal{K})$ substructures, 749
 $\text{Iso}(\mathcal{K})$ isomorphic copies, 749

$\text{Hom}(\mathcal{K})$ weak homomorphic images, 749
 $\text{ERP}(\mathcal{K})$ embeddings into reduced products, 749
 $\text{QV}(\mathcal{K})$ quasivariety, 749
 $\text{Var}(\mathcal{K})$ variety, 749

Chapter D3

(f, g) open cell between f and g , 761
 $[f, g]$ closed cell between f and g , 761
 $B(\bar{a}, \bar{b})$ box, 762
 $\text{Cn}(D)$ continuous functions, 776
 $\dim C$ dimension, 777

Chapter E2

$\text{dcl}_L(U)$ L -definitional closure, 819
 $\text{acl}_L(U)$ L -algebraic closure, 819
 $\text{dcl}_{\text{Aut}}(U)$ Aut-definitional closure, 821
 $\text{acl}_{\text{Aut}}(U)$ Aut-algebraic closure, 821
 \mathbb{M} the monster model, 829
 $A \equiv_U B$ having the same type over U , 830
 \mathfrak{M}^{eq} extension by imaginary elements, 831
 $\text{dcl}^{\text{eq}}(U)$ definable closure in \mathfrak{M}^{eq} , 831
 $\text{acl}^{\text{eq}}(U)$ algebraic closure in \mathfrak{M}^{eq} , 831
 T^{eq} theory of \mathbb{M}^{eq} , 833

$\text{Gb}(\mathfrak{p})$ Galois base, 841

Chapter E3

$I_{\text{cl}}(\mathfrak{A}, \mathfrak{B})$ elementary maps with closed domain and range, 877

Chapter E4

$\text{pMor}_{\mathcal{K}}(a, b)$ category of partial morphisms, 898
 $a \sqsubseteq_{\mathcal{K}} b$ forth property for objects in \mathcal{K} , 899
 $a \sqsubseteq_{\text{pres}}^{\kappa} b$ forth property for κ -presentable objects, 899
 $a \equiv_{\text{pres}}^{\kappa} b$ back-and-forth equivalence for κ -presentable objects, 899
 $\text{Sub}_{\kappa}(a)$ κ -presentable subobjects, 910
 $\text{atp}(\bar{a})$ atomic type, 922
 η^{eq} extension axiom, 922
 $T[\mathcal{K}]$ extension axioms for \mathcal{K} , 922
 $T_{\text{ran}}[\Sigma]$ random theory, 922
 $\kappa_n(\varphi)$ number of models, 924
 $\text{Pr}_{\mathfrak{M}}^{\#}[\mathfrak{M} \models \varphi]$ density of models, 924

Chapter E5

$[I]^{\kappa}$ increasing κ -tuples, 929
 $\kappa \rightarrow (\mu)_{\lambda}^{\nu}$ partition theorem, 929
 $\text{pf}(\eta, \zeta)$ prefix of ζ of length $|\eta|$, 934
 $\mathfrak{T}_*(\kappa^{<\alpha})$ index tree with small signature, 934
 $\mathfrak{T}_n(\kappa^{<\alpha})$ index tree with large signature, 934
 $\langle\langle X \rangle\rangle_n$ substructure generated in $\mathfrak{T}_n(\kappa^{<\alpha})$, 934
 $\text{Lvl}(\bar{\eta})$ levels of $\bar{\eta}$, 935
 \approx_* equal atomic types in \mathfrak{T}_* , 936
 \approx_n equal atomic types in \mathfrak{T}_n , 936
 $\approx_{n,k}$ refinement of \approx_n , 936
 $\approx_{\omega,k}$ union of $\approx_{n,k}$, 936
 $\bar{a}[\bar{i}]$ $\bar{a}^{i_0} \dots \bar{a}^{i_{n-1}}$, 945
 $\text{tp}_{\Delta}(\bar{a}/U)$ Δ -type, 945
 $\text{Av}((\bar{a}^i)_i/U)$ average type, 947
 $\llbracket \varphi(\bar{a}^i) \rrbracket$ indices satisfying φ , 956
 $\text{Av}_1((\bar{a}^i)_i/C)$ unary average type, 966

Chapter E6

$\mathfrak{Emb}(\mathcal{K})$ embeddings between structures in \mathcal{K} , 969
 p^F image of a partial isomorphism under F , 972
 $\text{Th}_L(F)$ theory of a functor, 975
 \mathfrak{Q}^{α} inverse reduct, 979
 $\mathcal{R}(\mathfrak{M})$ relational variant of \mathfrak{M} , 981

$\text{Av}(F)$ average type, 990

Chapter E7

$\text{ln}(\mathcal{K})$ Löwenheim number, 999
 $\mathfrak{A} \leq_{\mathcal{K}} \mathfrak{B}$ \mathcal{K} -substructure, 1000
 $\text{hn}(\mathcal{K})$ Hanf number, 1007
 \mathcal{K}_{κ} structures of size κ , 1008
 $I_{\mathcal{K}}^{\kappa}(\mathfrak{A}, \mathfrak{B})$ \mathcal{K} -embeddings, 1012
 $\mathfrak{A} \sqsubseteq_{\mathcal{K}}^{\kappa} \mathfrak{B}$ $I_{\mathcal{K}}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \sqsubseteq_{\text{iso}}^{\kappa} \mathfrak{B}$, 1012
 $\mathfrak{A} \equiv_{\mathcal{K}}^{\kappa} \mathfrak{B}$ $I_{\mathcal{K}}^{\kappa}(\mathfrak{A}, \mathfrak{B}) : \mathfrak{A} \equiv_{\text{iso}}^{\kappa} \mathfrak{B}$, 1012

Chapter F1

$\langle\langle X \rangle\rangle_D$ span of X , 1035
 $\text{dim}_{\text{cl}}(X)$ dimension, 1041
 $\text{dim}_{\text{cl}}(X/U)$ dimension over U , 1041

Chapter F2

$\text{rk}_{\Delta}(\varphi)$ Δ -rank, 1077
 $\text{rk}_{\mathbb{M}}^{\bar{s}}(\varphi)$ Morley rank, 1077
 $\text{deg}_{\mathbb{M}}^{\bar{s}}(\varphi)$ Morley degree of φ , 1080
(MON) Monotonicity, 1088
(NOR) Normality, 1088
(LRF) Left Reflexivity, 1088
(LTR) Left Transitivity, 1088
(FIN) Finite Character, 1089
(SYM) Symmetry, 1089
(BMON) Base Monotonicity, 1089
(SRB) Strong Right Boundedness, 1089

$\text{cl}_\sqrt{}$	closure operation associated with $\sqrt{}$, 1095
(INV)	Invariance, 1101
(DEF)	Definability, 1101
(EXT)	Extension, 1102
$A \stackrel{\text{df}}{\sqrt{U}} B$	definable over, 1103
$A \stackrel{\text{at}}{\sqrt{U}} B$	isolated over, 1103
$A \stackrel{s}{\sqrt{U}} B$	non-splitting over, 1103
$\mathfrak{p} \leq \sqrt{\mathfrak{q}}$	$\sqrt{}$ -free extension, 1108
$A \stackrel{\text{u}}{\sqrt{U}} B$	finitely satisfiable, 1109
$\text{Av}(u/B)$	average type of u , 1109
(LLOC)	Left Locality, 1114
(RLOC)	Right Locality, 1114
$\text{loc}(\sqrt{})$	right locality cardinal of $\sqrt{}$, 1114
$\text{loc}_o(\sqrt{})$	finitary right locality cardinal of $\sqrt{}$, 1114
κ^{reg}	regular cardinal above κ , 1114
$\text{fc}(\sqrt{})$	length of $\sqrt{}$ -forking chains, 1115
(SFIN)	Strong Finite Character, 1116
$\sqrt{}^*$	forking relation to $\sqrt{}$, 1118

Chapter F3

$A \stackrel{\text{d}}{\sqrt{U}} B$	non-dividing, 1131
$A \stackrel{\text{f}}{\sqrt{U}} B$	non-forking, 1131
$A \stackrel{\text{i}}{\sqrt{U}} B$	globally invariant over, 1140

Chapter F4

$\text{alt}_\varphi(\bar{a}_i)_{i \in I}$	φ -alternation number, 1159
$\text{rk}_{\text{alt}}(\varphi)$	alternation rank, 1159
$\text{in}(\sim)$	intersection number, 1170
$\bar{a} \approx_U^{\text{ls}} \bar{b}$	indiscernible sequence starting with \bar{a}, \bar{b}, \dots , 1174
$\bar{a} \equiv_U^{\text{ls}} \bar{b}$	Lascar strong type equivalence, 1174
$\text{CF}((\bar{a}_i)_{i \in I})$	cofinal type, 1200
$\text{Ev}((\bar{a}_i)_{i \in I})$	eventual type, 1205
$\text{rk}_{\text{dp}}(\bar{a}/U)$	dp-rank, 1218

Chapter F5

(LEXT)	Left Extension, 1234
$A \stackrel{\text{fi}}{\sqrt{U}} B$	combination of $\stackrel{\text{li}}{\sqrt{}}$ and $\stackrel{\text{f}}{\sqrt{}}$, 1245
$A \stackrel{\text{sli}}{\sqrt{U}} B$	strict Lascar invariance, 1245
(WIND)	Weak Independence Theorem, 1258
(IND)	Independence Theorem, 1259

Chapter G1

$\bar{a} \downarrow_U^{\text{i}} B$	unique free extension, 1280
$\text{mult}_{\sqrt{}}(\mathfrak{p})$	$\sqrt{}$ -multiplicity of \mathfrak{p} , 1285
$\text{mult}(\sqrt{})$	multiplicity of $\sqrt{}$, 1285
$\text{st}(T)$	minimal cardinal T is stable in, 1296

Chapter G2

(RSH)	Right Shift, 1303
$\text{lbm}(\sqrt{})$	left base-monotonicity cardinal, 1303
$A[I]$	$\bigcup_{i \in I} A_i$, 1312
$A[<\alpha]$	$\bigcup_{i < \alpha} A_i$, 1312
$A[\leq\alpha]$	$\bigcup_{i \leq \alpha} A_i$, 1312

$A \perp_U^{\text{do}} B$	definable orthogonality, 1335
$A \stackrel{\text{si}}{\sqrt{U}} B$	strong independence, 1338
$\Upsilon_{\kappa\lambda}$	unary signature, 1344
$\text{Un}(\kappa, \lambda)$	class of unary structures, 1344
$\text{Lf}(\kappa, \lambda)$	class of locally finite unary structures, 1344

Index

- abelian group, 389
- abstract elementary class, 999
- abstract independence relation, 1088
- κ -accessible category, 333
- accumulation, 12
- accumulation point, 368
- action, 394
- acyclic, 523
- addition of cardinals, 116
- addition of ordinals, 89
- adjoint functors, 238
- affine geometry, 1041
- aleph, 115
- algebraic, 149, 819
- algebraic class, 1000
- algebraic closure, 819
- algebraic closure operator, 51
- algebraic diagram, 503
- algebraic elements, 422
- algebraic field extensions, 422
- algebraic logic, 491
- algebraic prime model, 698
- algebraically closed, 819
- algebraically closed field, 422, 714
- algebraically independent, 422
- almost strongly minimal theory, 1060
- alternating path in a category, 276
- alternating-path equivalence, 277
- φ -alternation number, 1159
- alternation rank of a formula, 1159
- amalgamation class, 1009
- amalgamation property, 914, 1008
- amalgamation square, 657
- Amalgamation Theorem, 525
- antisymmetric, 40
- arity, 28, 29, 149
- array, 1227
- array property, 1227
- array-dividing, 1233
- associative, 31
- asynchronous product, 757
- atom, 449
- atom of a lattice, 218
- atomic, 218
- atomic diagram, 503
- atomic structure, 859
- atomic type, 922
- atomless, 218
- automorphism, 156
- automorphism group, 390
- average type, 947
- average type of an Ehrenfeucht-Mostowski functor, 990

- average type of an indiscernible system, 954
 average type of an ultrafilter, 1109
 Axiom of Choice, 109, 462
 Axiom of Creation, 19, 462
 Axiom of Extensionality, 5, 462
 Axiom of Infinity, 24, 462
 Axiom of Replacement, 133, 462
 Axiom of Separation, 10, 462
 axiom system, 458
 axiomatisable, 458
 axiomatise, 458
- back-and-forth property, 582, 897
 back-and-forth system, 582
 Baire, property of —, 367
 ball, 346
 $\sqrt{\quad}$ -base, 1234
 base monotonicity, 1089
 base of a partial morphism, 898
 base projection, 898
 base, closed —, 348
 base, open —, 348
 bases for a stratification, 1342
 basic Horn formula, 740
 basis, 110, 1038, 1041
 beth, 127
 Beth property, 652, 826
 bidefinable, 889
 biindiscernible family, 1225
 biinterpretable, 895
 bijective, 31
 boolean algebra, 200, 459, 494
 boolean closed, 494
 boolean lattice, 200
 boolean logic, 448, 466
 bound variable, 454
- boundary, 347, 762
 κ -bounded, 602
 bounded equivalence relation, 1179
 bounded lattice, 197
 bounded linear order, 587
 bounded logic, 622
 box, 762
 branch, 191
 branching degree, 193
- canonical base, 838
 canonical definition, 835
 weak —, 851
 canonical diagram, 341
 canonical parameter, 835
 weak —, 851
 canonical projection from the \mathcal{P} -completion, 313
 Cantor discontinuum, 355, 538
 Cantor normal form, 100
 Cantor-Bendixson rank, 369, 381
 cardinal, 113
 cardinal addition, 116
 cardinal exponentiation, 116, 127
 cardinal multiplication, 116
 cardinality, 113, 333
 cardinality quantifier, 486
 cartesian product, 27
 categorical, 881, 913
 category, 162
 δ -cell, 777
 cell decomposition, 779
 Cell Decomposition Theorem, 780
 chain, 42
 L -chain, 505
 chain condition, 1253
- chain condition for Morley sequences, 1263
 chain in a category, 272
 chain topology, 354
 chain-bounded formula, 1174
 Chang's reduction, 536
 character, 105
 characteristic, 715
 characteristic of a field, 417
 choice function, 106
 Choice, Axiom of —, 109, 462
 class, 9, 54
 clopen set, 345
 =-closed, 516
 closed base, 348
 closed function, 350
 closed interval, 761
 closed set, 51, 53, 345
 closed subbase, 348
 closed subset of a construction, 875, 1313
 closed unbounded set, 135
 closed under relativisations, 618
 closed under substitutions, 618
 closure operator, 51, 110
 closure ordinal, 81
 closure space, 53
 closure under reverse ultrapowers, 739
 closure, topological —, 347
 co-chain-bounded relation, 1179
 cocone, 257
 cocone functor, 262
 codomain of a partial morphism, 898
 codomain projection, 898
 coefficient, 402
 cofinal, 124
- cofinality, 124
 Coincidence Lemma, 235
 colimit, 257
 comma category, 170
 commutative, 389
 commutative ring, 401
 commuting diagram, 164
 comorphism of logics, 482
 compact, 356, 617
 compact, countably —, 617
 Compactness Theorem, 519, 535
 compactness theorem, 724
 compatible, 477
 complement, 200
 complete, 466
 κ -complete, 602
 complete partial order, 43, 50, 53
 complete type, 531
 completion of a diagram, 311
 (λ, κ) -completion of a diagram, 311
 (λ, κ) -completion of a partial order, 305
 composition, 30
 composition of links, 280
 concatenation, 189
 condition of filters, 727
 cone, 261
 confluence property, 1204
 confluent family of sequences, 1204
 congruence relation, 176
 conjugacy class, 395
 conjugate, 821
 conjugation, 395
 conjunction, 449, 494
 conjunctive normal form, 471
 connected category, 276
 connected, definably —, 765

- consequence, 464, 492, 525
 consistence of filters with conditions,
 727
 consistency over a family, 1227
 consistent, 458
 constant, 29, 149
 constructible set, 873
 $\sqrt{\quad}$ -constructible set, 1312
 construction, 873
 $\sqrt{\quad}$ -construction, 1312
 continuous, 46, 134, 350
 contradictory formulae, 632
 contravariant, 168
 convex equivalence relation, 1170
 coset, 390
 countable, 110, 115
 countably compact, 617
 covariant, 167
 cover, 356
 Creation, Axiom of —, 19, 462
 cumulative hierarchy, 18
 cut, 22

 deciding a condition, 727
 definability of independence
 relations, 1101
 definable, 819
 definable expansion, 477
 definable orthogonality, 1335
 definable Skolem function, 847
 definable structure, 889
 definable type, 574, 1103
 definable with parameters, 763
 definably connected, 765
 defining a set, 451
 definition of a type, 574
 definitional closed, 819
 definitional closure, 819
 degree of a polynomial, 403
 dense class, 1261
 dense linear order, 604
 κ -dense linear order, 604
 dense order, 459
 dense set, 365
 dense sets in directed orders, 251
 dense subcategory, 286
 dependence relation, 1035
 dependent, 1035
 dependent set, 110
 derivation, 402
 diagonal functor, 258
 diagonal intersection, 137
 diagram, 255, 260
 L -diagram, 503
 Diagram Lemma, 503, 638
 difference, 11
 dimension, 1041
 dimension function, 1042
 dimension of a cell, 777
 dimension of a vector space, 413
 direct limit, 256
 direct power, 409
 direct product, 243
 direct sum of modules, 409
 directed, 250
 directed colimit, 255
 directed diagram, 255
 κ -directed diagram, 255
 directed limit, 260
 discontinuum, 355
 discrete linear order, 587
 discrete topology, 346
 disintegrated matroid, 1048
 disjoint union, 38

 disjunction, 449, 494
 disjunctive normal form, 471
 distributive, 200
 dividing, 1131
 dividing chain, 1143
 dividing κ -tree, 1150
 divisible closure, 711
 divisible group, 710
 domain, 28, 151
 domain of a partial morphism, 898
 domain projection, 898
 dp-rank, 1218
 dual categories, 172

 Ehrenfeucht-Fraïssé game, 593, 596
 Ehrenfeucht-Mostowski functor, 990,
 1006
 Ehrenfeucht-Mostowski model, 990
 element of a set, 5
 elementary diagram, 503
 elementary embedding, 497, 502
 elementary extension, 502
 elementary map, 497
 elementary substructure, 502
 elimination
 uniform — of imaginaries, 844
 elimination of finite imaginaries, 857
 elimination of imaginaries, 845
 elimination set, 694
 embedding, 44, 156, 498
 Δ -embedding, 497
 \mathcal{K} -embedding, 999
 elementary —, 497
 embedding of a tree into a lattice, 224
 embedding of logics, 482
 embedding of permutation groups,
 890

 embedding, elementary —, 502
 endomorphism ring, 408
 entailment, 464, 492
 epimorphism, 165
 equivalence class, 54
 equivalence formula, 830
 equivalence of categories, 172
 equivalence relation, 54, 459
 L -equivalent, 466
 α -equivalent, 581, 596
 equivalent categories, 172
 equivalent formulae, 464
 Erdős-Rado theorem, 932
 Euklidean norm, 345
 even, 927
 exchange property, 110
 existential, 498
 existential closure, 703
 existential quantifier, 449
 existentially closed, 703
 expansion, 155, 1002
 expansion, definable —, 477
 explicit definition, 652
 exponentiation of cardinals, 116, 127
 exponentiation of ordinals, 89
 extension, 152, 1102
 Δ -extension, 502
 extension axiom, 922
 $\sqrt{\quad}$ -extension base, 1234
 extension of fields, 418
 extension, elementary —, 502
 Extensionality, Axiom of —, 5, 462

 factorisation, 180
 Factorisation Lemma, 158
 factorising through a cocone, 321
 faithful functor, 167

family, 37
 field, 401, 461, 502, 714
 field extension, 418
 field of a relation, 29
 field of fractions, 415
 field, real —, 430
 field, real closed —, 433
 filter, 205, 209, 534
 κ -filtered category, 289
 κ -filtered colimit, 289
 κ -filtered diagram, 289
 final segment, 41
 κ -finitary set of partial isomorphisms, 602
 finite, 115
 finite character, 51, 105, 1089
 strong —, 1116
 finite equivalence relation, 1170
 finite intersection property, 213
 finite occurrence property, 617
 finite, being — over a set, 780
 finitely axiomatisable, 458
 finitely branching, 193
 finitely generated, 154
 finitely presentable, 321
 finitely satisfiable type, 1109
 first-order interpretation, 450, 479
 first-order logic, 449
 fixed point, 48, 81, 134, 661
 fixed-point induction, 77
 fixed-point rank, 679
 Fodor
 Theorem of —, 139
 follow, 464
 forcing, 727
 forgetful functor, 168, 238
 forking chain, 1143

$\sqrt{\quad}$ -forking chain, 1115
 $\sqrt{\quad}$ -forking formula, 1108
 forking relation, 1101
 $\sqrt{\quad}$ -forking type, 1108
 formal power series, 402
 formula, 448
 forth property for partial morphisms, 899
 foundation rank, 194
 founded, 13
 Fraïssé limit, 916
 free algebra, 236
 free extension of a type, 1108
 $\sqrt{\quad}$ -free extension of a type, 1108
 free model, 745
 free structures, 754
 $\sqrt{\quad}$ -free type, 1108
 free variables, 235, 454
 full functor, 167
 full subcategory, 169
 function, 29
 functional, 29, 149
 functor, 167

 Gaifman graph, 609
 Gaifman, Theorem of —, 615
 Galois base, 838
 Galois saturated structure, 1015
 Galois stable, 1015
 Galois type, 1001
 game, 79
 generalised product, 756
 κ -generated, 259, 969
 generated substructure, 154
 generated, finitely —, 154
 generating, 41
 generating a sequence by a type, 1164

generating an ideal, 404
 generator, 154, 745
 geometric dimension function, 1042
 geometric independence relation, 1089
 geometry, 1041
 global type, 1119
 graduated theory, 702, 787
 graph, 39
 greatest element, 42
 greatest fixed point, 661
 greatest lower bound, 42
 greatest partial fixed point, 663
 group, 34, 389, 460
 group action, 394
 group, ordered —, 710
 guard, 451

 Hanf number, 622, 642, 1007
 Hanf's Theorem, 610
 Hausdorff space, 355
 having κ -directed colimits, 257
 height, 192
 height in a lattice, 218
 Henkin property, 862
 Henkin set, 862
 Herbrand model, 515, 862
 hereditary, 12
 κ -hereditary, 914, 969
 hereditary finite, 7
 Hintikka formula, 590, 591
 Hintikka set, 517, 862, 863
 history, 15
 hom-functor, 263
 homeomorphism, 350
 homogeneous, 791, 929
 \approx -homogeneous, 936

κ -homogeneous, 608, 791
 homogeneous matroid, 1048
 homomorphic image, 157, 749
 homomorphism, 156, 498
 Homomorphism Theorem, 183
 homotopic interpretations, 894
 honest definition, 1163
 Horn formula, 740

 ideal, 205, 209, 404
 idempotent link, 317
 idempotent morphism, 317
 identity, 163
 image, 31
 imaginaries
 uniform elimination of —, 844
 imaginaries, elimination of —, 845
 imaginary elements, 830
 implication, 451
 implicit definition, 652
 inclusion functor, 169
 inclusion link, 281
 inclusion morphism, 495
 inconsistent, 458
 k -inconsistent, 1131
 increasing, 44
 independence property, 956
 independence relation, 1088
 independence relation of a matroid, 1087
 Independence Theorem, 1259
 independent, 1035
 $\sqrt{\quad}$ -independent family, 1295
 independent set, 110, 1041
 index map of a link, 280
 index of a subgroup, 390
 indiscernible sequence, 946

- indiscernible system, 953, 1343
- induced substructure, 152
- inductive, 77
- inductive completion, 295
- inductive completion of a category, 285
- inductive fixed point, 81, 661, 662
- inductively ordered, 81, 105
- infimum, 42, 197
- infinitary first-order logic, 449
- infinitary second-order logic, 487
- infinite, 115
- Infinity, Axiom of —, 24, 462
- inflationary, 81
- inflationary fixed-point logic, 669
- initial object, 166
- initial segment, 41
- injective, 31
- κ -injective structure, 1012
- inner vertex, 191
- insertion, 39
- inspired by, 954
- integral domain, 415, 717
- interior, 347, 762
- interpolant, 657
- interpolation closure, 653
- interpolation property, 651
- Δ -interpolation property, 651
- interpretation, 448, 450, 479
- intersection, 11
- intersection number, 1170
- interval, 761
- invariance, 1101
- invariant class, 1261
- invariant over a subset, 1331
- U -invariant relation, 1179
- invariant type, 1103
- inverse, 30, 165
- inverse diagram, 260
- inverse limit, 260
- inverse reduct, 979
- irreducible polynomial, 420
- irreflexive, 40
- $\sqrt{\text{—}}$ -isolated, 1303
- isolated point, 368
- isolated type, 859, 1103
- isolation relation, 1303
- isomorphic, 44
- α -isomorphic, 585, 596
- isomorphic copy, 749
- isomorphism, 44, 156, 165, 172, 498
- isomorphism, partial —, 581
- joint embedding property, 1009
- κ -joint embedding property, 914
- Jónsson class, 1009
- Karp property, 617
- kernel, 158
- kernel of a ring homomorphism, 406
- label, 231
- large subsets, 829
- Lascar invariant type, 1184
- Lascar strong type, 1174
- lattice, 197, 459, 494
- leaf, 191
- least element, 42
- least fixed point, 661
- least fixed-point logic, 668
- least partial fixed point, 662
- least upper bound, 42
- left extension, 1234
- left ideal, 404
- left local, 1114
- left reflexivity, 1088
- left restriction, 31
- left transitivity, 1088
- left-narrow, 57
- length, 189
- level, 192
- level embedding function, 935
- levels of a tuple, 935
- lexicographic order, 189, 1029
- lifting functions, 659
- limit, 59, 261
- limit stage, 19
- limiting cocone, 257
- limiting cone, 261
- Lindenbaum algebra, 493
- Lindenbaum functor, 492
- Lindström quantifier, 486
- linear independence, 410
- linear matroid, 1041
- linear order, 40
- linear representation, 691
- link between diagrams, 280
- literal, 449
- local, 612
- local character, 1113
- local enumeration, 776
- κ -local functor, 969
- local independence relation, 1114
- localisation morphism, 495
- localisation of a logic, 495
- locality, 1114
- locality cardinal, 1312
- locally compact, 356
- locally finite matroid, 1048
- locally modular matroid, 1048
- logic, 448
- logical system, 489
- Łoś' theorem, 721
- Łoś-Tarski Theorem, 690
- Löwenheim number, 622, 642, 645, 999
- Löwenheim-Skolem property, 617
- Löwenheim-Skolem-Tarski Theorem, 524
- lower bound, 42
- lower fixed-point induction, 662
- map, 29
- Δ -map, 497
- map, elementary —, 497
- mapping, 29
- matroid, 1040
- maximal element, 42
- maximal ideal, 415
- maximal ideal/filter, 205
- maximally φ -alternating sequence, 1159
- meagre, 366
- membership relation, 5
- minimal, 13, 57
- minimal element, 42
- minimal polynomial, 423
- minimal rank and degree, 226
- minimal set, 1053
- model, 448
- model companion, 703
- model of a presentation, 745
- model-complete, 703
- κ -model-homogeneous structure, 1012
- modular, 200
- modular lattice, 218
- modular law, 220, 221

- modular matroid, 1048
- modularity, 1099
- module, 407
- monadic second-order logic, 487
- monoid, 31, 191, 389
- monomorphism, 165
- monotone, 762
- monotonicity, 1088
- monster model, 829
- Morley degree, 1080
- Morley rank, 1077
- Morley sequence, 1123
- Morley-free extension of a type, 1080
- morphism, 162
- morphism of logics, 482
- morphism of matroids, 1048
- morphism of partial morphisms, 898
- morphism of permutation groups, 889
- multiplication of cardinals, 116
- multiplication of ordinals, 89
- multiplicity of a type, 1285
- mutually indiscernible sequences, 1213

- natural isomorphism, 172
- natural transformation, 172
- negation, 449, 493
- negation normal form, 473
- negative occurrence, 668
- neighbourhood, 345
- neutral element, 31
- node, 191
- normal subgroup, 391
- normality, 1088
- nowhere dense, 366

- o-minimal, 764, 960
- object, 162
- occurrence number, 622
- oligomorphic, 394, 881
- omitting a type, 532
- omitting types, 536
- open base, 348
- open cover, 356
- open dense order, 459
- open interval, 761
- Open Mapping Theorem, 1282
- open set, 345
- open subbase, 349
- opposite category, 166
- opposite functor, 168
- opposite lattice, 206
- opposite order, 40
- orbit, 394
- order, 458
- order property, 571
- order topology, 353, 762
- order type, 64, 945
- orderable ring, 430
- ordered group, 710
- ordered pair, 27
- ordered ring, 429
- ordinal, 64
- ordinal addition, 89
- ordinal exponentiation, 89
- ordinal multiplication, 89
- ordinal, von Neumann —, 69

- pair, 27
- parameter equivalence, 835
- parameter-definable, 763
- partial fixed point, 662
- partial fixed-point logic, 669

- partial function, 29
- partial isomorphism, 581
- partial isomorphism modulo a filter, 732
- partial morphism, 898
- partial order, 40, 458
- partial order, strict —, 40
- partition, 55, 222
- partition degree, 226
- partition rank, 222
- partitioning a relation, 780
- path, 191
- path, alternating — in a category, 276
- Peano Axioms, 488
- pinning down, 622
- point, 345
- polynomial, 403
- polynomial function, 420
- polynomial ring, 403
- positive existential, 498
- positive occurrence, 668
- positive primitive, 740
- power set, 21
- predicate, 28
- predicate logic, 448
- prefix, 189
- prefix order, 189
- preforking relation, 1101
- prelattice, 209
- prenex normal form, 473
- preorder, 208, 492
- κ -presentable, 321
- presentation, 745
- preservation by a function, 497
- preservation in products, 740
- preservation in substructures, 500
- preservation in unions of chains, 501

- preserving a property, 168, 266
- preserving fixed points, 659
- $\sqrt{\kappa}$ -prime, 1320
- prime field, 417
- prime ideal, 209, 406
- prime model, 872
- prime model, algebraic, 698
- primitive formula, 703
- principal ideal/filter, 205
- Principle of Transfinite Recursion, 75, 133
- product, 27, 37, 749
- product of categories, 170
- product of linear orders, 86
- product topology, 361
- product, direct —, 243
- product, generalised —, 756
- product, reduced —, 246
- product, subdirect —, 244
- projection, 37, 640
- projection along a functor, 265
- projection along a link, 281
- projection functor, 170
- projective class, 640
- projective geometry, 1047
- projectively reducible, 641
- projectively κ -saturated, 808
- proper, 205
- property of Baire, 367
- pseudo-elementary, 641
- pseudo-saturated, 811

- quantifier elimination, 694, 716
- quantifier rank, 457
- quantifier-free, 457
- quantifier-free formula, 498
- quantifier-free representation, 1344

- quasi-dividing, 1237
- quasivariety, 749
- quotient, 180

- Rado graph, 922
- Ramsey's theorem, 930
- random graph, 922
- random theory, 922
- range, 29
- rank, 73, 194
- Δ -rank, 1077
- rank, foundation \neg , 194
- real closed field, 433, 715
- real closure of a field, 433
- real field, 430
- realising a type, 532
- reduced product, 246, 749
- reduct, 155
- μ -reduct, 241
- refinement of a partition, 1342
- reflecting a property, 168, 266
- reflexive, 40
- regular, 125
- regular filter, 723
- regular logic, 618
- relation, 28
- relational, 149
- relational variant of a structure, 980
- relativisation, 478, 618
- relativised projective class, 645
- relativised projectively reducible, 645
- relativised quantifiers, 451
- relativised reduct, 645
- Replacement, Axiom of \neg , 133, 462
- replica functor, 983
- representation, 1344
- restriction, 30
- restriction of a filter, 246
- restriction of a Galois type, 1019
- restriction of a logic, 495
- restriction of a type, 564
- retract of a logic, 551
- retraction, 165
- retraction of logics, 550
- reverse ultrapower, 739
- right local, 1114
- right shift, 1303
- ring, 401, 461
- ring, orderable \neg , 430
- ring, ordered \neg , 429
- root, 191
- root of a polynomial, 420
- Ryll-Nardzewski Theorem, 881

- satisfaction, 448
- satisfaction relation, 448, 450
- satisfiable, 458
- saturated, 797
- κ -saturated, 672, 797
- $\sqrt{\quad}$ - κ -saturated, 1320
- κ -saturated, projectively \neg , 808
- Scott height, 591
- Scott sentence, 591
- second-order logic, 487
- section, 165
- segment, 41
- semantics functor, 489
- semantics of first-order logic, 450
- semi-strict homomorphism, 156
- semilattice, 197
- sentence, 454
- separated formulae, 632
- Separation, Axiom of \neg , 10, 462
- sequence, 37
- shifting a diagram, 317
- signature, 149, 151, 239, 240
- simple structure, 416
- simple theory, 1141
- simply closed, 698
- singular, 125
- size of a diagram, 255
- skeleton of a category, 270
- skew embedding, 943
- skew field, 401
- Skolem axiom, 509
- Skolem expansion, 1003
- Skolem function, 509
- definable \neg , 847
- Skolem theory, 509
- Skolemisation, 509
- small subsets, 829
- sort, 151
- spanning, 1038
- special model, 811
- specification of a dividing chain, 1143
- specification of a dividing κ -tree, 1150
- specification of a forking chain, 1143
- spectrum, 374, 535, 538
- spectrum of a ring, 406
- spine, 985
- splitting type, 1103
- stabiliser, 395
- stability spectrum, 1296
- κ -stable formula, 568
- κ -stable theory, 577
- stably embedded set, 1162
- stage, 15, 77
- stage comparison relation, 679
- stationary set, 138
- stationary type, 1278
- Stone space, 378, 535, 538
- $\sqrt{\quad}$ -stratification, 1312
- strict homomorphism, 156
- strict Horn formula, 740
- strict Δ -map, 497
- strict order property, 962
- strict partial order, 40
- strictly increasing, 44
- strictly monotone, 762
- strong γ -chain, 1021
- strong γ -limit, 1021
- strong finite character, 1116
- strong limit cardinal, 812
- strong right boundedness, 1089
- strongly homogeneous, 791
- strongly κ -homogeneous, 791
- strongly independent, 1338
- strongly local functor, 985
- strongly minimal set, 1053
- strongly minimal theory, 1060, 1156
- structure, 149, 151, 241
- subbase, closed \neg , 348
- subbase, open \neg , 349
- subcategory, 169
- subcover, 356
- subdirect product, 244
- subdirectly irreducible, 244
- subfield, 417
- subformula, 454
- subset, 5
- subspace topology, 350
- subspace, closure \neg , 350
- substitution, 238, 469, 618
- substructure, 152, 749, 969
- Δ -substructure, 502
- \mathcal{K} -substructure, 1000
- substructure, elementary \neg , 502
- substructure, generated \neg , 154

substructure, induced —, 152
 subterm, 232
 subtree, 192
 successor, 59, 191
 successor stage, 19
 sum of linear orders, 85
 superset, 5
 supersimple theory, 1300
 superstable theory, 1300
 supremum, 42, 197
 surjective, 31
 symbol, 149
 symmetric, 40
 symmetric group, 393
 symmetric independence relation,
 1089
 syntax functor, 489
 system of bases for a stratification,
 1342

 T_0 -space, 538
 Tarski union property, 618
 tautology, 458
 term, 231
 term algebra, 236
 term domain, 231
 term, value of a —, 235
 term-reduced, 470
 terminal object, 166
 L -theory, 465
 theory of a functor, 975
 topological closure, 347, 762
 topological closure operator, 51, 347
 topological group, 398
 topological space, 345
 topology, 345
 topology of the type space, 537

torsion element, 709
 torsion-free, 709
 total order, 40
 totally disconnected, 355
 totally indiscernible sequence, 946
 totally transcendental theory, 578
 transcendence basis, 422
 transcendence degree, 422
 transcendental elements, 422
 transcendental field extensions, 422
 transfinite recursion, 75, 133
 transitive, 12, 40
 transitive action, 394
 transitive closure, 55
 transitive dependence relation, 1035
 transitivity, left —, 1088
 translation by a functor, 265
 tree, 191
 φ -tree, 572
 tree property, 1149
 tree property of the second kind, 1227
 tree-indiscernible, 954
 trivial filter, 205
 trivial ideal, 205
 trivial topology, 346
 tuple, 28
 Tychonoff, Theorem of —, 363
 type, 564
 L -type, 531
 E -type, 808
 α -type, 532
 \bar{s} -type, 532
 type of a function, 151
 type of a relation, 151
 type space, 537
 type topology, 537
 type, average —, 947

type, average — of an indiscernible
 system, 954
 type, complete —, 531
 type, Lascar strong —, 1174
 types of dense linear orders, 533

 ultrafilter, 209, 534
 κ -ultrahomogeneous, 910
 ultrapower, 247
 ultraproduct, 247, 801
 unbounded class, 1007
 uncountable, 115
 uniform dividing chain, 1143
 uniform dividing κ -tree, 1151
 uniform elimination of imaginaries,
 844
 uniform forking chain, 1143
 uniformly finite, being — over a set,
 780
 union, 21
 union of a chain, 505, 692
 union of a cocone, 297
 union of a diagram, 296
 unit of a ring, 415
 universal, 498
 κ -universal, 797
 universal quantifier, 449
 universal structure, 1012
 universe, 149, 151
 unsatisfiable, 458
 unstable, 568, 578
 upper bound, 42
 upper fixed-point induction, 662

 valid, 458

value of a term, 235
 variable, 240
 variable symbols, 449
 variables, free —, 235, 454
 variety, 749
 Vaughtian pair, 1061
 vector space, 407
 vertex, 191
 von Neumann ordinal, 69

 weak γ -chain, 1021
 weak γ -limit, 1021
 weak canonical definition, 851
 weak canonical parameter, 851
 weak elimination of imaginaries, 851
 weak homomorphic image, 157, 749
 Weak Independence Theorem, 1258
 weakly bounded independence
 relation, 1195
 weakly regular logic, 618
 well-founded, 13, 57, 81, 109
 well-order, 57, 109, 133, 602
 well-ordering number, 622, 642
 well-ordering quantifier, 486, 487
 winning strategy, 594
 word construction, 976, 981

 Zariski logic, 447
 Zariski topology, 346
 zero-dimensional, 355
 zero-divisor, 415
 Zero-One Law, 926
 ZFC, 461
 Zorn's Lemma, 110

The Roman and Fraktur alphabets							
<i>A</i>	<i>a</i>	𝐀	𝐚	<i>N</i>	<i>n</i>	𝐸	<i>n</i>
<i>B</i>	<i>b</i>	𝐁	𝐛	<i>O</i>	<i>o</i>	𝐎	<i>o</i>
<i>C</i>	<i>c</i>	𝐶	𝐜	<i>P</i>	<i>p</i>	𝐏	<i>p</i>
<i>D</i>	<i>d</i>	𝐷	𝐝	<i>Q</i>	<i>q</i>	𝐐	<i>q</i>
<i>E</i>	<i>e</i>	𝐸	𝐞	<i>R</i>	<i>r</i>	𝐫	<i>r</i>
<i>F</i>	<i>f</i>	𝐅	𝐟	<i>S</i>	<i>s</i>	𝐒	<i>f</i> <i>ſ</i>
<i>G</i>	<i>g</i>	𝐆	𝐠	<i>T</i>	<i>t</i>	𝐓	<i>t</i>
<i>H</i>	<i>h</i>	𝐇	𝐇	<i>U</i>	<i>u</i>	𝐔	<i>u</i>
<i>I</i>	<i>i</i>	𝐈	𝐢	<i>V</i>	<i>v</i>	𝐕	<i>v</i>
<i>J</i>	<i>j</i>	𝐉	𝐣	<i>W</i>	<i>w</i>	𝐖	<i>w</i>
<i>K</i>	<i>k</i>	𝐊	𝐤	<i>X</i>	<i>x</i>	𝐗	<i>x</i>
<i>L</i>	<i>l</i>	𝐋	𝐥	<i>Y</i>	<i>y</i>	𝐘	<i>y</i>
<i>M</i>	<i>m</i>	𝐌	𝐦	<i>Z</i>	<i>z</i>	𝐙	<i>z</i>

The Greek alphabet					
<i>A</i>	α	alpha	<i>N</i>	ν	nu
<i>B</i>	β	beta	<i>Ξ</i>	ξ	xi
<i>Γ</i>	γ	gamma	<i>Ο</i>	o	omicron
<i>Δ</i>	δ	delta	<i>Π</i>	π	pi
<i>E</i>	ϵ	epsilon	<i>Ρ</i>	ρ	rho
<i>Z</i>	ζ	zeta	<i>Σ</i>	σ	sigma
<i>H</i>	η	eta	<i>Τ</i>	τ	tau
<i>Θ</i>	θ	theta	<i>Υ</i>	υ	upsilon
<i>I</i>	ι	iota	<i>Φ</i>	ϕ	phi
<i>K</i>	κ	kappa	<i>Χ</i>	χ	chi
<i>Λ</i>	λ	lambda	<i>Ψ</i>	ψ	psi
<i>M</i>	μ	mu	<i>Ω</i>	ω	omega